



000352-00541-060499

Printed: 10:56:29 Friday, 25 Jun, 1999 # 52 / 3503

**Systems Integration Group, Inc.**



PM 25

352-541

6/4/99  
SUPPLEMENTAL LABELING

163

DuPont Agricultural Products

"..... A Growing Partnership With Nature"



**ASSURE® II HERBICIDE**  
FOR POSTEMERGENCE  
GRASS CONTROL IN CERTAIN NON  
FOOD/NON FEED CROPS GROWN UNDER  
CONTRACT FOR SEED PRODUCTION ONLY  
IN ID, MT, OR, WA, AND WY

## ASSURE® II HERBICIDE

EPA REG. NO. 352-541

**FOR POSTEMERGENCE GRASS CONTROL IN CERTAIN NON FOOD/NON FEED CROPS  
GROWN UNDER CONTRACT FOR SEED PRODUCTION ONLY  
IN THE STATES OF IDAHO, MONTANA, OREGON, WASHINGTON, AND WYOMING**

### DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

DuPont ASSURE® II Herbicide is a selective postemergence herbicide recommended for control of annual and perennial grasses in alfalfa, onion, carrot, garlic, Swiss chard, spinach, radish, Chinese cabbage, and red beets grown specifically under contract as non food/non feed crops for seed production only. See "Restrictions" portion of label before using. Applied at recommended rates and timings, ASSURE® II Herbicide will control emergent grasses. Subsequent flushes of grasses require additional treatment.

### HOW TO USE

#### Ground Application

#### Broadcast Application

- Use flat fan or hollow cone nozzles at 25-60 psi.
- Do not use flood, rain drop, whirl chamber, or any other nozzle types that produce coarse, large spray droplets. In addition, do not use controlled droplet applicator (CDA) type nozzles as poor weed control or excessive spray drift may result.
- Use a minimum of 10 gal of water per acre
- Increase spray volume and pressure as weed or crop density and size increase.
- Do not exceed 40 gal of water per acre or control will be reduced.
- Adjust the boom and nozzle height according to the nozzle manufacturer's specifications to obtain proper spray coverage.

#### Aerial Application

- Use nozzle types and arrangements that provide optimum spray distribution and maximum coverage.
- Use a minimum of 3 gal of water per acre.
- Do not apply during a temperature inversion, when winds are gusty, or when other conditions favor poor coverage and/or off-target spray movement.

See full label for spray drift management and sprayer cleanout directions.

### Spray Additives

Always include a nonphytotoxic petroleum based crop oil concentrate at 1% v/v (4 qts/100 gals) or a nonionic surfactant at 0.25% v/v (1 qt/100 gals). Crop oil concentrate is the preferred adjuvant in arid areas.

### Tank Mix Applications


Tank mixtures of ASSURE® II with any pesticide or spray adjuvant is not recommended except as directed on this label or on other supplemental labels.

Tank mixes of ASSURE® II with postemergence broadleaf herbicides may result in reduced grass control. If grass control is reduced, an additional application of ASSURE® II may be required after grass plants begin to develop new leaves.

### Sequential Applications with Post Broadleaf Herbicides

**NOTE:** Reduction in grass control is possible when ASSURE® II is applied immediately prior to or sequentially after an application of a post broadleaf herbicide. Observe the following recommendations:

- After applying ASSURE® II, wait a minimum of 24 hours before applying a post broadleaf herbicide.
- In fields treated with a post broadleaf herbicide, reduced control may result if applications of ASSURE® II are made prior to grass plants beginning to develop new leaves (generally 5-7 days after the post broadleaf herbicide application).



WEEDS CONTROLLED AND RATE SELECTION		
	Size at Application (in)	ASSURE® II Applied Alone (oz product/A)
<b>Annual Grasses**</b>		
Corn, Volunteer ( <i>Zea mays</i> )	6-18	5 - 8 oz.
Foxtail, Giant ( <i>Setaria faberi</i> )	2-4 (pretiller)	
Johnsongrass, Seedling ( <i>Sorghum halepense</i> )	2-8	
Shattercane ( <i>Sorghum bicolor</i> )	6-12	
Wild Proso Millet ( <i>Panicum miliaceum</i> )	2-6	
Crowfootgrass ( <i>Dactyloctenium aegyptium</i> )	2-6	7 - 8 oz.
Fall Panicum ( <i>Panicum dichotomiflorum</i> )	2-6	
Field Sandbur ( <i>Cenchrus incertus</i> )	2-6	
Foxtail, Bristly ( <i>Setaria verticillata</i> )	2-4	
Foxtail, Giant ( <i>Setaria faberi</i> )	2-8	
Foxtail, Green ( <i>Setaria viridis</i> )	2-4	
Foxtail, Yellow ( <i>Setaria lutescens</i> )	2-4	
Goosegrass ( <i>Eleusine indica</i> )	2-6‡	
Itchgrass ( <i>Rottboellia exaltata</i> )	2-8	
Sprangletop ( <i>Leptochloa filiformis</i> )	2-6	
Volunteer Barley ( <i>Hordeum vulgare</i> )	2-6	
Volunteer Oats ( <i>Avena sativa</i> )	2-6	
Volunteer Rye ( <i>Secale cereale</i> )	2-6	
Volunteer Wheat ( <i>Triticum aestivum</i> )	2-6	
Wild Oat ( <i>Avena fatua</i> )	2-6	
Witchgrass ( <i>Panicum capillare</i> )	2-6	
Barnyardgrass ( <i>Echinochloa crus-galli</i> )	2-6	8 - 10 oz.
Crabgrass, Large ( <i>Digitaria sanguinalis</i> )	2-6‡	
Crabgrass, Smooth ( <i>Digitaria ischaemum</i> )	2-6‡	
Junglerice ( <i>Echinochloa colonum</i> )	2-6	
Texas Panicum ( <i>Panicum texanum</i> )	2-4	
Red Rice ( <i>Oryza sativa</i> )	1-4	9 - 10 oz.
Woolly Cupgrass ( <i>Eriochloa villosa</i> )	2-4§	10 oz.
Broadleaf Signalgrass ( <i>Brachiaria platyphylla</i> )	2-6	
<b>Perennial Grasses**</b>		
Wirestem Muhly ( <i>Muhlenbergia frondosa</i> )	4-8	8 - 10 oz.
Bermudagrass ( <i>Cynodon dactylon</i> )	3" tall (or up to 6" runners)	10 - 12 oz.
Johnsongrass, Rhizome ( <i>Sorghum halepense</i> )	10-24	
Quackgrass ( <i>Agropyron repens</i> )	6-10	
<p>** For annual and perennial grasses, up to 12 oz per acre may be applied, based on local recommendations. Under arid conditions the higher use rate is recommended.</p> <p>‡ Length of lateral growth.</p> <p>§ Size in height or diameter, whichever is more restrictive. Applications to plants with more than three tillers may result in unsatisfactory control.</p>		

**Precautions**

- Rainfall within 1 hour of application will reduce grass control from ASSURE® II.
- Applications to grassy weeds suffering stress from lack of moisture, cold, herbicide injury, and insect or disease injury may result in reduced control. A sequential application of ASSURE II at 6-7 oz per acre after growth resumes may be necessary for satisfactory control.

**Restrictions**

- Do not apply Assure II within 14 days of anticipated bloom.
- The maximum use rate of ASSURE® II is 25 oz per acre per season.
- After using ASSURE® II, do not divert any portion of crop (seed, sprouts, screenings, forage, hay, etc.) to use for human or animal consumption. Grazing of treated crop is prohibited.
- Do not make more than 2 applications per acre per season. Application intervals should be greater than 7 days apart to allow regrowth to occur.
- Do not apply ASSURE® II through any type of irrigation system.
- Most grass crops, including wheat, barley, rye, oats, sorghum, rice, and corn are highly sensitive to ASSURE® II Herbicide and all direct or indirect contact (such as spray drift) should be avoided.
- All seed crops treated with "Assure" II are to be tagged at the processing facility, "Not for Human or Animal Consumption". It shall be the growers' responsibility to notify the processing facility of any seed crop that has been treated with ASSURE® II.

**Resistance**

Biotypes of certain weeds listed on this label are resistant to Assure II, and other herbicides with the same mode of action,\* even at exaggerated application rates. Biotypes are naturally occurring individuals of a species identical in appearance but with slightly different genetic compositions; the mode of action of a herbicide is the chemical interaction that interrupts a biological process necessary for plant growth and development.

If weed control is unsatisfactory, it may be necessary to respray problem areas using a product with a different mode of action. If resistant weed biotypes (such as Wild Oats), are suspected or known to be present, consider using a planned herbicide rotation program to help control these biotypes. To better manage weed resistance when using ASSURE II use a combination of tillage and sequential herbicide applications that have a different mode of action than ASSURE II, to control escaped weeds. Do not let weed escapes go to seed.

Consult your agricultural dealer, consultant, applicator, and/or appropriate state agricultural extension service representative for specific alternative herbicide recommendations available in your area. It is advisable to keep accurate records of pesticides applied to individual fields to help obtain information on the spread and dispersal of resistant biotypes.

\* Naturally occurring weed biotypes that are resistant to Hoelon or Poast, will also be resistant to Assure II.

**IMPORTANT**

**BEFORE USING "ASSURE" II, READ AND FOLLOW ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA-REGISTERED LABEL.**

This bulletin contains new or supplemental instructions for use of this product which do not appear on the EPA-registered package label. Follow the instructions carefully.

This labeling must be in the possession of the user at the time of pesticide application.

D-12159