



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

MAY 31 1994

OFFICE OF
PREVENTION, PESTICIDES AND
TOXIC SUBSTANCES

Nancy B. Lomax
E.I. DU PONT DE NEMOURS AND CO., INC.
Barley Mill Plaza, Walker's Mill
Wilmington, DE 19880

Subject: Label Amendment Submission of 05/20/93 in Response to PR Notice 93-7
EPA Reg. No. 352-544
DUPONT NEW GEMINI HERBICIDE

Dear Registrant:

The labeling cited above and submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, is accepted subject to the comments reflected on the enclosed sheet. A copy of your proposed labeling stamped "ACCEPTED WITH COMMENTS" is enclosed.

WHAT THIS ACCEPTANCE MEANS:

Based on your certification, the Agency has accepted the labeling changes that are necessary to comply with the Worker Protection Standard (WPS) labeling requirements of 40 CFR part 156, subpart K, described in PR Notices 93-7 and 93-11. Any other labeling changes submitted in connection with this amendment application but not directly related to compliance with the WPS have not been reviewed or accepted by the Agency. If you wish to make such changes, you must submit a separate amendment application proposing them. If your product is currently suspended, the acceptance of this labeling amendment does not affect the suspension in any way.

WHAT YOU NEED TO DO NEXT:

By the next label printing make all the specified changes to your labeling. Send to EPA one (1) copy of the final printed labeling:

- BEFORE selling or distributing any product bearing the final printed labeling
- AND
- WITHIN one year from date of this acceptance.



Recycled/Recyclable
Printed with Soy/Canola Ink on paper that
contains at least 50% recycled fiber


Submit the final printed labeling via the U.S. Postal Service to:

Document Processing Desk (FIN-LABEL)
Office of Pesticide Programs (7505C)
U.S. Environmental Protection Agency
401 M Street, SW
Washington, D.C. 20460-0001

Hand or courier deliveries of final printed labeling may be made to:

Document Processing Desk (FIN-LABEL)
Office of Pesticide Programs
Room 266A, Crystal Mall 2
1921 Jefferson Davis Highway
Arlington, VA 22202

Sincerely,



Jim Tompkins, Deputy Chief
Registration Support Branch
Registration Division (7505W)

Attachment

119

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
Office of Pesticide Programs
Registration Division

Nancy B. Lomax
E. I. DU PONT DE NEMOURS AND CO, INC.
BARLEY MILL PLAZA, WALKER'S MILL
WILMINGTON DE 19880

Comment for: EPA Reg Nr.352-544
DUPONT NEW GEMINI HERBICIDE

The following specific comments pertain to your WPS labeling submission concerning the product cited above:

User Safety Recommendations must either be placed in a box or printed on the label in a contrasting color from surrounding text.

Add the following User Safety Recommendation to the label in the location shown in Section A on Part 1 of the Product Worksheet in Supplement Three-A to PR Notice 93-7: "Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing."

Correct the typographical errors circled on your proposed label.



4,59
GEMINI®

HERBICIDE

DISPERSIBLE GRANULES

BY WEIGHT

ACTIVE INGREDIENTS 60%

Linuron: [3-(3,4-dichlorophenyl)-1-methoxy-1-methylurea] 55.4%

Chlorimuron Ethyl: Ethyl 2-[[[(4-chloro-6-methoxypyrimidin-2-yl)amino]-carbonyl]amino]sulfonyl]benzoate 4.6%

INERT INGREDIENTS 40%

TOTAL 100%

EPA Reg. No. 352-544

U.S. Pat. 4,394,506 & 4,547,215

KEEP OUT OF REACH OF CHILDREN

CAUTION

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION! Harmful if swallowed or absorbed through skin. Causes moderate eye injury. Avoid contact with skin, eyes and clothing.

STATEMENT OF PRACTICAL TREATMENT

In case of eye contact, flush with plenty of water and get medical attention if irritation persists.

In case of skin contact, wash with soap and water and get medical attention if irritation persists. Wash thoroughly with soap and water after handling. Remove and wash contaminated clothing before reuse.

If swallowed, call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person.

For medical emergencies involving this product, call toll free 1-800-441-3637.

PERSONAL PROTECTIVE EQUIPMENT

Applicators and other handlers must wear:

- Coveralls over short-sleeve shirt and short pants.
- Waterproof gloves.
- Shoes plus socks.
- Chemical-resistant headgear for overhead exposure.

ACCEPTED
with COMMENTS
in EPA Letter Dated

MAR 31 1984

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

352-544

Discard clothing or other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENGINEERING CONTROLS STATEMENTS

When handlers use closed systems or enclosed cabs in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR part 170 Section 170.240 (d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

USERS SHOULD: Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.

ENVIRONMENTAL HAZARDS

Do not apply directly to water or wetlands (swamps, bogs, marshes and potholes). Do not contaminate water by cleaning of equipment or disposal of wastes.

BEST AVAILABLE COPY

IMPORTANT

Injury to or loss of desirable trees or vegetation may result from failure to observe the following: Do not apply or drain or flush equipment on or near desirable trees or other plants, or on areas where their roots may extend, or in locations where the chemical may be washed or moved into contact with their roots. Do not use on lawns, walks, driveways, tennis courts, or similar areas. Prevent drift of spray to desirable plants. Do not contaminate any body of water. Keep from contact with fertilizers, insecticides, fungicides, and seeds during storage.

PRIOR TO USING "GEMINI", CONSIDERATION SHOULD BE GIVEN TO CROP ROTATION PLANS.

Crops other than soybeans can be extremely sensitive to low concentrations of "Gemini" remaining in the soil the next planting season. Choice of rotational crop is restricted following application of "Gemini". (See "ROTATIONAL CROP GUIDELINES".)

Thoroughly clean all traces of "Gemini" from application equipment immediately after use and prior to spraying crops other than soybeans. Failure to remove even small amounts of "Gemini" from application equipment may result in injury to subsequently sprayed crops.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls over short-sleeve shirt and short pants.
- Waterproof gloves.
- Shoes plus socks.
- Chemical-resistant headgear for overhead exposure.

549
Do not apply this product through any type of irrigation system.

FOR USE ON SOYBEANS ONLY.

GENERAL INFORMATION

Du Pont "Gemini" Herbicide is a dispersible granule formulation to be mixed with water and sprayed for selective preemergence weed control in soybeans. When applied according to the instructions on this label, it will control many broadleaf weeds and provide partial control of nutsedge and annual grasses.

Preemergence and preplant incorporated applications of "Gemini" require rainfall or sprinkler irrigation to activate the herbicide. Degree of control and duration of effect depend on: rate used, weed spectrum, growing conditions at and following time of treatment, soil pH, texture, organic matter, moisture, and precipitation.

Because most crops are highly sensitive to "Gemini", all direct or indirect contact (such as spray drift) to crops or to land scheduled to be planted to crops other than soybeans should be avoided.

BIOLOGICAL ACTIVITY

"Gemini" rapidly inhibits the growth of susceptible weeds. Following activation of preplant incorporated or preemergence treatment, susceptible weeds may germinate and emerge, but growth then ceases and leaves become yellow 3-5 days after emergence. Death of leaf tissue and growing point will follow in some species while others remain green but stunted and noncompetitive. "Gemini" provides partial control of some annual grasses when used preplant or preemergence but other products may be needed to ensure adequate grass control.

WEEDS CONTROLLED

When used as directed "Gemini" will provide control of the following weeds:

Carpetweed	Pigweed species
Cocklebur	Palmer
Florida beggarweed	Redroot
Jimsonweed	Smooth
Lambsquarters	Spiny amaranth
*Morningglory, annual	Tall waterhemp
Ivyleaf	Prickly sida (teaweed)
Entireleaf	Purslane
Pitted	Ragweed, common
Smallflower	*Ragweed, giant
Tall	Smartweed
Mustards	Sunflower
	Velvetleaf

When used as directed "Gemini" will provide partial control of the following weeds:

Annual grasses	Sicklepod*
Burcucumber	Wild poinsettia
Hemp sesbania	Yellow nutsedge
Purple nutsedge	

- * Large seeded weeds germinating deep in the soil such as morning-glory and giant ragweed may require a cultivation or an application of Du Pont "Classic" Herbicide postemergence for season long

BEST AVAILABLE COPY

control. Consult "Classic" label for application directions, recropping intervals and use restrictions.

NOTE: Consult Lasso', Dual', Prowl', Sonalan', Surflan', or Treflan' labels for additional weeds controlled when "Gemini" is used in conjunction with these herbicides.

APPLICATION INFORMATION

SPRAY TANK PREPARATION

It is important that spray equipment is clean and free of existing pesticide deposits before using "Gemini". Follow the spray tank cleanout procedures specified on the label of product previously sprayed. If no cleanout procedure is provided, follow the cleanout procedure below for all application equipment.

1. Thoroughly rinse sprayer, tanks, boom, and hoses with clean water.
2. Partially fill the tank with water and add one of the cleaning agents listed in the SPRAYER CLEANUP section of this label. Complete filling the tank and flush the cleaning solution through the boom and hoses. Let stand for 15 minutes with agitation or recirculation and then drain the tank after flushing the hoses, boom, and nozzles.
3. Thoroughly rinse sprayer, tanks, boom, and hoses with clean water.
4. Follow label directions of the product previously sprayed for rinsate disposal.

NOTE: During an extended period where spraying or mixing equipment will be used to apply multiple loads of "Gemini", at the end of each day of spraying partially fill the tank with fresh water, flush the boom and hoses and allow to sit overnight.

EQUIPMENT / SPRAY VOLUMES

Apply uniformly by ground equipment with a properly calibrated low pressure (20 to 40 psi) fixed-boom sprayer equipped with fan-type nozzles and screens no finer than 50 mesh. Use 10 to 40 gals. of water per acre.

Continuous agitation in the spray tank is required to keep the material in suspension. Avoid overlap, and shut off spray booms while starting, turning, sowing or stopping, or injury to the crop may result.

MIXING INSTRUCTIONS

Fill tank 1/4 full with water. Start agitation system, add "Gemini" and continue adding water. Add separately each additional component of any tank mix while adding water. Continue agitation throughout. If poor mixing should occur with any component, premix the component with two parts water before adding to the spray tank.

A fertilizer solution may be used in the spray mixture. Small quantities should be tested for compatibility by the following procedures before full scale mixing.

1. Put 1 pint of fertilizer solution in a quart jar.
2. Mix 2 teaspoons "Gemini" with 2 tablespoons of water; mix thoroughly and add to fertilizer solution.
3. Close jar and shake well.
4. If other herbicides are to be used in the mixture, premix 2 teaspoons of wettable powder or 1 teaspoon of liquid with

2 tablespoons of water; add to "Gemini"/fertilizer solution mixture.

5. Close jar and shake well.
6. Watch mixture for several seconds; check again in 30 minutes.
7. If mixture does not separate, foam, gel, or become lumpy, it may be used.
8. Mixing ability may be improved by adding compatibility agents such as Kalo Laboratories' Compex³ or Witco Chemicals' Sponto⁶ 168D. Follow directions on container.

Provided the above procedure shows the mixture to be compatible, prepare the tank mixture as follows: Add the fertilizer solution to the spray tank first, with the agitator running, add the required amount of "Gemini" and thoroughly mix. For tank mixtures with other herbicides, follow directions above.

When tank mixing with other herbicides, all applicable directions, restrictions and precautions for the additional herbicides are also to be followed. Use "Gemini" spray preparations the same day as mixed or product degradation may occur. Thoroughly reagituate before using if allowed to settle.

APPLICATION METHODS

If initial seeding fails to produce a stand, treated fields may be replanted to soybeans. Do not rework soil; do not retreat field with a second application as injury to the crop may result. Do not replant treated areas to any crop other than soybeans within four months after treatment as injury to subsequent crops may result.

EARLY PREPLANT OR PREPLANT INCORPORATED (SEE RATE CHART)

"Gemini" may be applied at planting or up to 30 days prior to planting either preemergence or preplant incorporated. If incorporated, "Gemini" should be uniformly incorporated no deeper than the top 1-2" of soil prior to planting soybeans. Use equipment suited for proper incorporation into the top 1-2 inches of soil e.g.: Do-all, field cultivator, Roterra. If tank-mixed with a grass herbicide, such as a DNA like "Treflan", "Sonalan", or acetanilide such as "Dual", or "Lasso", follow label instructions for proper incorporation of the grass herbicide into the top 1 to 2" of soil. Improper soil incorporation can result in erratic weed control or crop injury. Rainfall* (~1") is required to activate preplant incorporated herbicides such as "Gemini" before weed emergence.

Use of a disk for incorporation is not recommended.

"Gemini" + "Treflan", "Prowl", "Dual" or "Sonalan" - Apply within 2 weeks of planting. "Gemini" + "Lasso" - Apply within 7 days of planting.

PREEMERGENCE (SEE RATE CHART)

Preemergence application: Apply before soybeans emerge. Do not spray over top of emerged soybeans as severe injury or death of crop will occur.

Preemergence activity of "Gemini" requires rainfall* to activate. Best results are obtained if treatment is made to moist soil and followed by rainfall or irrigation (~1") before weeds germinate. On dry soil more moisture is required for

BEST AVAILABLE COPY

activation (1 to 2") before weed emergence. If moisture is insufficient to activate the herbicide, a rotary hoeing or shallow cultivation should be made after emergence of crop while weeds are small enough to be controlled by mechanical means. "Gemini" can be applied preemergence in combination with "Lasso", "Dual", "Prowl", "Surflan", or following the use of a preplant grass material such as "Treflan" or "Sonalan." Consult labels for rates and use instructions.

*Rainfall required to activate ppi or pre treatments depends on soil moisture when activating moisture is received. Several small rainfalls of less than 1/4" are not as beneficial as one large rain of 1/2 inch or more for activation.

**MINIMUM TILLAGE OR NO-TILL
(SEE RATE CHART)**

"Gemini" applied preemergence to soybeans but postemergence to broadleaf weeds and small annual grasses will provide some burndown of existing weeds as well as normal preemergence broadleaf control. When used for burndown add 0.25% (1 qt./100 gal) surfactant such as "Surfactant WK" or crop oil concentrate at 1% v/v (1 gal/100 gal). For added postemergence control especially on grasses, "Gemini" can be applied with Roundup', paraquat, or Bronco'. For added preemergence grass control "Gemini" can be mixed with "Dual", "Surflan" or "Lasso." Consult labels for rates and use instructions. Maintain constant agitation of spray mixture.

RATE CHART

Apply as follows:

Soil Texture (1/2 - 3% Organic Matter)	Rate (Ounces Per Acre)
Coarse:	
Loamy sand	12-16
Medium:	
Loam, Silty loam, Silt	16-20
Fine:	
Silty Clay loam, Clay loam, Clay	20-24

Do not use on sand or loamy sand nor on any soil with less than 1/2 percent organic matter.

Do not apply "Gemini" on soils with pH greater than 7.0 except in those states as directed in the "ROTATIONAL CROP GUIDELINES" where application can be made to soils with a pH up to 7.5.

**REDUCED RATE FOLLOWED BY
"CLASSIC" HERBICIDE**

"Gemini" Herbicide is recommended preplant incorporated or preemergence at reduced rates followed by "Classic" Herbicide postemergence for broadleaf weed control in soybeans. This treatment will control those broadleaf weeds listed on the "Gemini" and "Classic" Herbicide product labels.

Apply "Gemini" preplant incorporated or preemergence at 8-10 ounces product per acre to any soil texture with 1/2 to 3% organic matter as directed on the product label. Follow the "Gemini" preplant incorporated or preemergence reduced rate with a postemergence application of "Classic"

Herbicide at its labeled rate when weeds are small and actively growing. (Usually 14-28 days after preplant incorporated or preemergence application of "Gemini" reduced rate.)

For grass weed control "Gemini" can be tank mixed with Dual, Lasso, Sonalan, Treflan or Surflan. Consult labels for rates and use instructions of these products.

Dosage rates listed are for broadcast rates. For band applications use proportionately less.

Plant seed 1 1/2 to 2" deep on a flat or raised seedbed only, or crop injury may occur.

Soybean stunting may occur if excessive rainfall occurs after application. Injury is more prevalent under poor drainage or compacted conditions or when soil is saturated for long periods of time. Soybeans rapidly outgrow stunting once favorable growing conditions return.

Injury to soybeans may occur if "Gemini" is used on soils having a calcareous surface layer or a pH of 7.0 or higher, or if used in conjunction with soil-applied insecticides. Seedling disease, cold or hot weather, deep planting (more than 2"), excessive moisture, high salt concentration, or drought may weaken soybean seedlings and increase possibility of crop injury. Do not apply to Black Belt Soils of Alabama and Mississippi with a soil pH greater than 7.0 or a history of nutrient deficiency such as iron chlorosis, as injury may occur.

ROTATIONAL CROP GUIDELINES

Crops that can be planted after using "Gemini" are listed below. Choice of rotational crop is determined by geographical location (state), soil pH, rate and time interval elapsed since last application.

GROUP I

States of: AL, AR, FL, GA, I A, MS, TX

- Soil pH 7.0 or less

GROUP II

States of: DE, KY, MD, MO Bootheel, NJ, NC, PA, SC, TN, VA, WV

- Soil pH 6.8 or less

AND

Use rate 16-22 oz/A

GROUP III

All other states not listed in Group I or II

- Soil pH 7.0 or less

OR

States of: AL, AR, FL, GA, LA, MS, TX

- Soil pH 7.1-7.5

OR

States of: DE, KY, MD, MO Bootheel, NJ, NC, PA, SC, TN, VA, WV

- Soil pH 6.9-7.5

OR

Fields not meeting rate requirement in Group II

TIME INTERVAL IN MONTHS BEFORE REPLANTING ROTATIONAL CROP

Crops	Group I	Group II	Group III
Soybeans	Anytime	Anytime	Anytime
Cereal grains	4	4	4
Ryegrass	4	4	4
Field Corn *	9	10	18
Alfalfa	10	10	18
Clover	10	12	18
Cotton	10	10	18
Peanuts	10	10	18
Sorghum	10	10	18
Tobacco (transplant)	10	10	18
Tomato (transplant)	10	10	18

All other crops require a minimum 18 month rotational interval and a successful field bioassaya.

* The term "FIELD CORN" is defined to include only that corn grown for grain or silage, popcorn, and seed corn relative to the "Rotational Crop Guidelines" section of the label, but because seed corn inbred lines may vary in their sensitivity to trace amounts of herbicide carryover Du Pont cannot warrant that seed corn can be recropped without damage or yield loss. Users are advised to seek the advice of their seed corn company agronomist regarding inbred sensitivity to herbicides prior to planting any inbred lines.

Suggested interval (months) before initiating bioassay is 10 months for Group I, 12 months for Group II and 15 months for Group III bioassay interval.

a Field Bioassay - The field bioassay will detect small quantities of "Gemini" which can remain in the soil and injure rotational crops. A successful field bioassay must be completed the season before planting any crop not listed above. A successful field bioassay means growing to maturity a test strip of the crop(s) intended for production the following year. The test strip should cross the whole field, including knolls and low areas.

SPRAYER CLEANUP

To avoid subsequent injury to desirable crops, thoroughly clean all mixing and spray equipment immediately following applications of "Gemini" as follows:*

1. Drain tank; thoroughly hose down the interior surfaces of the tank; then flush tank, boom, and hoses with clean water for a minimum of 5 minutes.
2. Partially fill the tank with water and add one of the cleaning agents listed below. Complete filling the tank with water, then flush the cleaning solution through the boom, hoses, and nozzles. Add water to completely fill the tank and allow to agitate or recirculate for at least 15 minutes. Again, flush the boom, hoses and nozzles, and drain the tank.
3. Repeat Step 2.
4. Remove the nozzles and screens and clean separately in a bucket containing water and the cleaning agent.
5. Thoroughly rinse the tank with clean water for a minimum of 5 minutes, flushing water through the boom and hoses.

NOTE: Use any of the following cleaning agents. Carefully read and follow the individual cleaning agent instructions.

1. One gallon of household ammonia (contains 3% active) per 100 gallons of water.
2. Nutra-Sol⁷
3. Loveland Tank and Equipment Cleaner⁸
4. Protank Cleaner⁹
5. Chem-Tank Cleaner and Neutralized¹⁰
6. Incide-Out¹¹
7. Tank-Aid¹²

PRECAUTIONS

Do not apply "Gemini" to soils with a pH greater than 7.0, except for those states listed in Group III of the "ROTATIONAL CROP GUIDELINES" where "Gemini" can be applied to soils up to 7.5 pH.

• CAUTION: Soil pH varies greatly, even within the same field. pH variations as much as 2 pH units are common. Composite soil samples taken across an entire field, such as those samples taken for soil fertility recommendations, may not detect areas of high pH (greater than pH 6.8). Sub-sampling areas likely to have pH values higher than the field average is recommended.

Soil pH may vary with soil type. Where different soil types are evident within a field, these types should be sampled separately.

Soil pH may vary due to conditions within a field. Such areas include, but are not limited to, areas bordered by limestone gravel roads, river bottoms subject to flooding, low areas in hardpan soils where evaporative ponds may occur, eroded hillsides, along drain tile lines, and areas where drainage ditch spoil has been spread. These areas should be sampled separately.

Soil pH may vary within the soil profile. For instance, fields in which lime has not been deeply incorporated may exhibit significantly higher pH values in the upper 3 inches of soil. Composite soil samples taken at a 6-8 inch depth may not reflect the elevated pH near the surface. In these cases shallow sampling, the upper 3 inches, is advised.

Determine soil pH by laboratory analysis using a 1:1 soil:water suspension.

- Do not graze treated fields or harvest for forage or hay.
- Do not use on sand, muck, or peat soils, nor on soils with less than 1/2% organic matter as crop injury may result.
- Do not apply "Gemini" after the soybean crop has emerged.
- Use of "Gemini" may affect rotational crops. Follow "ROTATIONAL CROP GUIDELINES".
- Do not apply to Black Belt Soils of Alabama and Mississippi with soil pH greater than 7.0, or on soils with a history of nutrient deficiencies such as iron chlorosis or injury may occur.
- When recommended tank mix combinations are used, consult the product label for replanting and recropping instructions and observe the directions that are the most restrictive.

- If Scepter®, Pursuit® or any product containing the active ingredients, imazaquin or imazethapyr (such as Squadron®, Partner®) is applied the same year as "Gemini", do not plant crops other than soybeans for at least 15 months from the last application. Crops listed in the "Gemini" Rotational Crop Guidelines" may be planted after 15 months; all other crops require a successful field bioassay.
- Do not apply to land that has been or will be treated with "Ally", "Finesse" or "Glean" herbicides in the states of Nebraska, Kansas, North Dakota and South Dakota.

- 1 Trademark of Monsanto Agricultural Products Co., St. Louis, MO
- 2 Trademark of Ciba-Geigy Corp., Greensboro, NC
- 3 Trademark of American Cyanamid Corp., Princeton, NJ
- 4 Trademark of Elanco Products Co., Indianapolis, IN
- 5 Trademark of Kalo Agricultural Chemicals, Inc., Overland Park, KS
- 6 Trademark of Witco Chemical Corp., Houston, TX
- 7 Compounded for Thomas G. Kilfoil Co., Inc., San Bruno, CA
- 8 Registered Trademark of Loveland Industries, Inc.
- 9 Manufactured for Cenex/Land 'O Lakes Agronomy Co.
- 10 Manufactured by Farmbelt Chemicals, Inc.
- 11 Trademark of Precision Laboratories, Inc.
- 12 Manufactured for Combelt Chemical Co.

STORAGE AND DISPOSAL

Storage: Store product in original container only, away from other pesticides, fertilizer, food, or feed.

Product Disposal: Do not contaminate water, food, or feed by storage or disposal. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

D-032093

NOTICE TO BUYER: Purchase of this material does not confer any rights under patents of countries outside of the United States.

NOTICE OF WARRANTY

Du Pont warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for the purposes stated on such label only when used in accordance with the directions under normal use conditions. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors such as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Du Pont. In no case shall Du Pont be liable for consequential, special or indirect damages resulting from the use or handling of this product. All such risks are assumed by the buyer. **DU PONT MAKES NO WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.**

BEST AVAILABLE COPY