

SUPPLEMENTAL LABELING**DuPont Agricultural
Products****MATRIX® HERBICIDE
ROTATION INTERVAL TO
CARROTS AND ONIONS**

"..... A Growing Partnership With Nature"

MATRIX® HERBICIDE

EPA Reg. No. 352-556

ROTATION INTERVAL TO CARROTS AND ONIONS**DIRECTIONS FOR USE**

MATRIX® Herbicide is a dry flowable formulation that selectively controls broadleaf weeds and grasses in potatoes. It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

CROP ROTATION

Carrots or onions grown under sprinkler irrigation can be planted 10 months after using MATRIX® herbicide on sprinkler irrigated potatoes grown in Morrow and Umatilla counties in Oregon, and in Benton, Franklin, Klickitat, and Walla Walla counties in Washington.

This rotation interval is for sand, loamy sand and sandy loam soils having not more than 1.5% organic matter where a minimum of 18 inches of sprinkler irrigation is used on the previous potato crop. Injury to the carrot or onion crop may occur if less than 18 inches of irrigation is used on the previous potato crop.

For tank mixtures, follow the most restrictive rotational crop guideline.

NOTE:


MATRIX® should not be used in a tankmix or sequential application program with other soil residual ALS-inhibiting herbicides on potatoes as the combined effects of these herbicides on the planting of subsequent crops have not been thoroughly investigated and crop injury may occur.

IMPORTANT

**BEFORE USING MATRIX® HERBICIDE,
READ AND FOLLOW ALL APPLICABLE
DIRECTIONS, RESTRICTIONS AND PRECAU-
TIONS ON THE EPA-REGISTERED LABEL.**

This bulletin contains new or supplemental instructions for use of this product which do not appear on the EPA-registered package label. Follow the instructions carefully.

This labeling must be in the possession of the user at the time of pesticide application.



DR-206 041902