


**DuPont Crop
Protection**



SUPPLEMENTAL LABELING

**MATRIX® HERBICIDE
LOW RATE USE OF
"EPTAM" IN POTATOES**

MATRIX® HERBICIDE

EPA Reg. No. 352-556

**LOW RATE USE OF "EPTAM" IN POTATOES -
FOR USE IN THE STATE OF OREGON**

GENERAL INFORMATION

DuPont MATRIX® Herbicide is a dry flowable formulation that selectively controls certain broadleaf weeds and grasses in potatoes. The best control is obtained when MATRIX® is applied to young, actively growing weeds. The degree and duration of control may depend on the weed spectrum and infestation intensity, the weed size at application and/or the environmental conditions at and following treatment.

MATRIX® is noncorrosive to equipment, nonflammable, and nonvolatile.

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

DIRECTIONS FOR USE

MATRIX® Herbicide can be tankmixed with "Eptam" herbicide and applied postemergence for improved control of common lambsquarter in potatoes.

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

POSTEMERGENCE APPLICATIONS

For postemergence applications, apply MATRIX® at 1.5 ounce per acre tankmixed with 1 pint of "Eptam 7-E" herbicide (or 0.88 ai/A of a comparable EPTC containing product registered for potatoes in Oregon) to young, actively growing weeds after crop emergence but before potatoes exceed 4 - 6" in height.

For best results with MATRIX® postemergence, rainfall or sprinkler irrigation of 1/3 to 1" (sandy soils apply at least 1/3", sandy loams apply at least 1/2", silt soils apply at least 3/4", clay soils apply at least 1"), no sooner than 4 hours, but not more than 5 days after application, will activate MATRIX® in the soil and help provide control of subsequent flushes of annual weeds.

Typically, small weeds (less than 1" in height or diameter) that are actively growing at application are most easily controlled. For control of common lambsquarter height should not exceed 3 inches. Postemergence applications can be made by ground or air.

Read and follow both product labels for this geographical area.

Adjuvants:

Postemergence applications of MATRIX® tankmixed with "Eptam" must include a Modified Seed Oil (MSO) and Ammonium nitrogen fertilizer.

Modified Seed Oil Adjuvants

- Apply at 1-% volume/volume (1 gal per 100-gal spray solution).
- Methylated seed oil (MSO) or Ethylated seed oil (ESO) are the preferred adjuvant.
- Oil adjuvants must contain at least 80% high quality, petroleum (mineral) or modified vegetable seed oil with at least 15% surfactant emulsifiers.

Ammonium Nitrogen Fertilizer

- An ammonium nitrogen fertilizer must be used in addition to the adjuvant. Use 2 lb/acre of a spray-grade ammonium sulfate (AMS).

IMPORTANT

**BEFORE USING MATRIX® HERBICIDE,
READ AND FOLLOW ALL APPLICABLE
DIRECTIONS, RESTRICTIONS AND PRECAU-
TIONS ON THE EPA-REGISTERED LABEL.**

This bulletin contains new or supplemental instructions for use of this product, which do not appear on the EPA registered package label.

Follow the instructions carefully.

This labeling must be in the possession of the user at the time of pesticide application.

"Eptam" is a registered trademark of Syngenta Crop Protection

DR-335 042803-2