

DuPont Agricultural Products

..... A Growing Partnership With Nature

ACCENT® SP HERBICIDE FOR WEED CONTROL IN FIELD CORN IN THE STATE OF CALIFORNIA ONLY

ACCENT® SP HERBICIDE

EPA REG. NO. 352-560

FOR WEED CONTROL IN FIELD CORN IN THE STATE OF CALIFORNIA ONLY

) DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

) DuPont "Accent" SP Herbicide is recommended for selected postemergence control of certain grass and broadleaf weeds in field corn in the state of California only.

Apply "Accent" SP at the rate of 2/3 oz product per acre (1 Soluble Packet per 4 treated acres) for selective postemergence grass weed control. Refer to the "Accent" SP product label for a complete list of weeds controlled and for recommended weed sizes at application. Weeds that exceed the listed weed sizes by up to 50% may be partially controlled with rates between 2/3 and 1 1/3 oz of "Accent" SP per acre.

DO NOT APPLY AERIALY IN CALIFORNIA.

) DO NOT USE AMMONIUM NITROGEN, LIQUID NITROGEN OR SPRAY GRADE NITROGEN AS A SPRAY ADJUVANT IN CALIFORNIA.

DO NOT APPLY ACCENT WITHIN 50 FEET OF SENSITIVE CROPS.

)

IMPORTANT


BEFORE USING "ACCENT" SP, READ AND FOLLOW ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA-REGISTERED LABEL.

This bulletin contains new or supplemental instructions for use of this product which do not appear on the EPA-registered package label. Follow the instructions carefully.

This labeling must be in the possession of the user at the time of pesticide application.

D-033094/051094

Expires 12/31/94



Reg # 352-560

PM-25

H-29  
Y 12




# Accent<sup>®</sup> SP

herbicide

---

RECEIVED  
 MAY 18 1994  
 Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under  
 EPA Reg. No. 352-560



®

*"..... A Growing Partnership With Nature"*

### ACCENT SP HIGHLIGHTS

- ACCENT SP provides selective postemergence grass control in field corn, field corn grown for seed, and popcorn.
- One 2-2/3 oz Soluble Packet treats four acres.
- Include an adjuvant as recommended in this label. See **Spray Additives**.
- The use of nitrogen fertilizer is recommended for certain grasses and for certain environmental conditions. See **Spray Additives**.
- ACCENT SP may be applied by ground (broadcast or band) or by air.
- For ground application, apply in 10 (light grass pressure) to 15 (heavy grass pressure) gal of water at 20-40 PSI using flat fan nozzles.
- Recommended tank mixes are specified on this label. See **Tank Mix Applications**.
- Apply to actively growing grasses at the recommended sizes. See **Rate**.
- Applications of ACCENT SP to grasses or corn under stress may affect the performance of ACCENT SP or in some cases reduce crop tolerance. See **Environmental Conditions**.
- Consult label text for complete instructions. Always read and follow label directions for use.

### TABLE OF CONTENTS

Precautionary Statements ..... 1

Application Information ..... 2

    When to Apply ..... 2

    Soil Insecticide Interaction Information ..... 2

    Timing to Crop Stage ..... 2

    Rate ..... 3

    Spray Adjuvants ..... 3

    Mixing Instructions ..... 4

    Tank Mix Applications ..... 4

    Sequential Applications ..... 5

Application Information ..... 6

    Ground Application ..... 6

    Aerial Application ..... 6

    Chemigation ..... 6

Environmental Conditions and

    Biological Activity ..... 6

    Cultivation ..... 6

    Crop Rotation ..... 7

    Sprayer Preparation/Cleanup ..... 7

    Spray Drift ..... 8

    Important Precautions ..... 8

    Storage and Disposal ..... 8

    Notice of Warranty ..... 9



## ACCENT® SP HERBICIDE

*For Use on Field Corn, Field Corn Grown for Seed, and Popcorn.*

This product is a water-dispersible granule containing 75% active ingredient by weight contained in a premeasured water-soluble packet.

<u>Active Ingredients</u>	<u>By Weight</u>
<i>nicosulfuron</i>	
<i>2-((( (4,6-Dimethoxypyrimidin-2-yl)aminocarbonyl)aminosulfonyl))- N,N-dimethyl-3-pyridinecarboxamide</i>	<i>75.0%</i>
<u>Inert Ingredients</u>	<u>25.0%</u>
<u>TOTAL</u>	<u>100.0%</u>

EPA Reg. No. 352 - 560

U.S. Patent No. 4,789,393

KEEP OUT OF REACH OF CHILDREN

### CAUTION

#### STATEMENT OF PRACTICAL TREATMENT

If in eyes: Flush eyes with plenty of water. Call a physician if irritation persists.

If on skin: Wash with plenty of soap and water.

If swallowed: No specific intervention is indicated as the compound is not likely to be hazardous by ingestion. However consult a physician if necessary.

*For medical emergencies involving this product, call toll-free 1-800-441-3637.*

#### PRECAUTIONARY STATEMENTS

##### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Caution! Causes moderate eye irritation. Harmful if absorbed through skin. Avoid contact with skin, eyes or clothing.

#### PERSONAL PROTECTIVE EQUIPMENT

Applicators and other handlers must wear:

Long-sleeved shirt and long pants.

Waterproof gloves.

Shoes plus socks.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

#### USER SAFETY RECOMMENDATIONS

**USERS SHOULD:** Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

#### ENVIRONMENTAL HAZARDS

Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment rinsewater. Do not apply where/when conditions could favor runoff. Do not apply if a severe storm is expected within 24 hours.

49 12

5912

## DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls.
- Waterproof gloves.
- Shoes plus socks.

ACCENT SP Herbicide should be used only in accordance with recommendations on this label or in supplemental Du Pont publications. Du Pont will not be responsible for losses or damage resulting from use of this product in any manner not specifically recommended by Du Pont.

### APPLICATION INFORMATION

Du Pont ACCENT SP Herbicide is a water-dispersible granule premeasured in a 2-2/3 oz packet that readily dissolves in water.

Apply ACCENT SP at the rate of 1 Soluble Packet per 4 treated acres for selective postemergence grass weed control.

For all application systems, use 50-mesh or larger strainer screens.

For use in California, see supplemental label "ACCENT SP For Use On Field Corn In California." Do not apply aerially in California.

Do not use ammonium nitrogen, liquid nitrogen or spray grade nitrogen as a spray adjuvant in California.

### WHEN TO APPLY

ACCENT SP performs best when applied to actively growing weeds at weed heights no greater than those given in the rate guide. See Rate.

### SOIL INSECTICIDE INTERACTION INFORMATION

Before using ACCENT SP, ensure that it is compatible with any other insecticides previously applied to the crop.

## CONVENTIONAL FIELD CORN, FIELD CORN GROWN FOR SEED, AND POPCORN

- ACCENT SP should not be applied if field corn was previously treated with Counter<sup>1</sup> 15G (all application methods), as severe crop injury may result.
- ACCENT SP should not be applied if field corn was previously treated with Counter<sup>1</sup> 20CP applied in-furrow at planting or over-the-row at cultivation.
- Applications of ACCENT SP to field corn previously treated with Counter 20CR applied T- and surface-banded on soil with less than or equal to 4.0% organic matter content may cause unacceptable injury. Du Pont will not be responsible for losses or damage resulting from such use.
- Application of ACCENT SP to field corn previously treated with Counter 20CR applied T- and surface-banded on soils with greater than 4% organic matter content, or to field corn previously treated with other organophosphate soil insecticides such as Thimet<sup>1</sup>, Dyfonate<sup>2</sup>, Lorsban<sup>3</sup>, etc. may result in temporary crop injury.
- For popcorn or field corn grown for seed, contact the seed supplier for full information on the use of ACCENT SP and its interaction with previously applied organophosphate insecticides.
- In all cases, the use of ACCENT SP on popcorn or field corn grown for seed that has been previously treated with Counter insecticide is prohibited.

### Herbicide-Resistant Field Corn

- ACCENT SP may be used on fields treated with Counter 15G or Counter 20CR (applied in-furrow, T- or surface-banded) if the field has been planted with an imidazolinone-resistant ("IR") hybrid such as Pioneer 3377 IR, Pioneer 3180IR, etc.
- For ACCENT SP applied to imazethapyr-tolerant ("IT") field corn hybrids, follow directions in Conventional Field Corn, Popcorn, and Field Corn Grown for Seed.

### TIMING TO CROP STAGE

#### Field Corn

- ACCENT SP may be broadcast or applied with drop nozzles to corn up to 24" tall (free-standing). The method of application depends on the crop:weed height differential.
- For corn 24-36" tall, apply ACCENT SP with drop nozzles only and avoid spraying into whorls of cornstalks.
- Do not apply to corn that is taller than 36" or that exhibits 10 collars, whichever is most restrictive.

#### Popcorn and Field Corn Grown for Seed

- ACCENT SP may be broadcast or applied with drop nozzles to popcorn or field corn grown for seed that is less than 20" tall (free-standing) or that exhibits 6 or fewer collars.
- Many seed companies have tested seed corn hybrids or popcorn hybrids for sensitivity to ACCENT SP and have reported excellent safety.

6912

- Not all seed corn inbreds or popcorn hybrids have been tested, nor does Du Pont have access to all seed company data. Consequently, Du Pont is not responsible for any crop injury arising from the use of ACCENT SP on field corn grown for seed or popcorn.
- Do not apply ACCENT SP to "White Dynamite" popcorn.
- See **Soil Insecticide Interaction Information** regarding the use of ACCENT SP on popcorn or field corn grown for seed that has been previously treated with a soil insecticide.
- Do not apply to popcorn or field corn grown for seed that is taller than 20" or that exhibits 6 collars, whichever is more restrictive.

**Timing to Weeds**

- Apply ACCENT SP when grasses are young and actively growing, but before they exceed the sizes indicated in the table below.
- Applications made to weeds at growth stages later than those listed below may vary from complete control to suppression. Yields may be reduced. Level of control will depend on the weed species, stage of growth, and environmental conditions.
- Treat heavy infestations of weeds before they become too competitive with the crop, especially where soil moisture and/or fertility are limited.
- ACCENT SP provides weed control via foliar absorption. ACCENT SP only controls those weeds that have emerged. For later-emerging weeds, a second application or a timely cultivation is required.
- Applications made to weeds larger than the size indicated or to weeds under stress may result in unsatisfactory control.

**RATE**

- Optimum control of the weeds listed can be achieved with 2/3 oz of ACCENT SP. Weeds that exceed the listed weed sizes by up to 50% may be partially controlled with rates between 2/3 and 1-1/3 oz of ACCENT SP per acre.
- In the WESTERN U.S. and WESTERN AREAS OF THE CORN BELT, arid conditions may require 1 oz of ACCENT SP to control difficult annual grasses such as green and yellow foxtail, wild proso millet, and sandbur. When selecting an adjuvant, use crop oil concentrate, or add liquid nitrogen fertilizer or ammonium sulfate to a nonionic surfactant.
- As weeds mature, their sensitivity to ACCENT SP decreases. As grassy weeds become mature (more than 3 tillers), they may not reach the size listed below, due to drought or other environmental factors. Grassy weeds that are maturing rapidly should be treated before they reach the stages listed below.

When applied as directed, ACCENT SP will control the following weeds:

	Height (Inches) at Application																	
Grasses	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18
Manisgrass 2-4"																		
Broadleaf signalgrass 1-2"																		
<b>Foxtails</b>																		
Birds 2-4"																		
Green 2-4"																		
Yellow 2-4"																		
Itchgrass 2-6"																		
<b>Johnsongrass</b>																		
Seedling 4-12"																		
Rhizome 8-18"																		
<b>Panicum</b>																		
Broomtop 1-3"																		
Fall 2-4"																		
Teas 1-3"																		
Quackgrass 4-10"																		
Ryegrass, Italian 2-6"																		
<b>Sandbur</b>																		
* Field 1-3"																		
* Longspur 1-3"																		
<b>Shattercane 4-12"</b>																		
Sorghum alba 4-12"																		
<b>Wild oats 2-6"</b>																		
* Wild proso millet 1-4"																		
* Woolly Cupgrass 2-4"																		

\* Size in inches or diameter, whichever is more restrictive

	Height (Inches) at Application																	
Broadleaves	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18
Paracucumber 1-3"																		
Jimsonweed 1-3"																		
<b>Morningglorys</b>																		
Tricolor 1-3"																		
Paint 1-2"																		
Tall 1-2"																		
<b>Pigweed</b>																		
Ruston 1-4"																		
Smooth 1-4"																		
Smartweed, annual 1-4"																		

**SPRAY ADJUVANTS**

Applications of ACCENT SP must include either a crop oil concentrate or a nonionic surfactant. The addition of ammonium nitrogen fertilizer is recommended. Additional information on adjuvant selection may be found in the bulletin "Approved Adjuvants For Use With DuPont Row Crop And Cereal Herbicides".

- Corn tolerance to ACCENT SP is very high, so adjuvant selection can be based solely on maximizing weed control.
- If another herbicide is tank mixed with ACCENT SP to increase the broadleaf weed spectrum, select adjuvants based on the adjuvant limitations of the other herbicide.

### **Crop Oil Concentrate (COC)**

- Apply at a rate (concentration) of 1.0% v/v (1 gal per 100 gal spray solution).
- The crop oil concentrate must be high quality, petroleum- or vegetable-seed oil-based product (methylated seed oil is considered a vegetable seed-based oil). It must contain at least 14% emulsifiers/surfactants.
- Crop oil concentrate is recommended for use when conditions have been hot and dry prior to ACCENT SP application.

### **Nonionic Surfactant (NIS)**

- Apply at a rate (concentration) of 0.25–0.5% v/v (1–2 qt per 100 gal spray solution). Use the higher rate in drought conditions to enhance control.
- At least 50% of the surfactant product must be active nonionic surfactant.
- Avoid products that do not accurately define their ingredients. Products must contain only EPA-exempt ingredients (40 CFR 1001).
- Biodegradable products are encouraged.
- Do not use products that change the pH of the spray tank solution.

### **Ammonium Nitrogen Fertilizer**

An ammonium nitrogen fertilizer may be added to either crop oil concentrate or nonionic surfactant to improve control under adverse conditions.

- Use a high-quality liquid nitrogen fertilizer such as 28–0–0 at a rate of 2–4 qt per acre or 10–34–0 at 1–2 qt per acre.
- A high-quality spray-grade ammonium sulfate (21–0–0) may be applied at a rate of 2–4 lb per acre in place of the liquid nitrogen fertilizer.
- Do not use liquid nitrogen fertilizers without either a crop oil concentrate or nonionic surfactant.
- Liquid nitrogen fertilizers should not be used as the total carrier solution.

### **MIXING INSTRUCTIONS**

Five Soluble Packets are contained in a waterproof, resealable bag. Three resealable bags are enclosed in each cardboard box. The individual Soluble Packets will dissolve completely in water. Open the outer resealable bag and remove the required number of Soluble Packets.

- To apply 2.5 oz product per acre, use one 2.5 oz Soluble Packet for every 4 treated acres.
- To apply 1 oz product per acre, use three 2.5 oz Soluble Packets for every 3 treated acres.

### **Soluble Packet Handling Precautions**

7/12

- The outer resealable bag is NOT soluble in water. DO NOT place it in the spray tank.
- Exposure to moisture or excessive handling of the Soluble Packets will cause them to break.
- Do not touch the packets with wet hands or place them on wet surfaces.
- Protect unused Soluble Packets by resealing them in the resealable bag.

### **Soluble Packet Use**

1. Fill the tank 1/4 to 1/3 full of water.
2. While agitating, add the required number of Soluble Packets.
3. Continue agitation while Soluble Packets dissolve (about 5 min).
4. Once the packets have dissolved, maintain agitation and continue filling tank with water. ACCENT SP should be thoroughly mixed with water before adding any other material.
5. As the tank is filling and after Soluble Packets have dissolved, add the required spray adjuvants (Crop Oil Concentrate, nonionic surfactant, liquid nitrogen fertilizer, or ammonium sulfate).
6. If the mixture is not continuously agitated, settling will occur. If settling occurs, thoroughly re-agitate before using.
7. Apply ACCENT SP spray mixture within 24 hours of mixing to avoid product degradation.
8. If ACCENT SP and a tank mix partner are to be applied in multiple loads, pre-slurry the ACCENT SP in clean water prior to adding to the tank. This will prevent the tank mix partner from interfering with the dissolution of the Soluble Packets.

### **TANK MIX APPLICATIONS**

#### **For Improved Control of Broadleaf Weeds**

- ACCENT SP may be tank mixed with other herbicides, listed below, for improved control of broadleaf weeds. See tank mix partner label for additional weeds controlled, weed sizes, and application conditions.
- For weeds listed on both ACCENT SP and tank mix partner product labels, follow the size guidelines that are least restrictive.
- Ammonium nitrogen fertilizer should be added to ACCENT SP tank mixes to control green and yellow foxtails to minimize or eliminate antagonism.
- An adjuvant must be used in all tank mixes. Consult table below for adjuvant recommendations. See *Spray Adjuvants* for adjuvant rate information.
- If antagonism occurs, complete control can be obtained with either a timely cultivation (see *Cultivation*) or a second application of ACCENT SP (see *Sequential Applications*).

8 9 12  
y

**Tank Mixes-Improved Broadleaf Control**

Product	Rate per Acre	Application Method	Adjuvant	Comments
Atrazine 4L or Atrazine 90DF	3/4 - 1-1/2 qt 0.83-1.66 lb	Broadcast on corn up to 12" tall.	COC	<ul style="list-style-type: none"> <li>Do not apply to corn taller than 12".</li> <li>Ammonium nitrogen fertilizer is recommended.</li> </ul>
Buctril *	1-1-1/2 pt	Broadcast or with drop nozzles on corn up to 24" tall; apply with drops only on corn up to 36" tall.	NIS	<ul style="list-style-type: none"> <li>Ammonium nitrogen fertilizer is recommended, especially where green or yellow foxtail is present.</li> <li>Do not apply ACCENT SP/ "Buctril" tank mixes on corn taller than 36" or that exhibits 10 collars, whichever is more restrictive.</li> <li>The substitution of COC for NIS may result in excessive crop injury.</li> </ul>
Buctril+atrazine	1-1/2-3 pt	Broadcast or with drop nozzles on corn up to 12" tall.	NIS	<ul style="list-style-type: none"> <li>Ammonium nitrogen fertilizer is recommended, especially where green or yellow foxtail is present.</li> <li>NIS is still required.</li> <li>Do not apply ACCENT SP/ "Buctril+ atrazine" tank mixes on corn taller than 12".</li> <li>The substitution of COC for NIS may result in excessive crop injury.</li> </ul>
Banvel <sup>†</sup>	1/2-1 pt  1/2 pt	Broadcast on corn to 8" tall.  Apply with drop nozzles only on corn 8-24" tall.	NIS	<ul style="list-style-type: none"> <li>Ammonium nitrogen fertilizer is recommended, especially where green or yellow foxtail is present.</li> <li>NIS is required.</li> <li>Do not apply ACCENT SP/ "Banvel" tank mixes to corn taller than 24".</li> <li>Do not substitute COC for NIS or excessive crop injury may occur.</li> </ul>
Marksman <sup>†</sup>	2-3 1/2 pt	Broadcast on corn up to 8" tall	NIS	<ul style="list-style-type: none"> <li>Ammonium nitrogen fertilizer is recommended, especially where green or yellow foxtail is present.</li> <li>NIS is still required.</li> <li>Do not substitute COC for NIS or excessive crop injury may occur.</li> </ul>

**TANK MIX APPLICATIONS**

**Additional Information**

- ACCENT SP should not be tank mixed with Basagran<sup>®</sup>, Laddok<sup>®</sup>, or Tandem<sup>®</sup> as severe crop injury may occur.
- ACCENT SP should not be tank mixed with 2, 4-D as severe grass control antagonism may occur.
- ACCENT SP should not be tank mixed with foliar-applied organophosphate insecticides such as Lorsban, malathion, parathion, etc., as severe crop injury may occur.
- To avoid crop injury or antagonism, apply Basagran, Laddok, Tandem, 2, 4-D, or organophosphate insecticides at least seven days before or 3 days after the application of ACCENT SP.
- ACCENT SP may be tank mixed with ASANA<sup>®</sup> XI or LANNATE<sup>®</sup> insecticides.

**SEQUENTIAL APPLICATIONS**

Annual grasses may have more than one flush of emerging seedlings. Also, regrowth of treated annual grasses may occur due to adverse environmental conditions following application. Perennial grasses may regrow from underground stems or roots, depending upon environmental conditions.

To control grasses under these conditions, a sequential application of ACCENT SP may be necessary. The combined dosage of the sequential applications cannot exceed 1 1/3 oz per acre of ACCENT SP.

**APPLICATION INFORMATION**

Many crops are highly sensitive to ACCENT SP. All direct or indirect contact (such as spray drift) with crops other than field corn, popcorn, or field crop soybean or field beans should be avoided.

**GROUND APPLICATION**  
(See also Spray Drift)

**Broadcast Application**

- Use a minimum of 10 gal of water per acre (GPA) for light, scattered stands of grass. Under heavy weed pressure, dense crop foliage or moisture stress, increase volume to at least 15 GPA.
- Use flat fan nozzles at 20–40 psi.
- Increase both spray volume and pressure as weed density and size increase.
- Do not use flood, hollow cone, rain drop, whirl chamber or controlled droplet applicator (CDA) nozzles. Unacceptable crop injury, excessive spray drift, or poor weed control may result.
- For proper spray coverage, adjust the boom and nozzle height according to manufacturers' specifications. For additional information on calibration see Du Pont's bulletin, "Application Accuracy for ACCENT Herbicide."
- Ensure that equipment is set up to avoid applying an excessive rate directly over the rows and into the corn plant whorl.
- Overlaps or starting, stopping, slowing, and turning while spraying may result in crop injury.

**Band Application**

- For band applications, use proportionately less spray mixture.
- To avoid crop injury, carefully calibrate the band applicator to not exceed the labeled rate.
- Carefully follow the manufacturer's instructions for nozzle type (flat fans), orientation, distance of nozzles from the crop and weeds, spray volumes, calibration and spray pressure.
- For additional information see Du Pont's bulletin, "Application Accuracy—Row Banders."

**AERIAL APPLICATION (EXCEPT CA)**  
(See also Spray Drift)

- Use nozzle types and arrangements that will provide optimum spray distribution and maximum coverage at 3 to 5 GPA.
- Do not apply during a temperature inversion, when winds are gusty, or when conditions favor poor coverage and/or off-target spray movement.

**Size of crop for aerial applications**

Field Corn	Seed Corn/ Popcorn	Minimum GPA	Target Grass
up to 24"	up to 20"	3-5	Shattercane Johnsongrass
up to 8" 8-16"	up to 8" 8-16"	3-5 5	Other Labelled Grasses

**CHEMIGATION**

ACCENT SP should not be applied through any type of irrigation system.

**ENVIRONMENTAL CONDITIONS  
AND BIOLOGICAL ACTIVITY**

9/12

ACCENT SP provides best results when applied to young, actively growing weeds. Success is heightened by warm, moist conditions (70 °F or more) and adequate soil moisture both before and after application. The degree and duration of control depend on: application rate, weed spectrum, weed size, growing conditions before and after treatment, soil moisture, precipitation, and adjuvants. Stress affects all weeds, but especially weeds such as field sandbur, woolly cupgrass, green and yellow foxtail, and wild proso millet. If weeds are under stress, delay application of ACCENT SP until the stress passes and weeds begin to grow again.

Treating weeds that exceed maximum label height or that are under stress may result in only partial control.

- ACCENT SP is rainfast in 4 hours.
- Applications made during or immediately following periods of large day/night temperature fluctuations or where daytime temperatures do not exceed 50 °F may decrease weed control.
- Ground application of ACCENT SP to dry, dusty fields may reduce weed control in wheel track areas.
- Poor weed control or crop injury may result from applications made to plants under stress from:
  - abnormally hot or cold weather
  - environmental conditions such as drought, water-saturated soils, hail damage, or frost
  - disease, insect, or nematode injury
  - prior herbicide, or carryover from a previous year's herbicide application

Delay application until stress passes and both weeds and corn resume growth. Severe stress from conditions immediately following application may also result in crop injury or poor weed control.

As weeds mature, their sensitivity to ACCENT SP decreases. As grassy weeds become mature (more than 3 tillers), they may be smaller than the size listed in Rate. When conditions exist where weeds are maturing rapidly, apply ACCENT SP to weeds that are smaller than those listed in Rate.

ACCENT SP rapidly inhibits the growth of susceptible weeds, reducing weed competition within as little as 6 hours after application. Susceptible plants are controlled in 7–21 days.

**CULTIVATION**

A timely cultivation may be necessary to control suppressed weeds, weeds beyond maximum size at application, or weeds that emerge after an application of ACCENT SP.

- Do not cultivate during the 10 days before ACCENT SP application
- Cultivation prior to application may decrease weed control by pruning roots and placing the weed under stress
- Optimum timing for cultivation is 7-14 days after ACCENT SP application

10912

**CROP ROTATION**

Rotational crops vary in their response to low concentrations of ACCENT SP remaining in the soil. ACCENT SP dissipates rapidly in warm, acidic, microbiologically active soils.

The amount of ACCENT SP which may be present in the soil depends on application rate, soil pH and organic matter content, elapsed time since application, crop production practices, and environmental factors.

Injury to rotational crops may occur in high-pH, cold soils if dry weather prevails between application and rotational crop planting. Consult your local Du Pont representative for additional guidelines.

Soil pH should be determined by laboratory analysis using the 1:1 soil:water suspension method on representative soil samples taken at 0-4" depth. Soil pH varies within fields; therefore, recropping should be based on the highest soil pH within each field. Consult local extension publications for recommended soil sampling procedures.

The following rotational intervals should be observed when using ACCENT SP:

**ACCENT SP Rotational Crop Guideline-1**

**No soil pH restrictions**

Crop Rotational	Interval in Months
Corn (Field, Seed)	Anytime
Corn (Pop, Sweet)*	10
Soybeans	10
Wheat (Winter)	4
Wheat (Spring)	8
Barley (Winter)	4
Barley (Spring)	8
Rye (Winter)	4
Dry Beans	10
Oats	8
Cotton	10
Peas, Snap Beans	10
Alfalfa**	12
Red Clover**	12
Other Crops	See Rotational Crop Guidelines 2 and 3

\* Except the sweet corn varieties "Merit", "Carnival", and "Sweet Success", for which the minimum time interval is 15 months

\*\*Except for the state of Kansas east of Highway 75, for Minnesota east and south of the Red River Valley and for the states east of the line formed by the western borders of Iowa, Missouri, Arkansas, and Louisiana, where the minimum time interval is 10 months

**ACCENT SP Rotational Crop Guideline-2**

**With soil pH > 6.5 restriction:**

Crop	Rotational Interval in Months	
	pH < 6.5	pH ≥ 6.5
Soybeans	10	18**
All other crops not listed in Rotational Guidelines 1 or 2	10	18**

\* Except on irrigated sites in Colorado, Wyoming, Nebraska, and Texas where the interval is 10 months on soils with pH < 7.5.

**ACCENT SP Rotational Crop Guideline-3**

**With soil pH 6.5 restrictions**

Crop	Rotational Interval in Months	
	pH < 6.5	pH ≥ 6.5
Sugar Beets*	10	18**
All other crops not listed in Rotational Guidelines 1 or 2	10	18**

\* Except on irrigated sites in Colorado, Wyoming, Nebraska, and Texas where the interval is 10 months on soils with pH < 7.5.

\*\*In North Dakota and northwest Minnesota, the cumulative precipitation in the 18 months following application must exceed 28" in order to rotate to sugarbeets or potatoes.

**SPRAYER PREPARATION/CLEANUP**

It is important that spray equipment is clean and free of previous pesticide deposits before using ACCENT SP and then properly cleaned out following application. Clean all application equipment before applying ACCENT SP. Follow the cleanup procedures specified on the label of the product previously sprayed. If no cleanup procedure is provided, use the procedure that follows. Immediately following applications of ACCENT SP, thoroughly clean all mixing and spray equipment to avoid subsequent crop injury.

**Notes**

- When cleaning spray equipment before applying ACCENT SP, read and follow label directions for proper rinsate disposal of the product previously sprayed.
- A steam cleaning of aerial spray tanks is recommended to dislodge any visible pesticide deposits.
- When spraying or mixing equipment will be used over an extended period to apply multiple loads of ACCENT SP, partially fill the tank with fresh water at the end of each day of spraying, flush the boom and hoses, and allow to sit overnight.

**Cleanup Procedure**

1. Drain the tank and thoroughly hose down the interior surfaces. Flush the tank, hoses, and boom with clean water for a minimum of 5 min.
2. Partially fill the tank with clean water and add one gal of household ammonia\* (containing 3% active) for every 100 gal of water. Finish filling the tank with water, then flush the cleaning solution through the hoses, boom, and nozzles. Add more water to completely fill the tank and allow to agitate/recirculate for at least 15 min. Again, flush the hoses, boom, and nozzles with the cleaning solution, then drain the tank.
3. Repeat Step 2.
4. Remove the nozzles and screens and clean separately in a bucket containing the cleaning agent and water.
5. Thoroughly rinse the tank with clean water for a minimum of 5 min, flushing the water through the boom and hoses.

\* Equivalent amount of an alternate strength cleaning solution or a tank cleaner recommended in the literature (e.g., Bulletin Sulfonixure Herbicide Application Equipment Cleanup) may be used.

### SPRAY DRIFT

Do not allow spray from either ground or aerial equipment to drift onto adjacent crops or land, as injury to other plants may occur. When spraying adjacent to crops other than corn, minimize the spray drift potential by following these practices.

- Stop spraying if wind speed becomes excessive. Spray drift can occur at wind speeds less than 10 MPH. Do not spray if sensitive crops are downwind. Do not spray if winds are gusty.
- High temperatures, drought, and low relative humidity increase the possibility of spray drift. Regardless of wind speed, extreme caution must be used when these conditions are present and sensitive crops or plants are nearby.
- Do not apply during a temperature inversion. An inversion is characterized by low air movement and an increase in air temperature with an increase in altitude. In humid regions, a fog or mist may form. An inversion may be detected by producing a smoke column and checking for a layering effect. Smoke-producing devices on aircraft are recommended. If not sure whether inversion conditions are present, consult with local weather services before making an application.

To further reduce drift from aerial or ground equipment:

- Use large-droplet size sprays to minimize drift. **DO NOT USE HOLLOW-CONE NOZZLES ON GROUND EQUIPMENT.** Do not use nozzles that produce very small droplets, such as airblast-type nozzles. For ground applications, nozzles should be oriented at an angle between straight down and straight back. For aerial applications, orient nozzles straight back along the windstream. If using flood-type nozzles on aircraft, orient the nozzles so spray is produced in direction of the airstream.
- Increase volume of spray mix per acre (for example, 10 GPA by air, 25 GPA by ground) by using higher-flow-rate nozzles.
- Reduce pressure (PSI). Do not exceed 40 psi when applying ACCENT SP. (Vehicle speed must also be reduced to maintain spray mix volume per acre.) Consult manufacturer's catalogs for details on correct calibration.
- Apply as close to target plants as possible while still maintaining a good spray pattern.

### IMPORTANT PRECAUTIONS

- Injury to or loss of desirable trees or vegetation may result from failure to observe the following:
- Do not apply ACCENT SP or drain or flush application equipment on or near desirable trees or other plants, or on areas where their roots may extend, or in locations where the chemical may be washed or moved into contact with their roots.
- Do not use on lawns, walks, driveways, tennis courts, or similar areas.
- Do not use on or near desirable plants.
- Do not use near any body of water.
- Thoroughly clean application equipment immediately after use. See "Sprayer Cleanup" section of the label for instructions.

- Do not graze or feed forage, hay, or straw from treated areas to livestock within 30 days of ACCENT SP application.
- Do not apply more than 1 1/3 oz per acre per season.
- In fields infested with Johnsongrass, or fields with a previous history of corn virus infection, a corn hybrid with a high degree of virus tolerance should be used. Consult your local seed corn representative for information on virus-tolerant hybrids.

### STORAGE AND DISPOSAL

Store product in original container only. Do not contaminate water, other pesticides, fertilizer, food or feed in storage. Store in a cool, dry place.

**Product Disposal:** Do not contaminate water, food, or feed by disposal. Waste resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**Container Disposal:** Do not reuse the outer box or the resealable plastic bag. When all water-soluble packets are used, the outer packaging should be clean and may be disposed of in a sanitary landfill or by incineration, or if allowed by state and local authorities, by open burning. If burned, stay out of smoke. If the resealable plastic bag contacts the formulated product in any way, the bag must be triple-rinsed with clean water. Add the rinsate to the spray tank and dispose of the outer wrap as described above.

**NOTICE TO BUYER:** Purchase of this material does not confer any rights under patents of countries outside of the United States.

12 of 12  
7

### NOTICE OF WARRANTY

Du Pont warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for the purposes stated on such label only when used in accordance with directions under normal use conditions. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Du Pont. In no case shall Du Pont be liable for consequential, special, or indirect damages resulting from the use or handling of this product. All such risks shall be assumed by the buyer. Du Pont will not be responsible for losses or damage resulting from application of ACCENT SP when applied to crops treated previously with "Counter" 20CR applied in a band at planting on soils of 4% organic matter content or less. DU PONT MAKES NO WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESSED OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

- 1 Registered trademark of American Cyanamid Co.;
  - 2 Registered trademark of ICI Americas, Inc.;
  - 3 Registered trademark of Dow Elanco;
  - 4 Registered trademark of Rhone-Poulenc Ag Company;
  - 5 Registered trademark of Sandoz Ltd.;
  - 6 Registered trademark of BASF Corporation;
  - 7 Registered trademark of E.I. Du Pont de Nemours & Co., Inc.
- AG-7721 9014/9034 1/6/94