




**DuPont Agricultural
Products**

"..... A Growing Partnership With Nature"



ACCENT® HERBICIDE
FOR WEED CONTROL
IN SWEET CORN FOR PROCESSING
IN THE STATES OF DELAWARE,
MARYLAND AND NEW JERSEY

ACCENT® HERBICIDE

EPA REG. NO. 352-560

**FOR WEED CONTROL IN SWEET CORN FOR PROCESSING
IN THE STATES OF DELAWARE, MARYLAND, AND NEW JERSEY**

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

DuPont "Accent" Herbicide may be applied to sweet corn grown under contract for processing. Consult the "Accent" label for additional requirements, restrictions and precautions.

WHEN TO APPLY

Applications of "Accent" may be applied broadcast or with drop nozzles (post-directed) on sweet corn up to 12 inches tall. For sweet corn 12 - 18 inches tall, apply only with drop nozzles. Do not apply to sweet corn taller than 18 inches OR THAT EXHIBITS 6 OR MORE COLLARS (V6), WHICHEVER IS MOST RESTRICTIVE. Only one application of "Accent" is allowed per year.

HOW TO USE

Apply "Accent" at 0.66 ounces per acre (or, when packed in water soluble packs, one soluble packet per 4 acres) on processing sweet corn for selective postemergence grass control. See Federal label for weeds controlled.

"Accent" performs best when applied to actively growing weeds at weed heights no greater than those given in the rate guide. Applications made to weeds larger than the size indicated or to weeds under stress may result in unsatisfactory control.

Adjuvants:

Always add either a crop oil concentrate (1% v/v = 1 gal/100 gal), methylated seed oil (1% v/v = 1 gal/100 gal) or nonionic surfactant (0.25% v/v = 1 qt/100 gal). The addition of an ammonium nitrogen fertilizer such as 28-0-0 (2 - 4 qt per acre), 10-34-0 (1 - 2 qt per acre) or ammonium sulfate (21-0-0) at 2 - 4 lb per acre is also recommended.

Broadcast Application:

Apply "Accent" using properly calibrated ground equipment only. Use flat fan nozzles at 20 - 40 psi and use a minimum of 10 gal water per acre. Increase both spray volume and pressure as weed density and size increase. Spray equipment must be properly set up to avoid applying an excessive rate directly over the rows and into the corn plant whorl.

Tank Mixes:

Do not tank mix "Accent" with any other product except those specified on this label. Atrazine can be tank mixed with "Accent" for improved control of broadleaf weeds. See the Atrazine label for additional information on rates to use and weeds controlled. Do not apply this tank mix to corn taller than 12 inches.

Rotational Crops:

Follow rotational intervals on "Accent" label.

RESTRICTIONS AND LIMITATIONS

"Accent" may be used only on the following sweet corn hybrids grown under contract for processing:

Bonus
Excalibur
GG-43
More(GG 55)
Reward
Viking
Zenith

Applying "Accent" to processing sweet corn under stress may result in crop injury. Stress may be caused by:

- abnormally hot or cold weather
- environmental conditions such as drought, water saturated soils, soil compaction, hail damage, or frost
- disease, insect, or nematode injury
- prior herbicide, or carryover from a previous year's herbicide application.

The user assumes all responsibility associated with applications of "Accent" to sweet corn.

Do not apply any organophosphate insecticide or any other herbicide within 7 days before or 3 days after an application of "Accent".

For product information call 1-888-6-DUPONT

© 1999 E. I. du Pont de Nemours and Company, Agricultural Products, Wilmington, Delaware 19898

2 9 7

Do not apply to sweet corn previously treated with any formulation of Counter¹ soil insecticide. Application of "Accent" to sweet corn previously treated with Thimet¹, Dyfonate², or Lorsban³ soil insecticide may cause unacceptable injury and is not recommended. IF A SOIL INSECTICIDE IS REQUIRED, DUPONT FORTRESS SOIL INSECTICIDE, AZTEC⁴ OR FORCE² ARE THE RECOMMENDED SOIL INSECTICIDES, AS THEY DO NOT INTERACT WITH ACCENT.

Applications within 4 hours of rain may reduce weed control.

A timely cultivation may be necessary to control weeds that emerge after an application of "Accent". Do not cultivate within 10 days prior to the application because weed control might be reduced by pruning roots and placing the weed under stress. Optimum timing for cultivation is 7 - 14 days after "Accent" application.

To avoid subsequent injury to crops other than corn, thoroughly clean all mixing and spray equipment immediately following application. Refer to the label for cleanout procedure.

Do not graze or feed forage, hay, or straw from treated areas to livestock within 30 days of "Accent" application.

IMPORTANT

BEFORE USING "ACCENT", READ AND FOLLOW ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA-REGISTERED LABEL.

This bulletin contains new or supplemental instructions for use of this product which do not appear on the EPA-registered package label. Follow the instructions carefully.

This labeling must be in the possession of the user at the time of pesticide application.

¹ Registered trademark of American Cyanamid Co.


² Registered trademark of Zeneca

³ Registered trademark of DowElanco

⁴ Registered trademark of Bayer

(Replaces H-63364 and H-63453)

D-011299




For product information call 1-888-6-DUPONT

© 1999 E. I. du Pont de Nemours and Company, Agricultural Products, Wilmington, Delaware 19898



**DuPont Agricultural
Products**

"..... A Growing Partnership With Nature"



3 7 7
SUPPLEMENTAL LABELING

**ACCENT® HERBICIDE
FOR WEED CONTROL IN SWEET CORN
FOR PROCESSING IN THE
STATES OF MINNESOTA, WISCONSIN,
ILLINOIS AND IOWA**

ACCENT® HERBICIDE

EPA REG. NO. 352-560

**FOR WEED CONTROL IN SWEET CORN FOR PROCESSING
IN THE STATES OF MINNESOTA, WISCONSIN, ILLINOIS AND IOWA**

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner consistent with its labeling.

DuPont "Accent" Herbicide may be applied to sweet corn grown under contract for processing. Consult the "Accent" label for additional requirements, restrictions and precautions.

WHEN TO APPLY

Applications of "Accent" may be applied broadcast or with drop nozzles (post-directed) on sweet corn up to 12 inches tall. For sweet corn 12 - 18 inches tall, apply only with drop nozzles. Do not apply to sweet corn taller than 18 inches OR THAT EXHIBITS 6 OR MORE COLLARS (V6), WHICHEVER IS MOST RESTRICTIVE. Only one application of "Accent" is allowed per year.

HOW TO USE

Apply "Accent" at 0.66 ounces per acre (or, when packed in water soluble packs, one soluble packet per 4 acres) on processing sweet corn for selective postemergence grass control. See Federal label for weeds controlled.

"Accent" performs best when applied to actively growing weeds at weed heights no greater than those given in the rate guide. Applications made to weeds larger than the size indicated or to weeds under stress may result in unsatisfactory control.

Adjuvants:

Always add either a crop oil concentrate (1% v/v = 1 gal/100 gal), methylated seed oil (1% v/v = 1 gal/100 gal) or nonionic surfactant (0.25% v/v = 1 qt/100 gal). The addition of an ammonium nitrogen fertilizer such as 28-0-0 (2 - 4 qt per acre), 10-34-0 (1 - 2 qt per acre) or ammonium sulfate (21-0-0) at 2 - 4 lb per acre is also recommended.

Broadcast Application:

Apply "Accent" using properly calibrated ground equipment only. Use flat fan nozzles at 20 - 40 psi and use a minimum of 10 gal water per acre. Increase both spray volume and pressure as weed density and size increase. Spray equipment must be properly set up to avoid applying an excessive rate directly over the rows and into the corn plant whorl.

Tank Mixes:

Do not tank mix "Accent" with any other product except those specified on this label. Atrazine can be tank mixed with "Accent" for improved control of broadleaf weeds. See the Atrazine label for additional information on rates to use and weeds controlled. Do not apply this tank mix to corn taller than 12 inches.

Rotational Crops:

Follow rotational intervals on "Accent" label.

RESTRICTIONS AND LIMITATIONS

"Accent" may be used only on the following sweet corn hybrids grown under contract for processing:

Bonus (GH 9056)	GG 43
Challenger	GG 46
Cornucopia	GG 55
Crisp'N'Sweet	GH 2628
DMC 20-04	GH 2690
DMC 20-35	HM701
Eliminator	Lumina
Excellency	Reward
Empire (GH 2759)	Sheba
Excalibur	Spirit
GG 5	Sprint (GH 2643)
GG 22	Viking
GG 23	Zenith
GG 40	

Applying "Accent" to processing sweet corn under stress may result in crop injury. Stress may be caused by:

- abnormally hot or cold weather
- environmental conditions such as drought, water saturated soils, soil compaction, hail damage, or frost
- disease, insect, or nematode injury
- prior herbicide, or carryover from a previous year's herbicide application.

For product information call 1-888-6-DUPONT

© 1999 E. I. du Pont de Nemours and Company, Agricultural Products, Wilmington, Delaware 19898

4.7.7

The user assumes all responsibility associated with applications of "Accent" to sweet corn.

Do not apply any organophosphate insecticide or any other herbicide within 7 days before or 3 days after an application of "Accent".

Do not apply to sweet corn previously treated with any formulation of Counter¹ soil insecticide. Application of "Accent" to sweet corn previously treated with Thimet¹, Dyfonate², or Lorsban³ soil insecticide may cause unacceptable injury and is not recommended. IF A SOIL INSECTICIDE IS REQUIRED, DUPONT FORTRESS SOIL INSECTICIDE, AZTEC⁴ OR FORCE² ARE THE RECOMMENDED SOIL INSECTICIDES, AS THEY DO NOT INTERACT WITH ACCENT.

Applications within 4 hours of rain may reduce weed control.

A timely cultivation may be necessary to control weeds that emerge after an application of "Accent". Do not cultivate within 10 days prior to the application because weed control might be reduced by pruning roots and placing the weed under stress. Optimum timing for cultivation is 7 - 14 days after "Accent" application.

To avoid subsequent injury to crops other than corn, thoroughly clean all mixing and spray equipment immediately following application. Refer to the label for cleanout procedure.

Do not graze or feed forage, hay, or straw from treated areas to livestock within 30 days of "Accent" application.

IMPORTANT

BEFORE USING "ACCENT", READ AND FOLLOW ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA-REGISTERED LABEL.

This bulletin contains new or supplemental instructions for use of this product which do not appear on the EPA-registered package label. Follow the instructions carefully.

This labeling must be in the possession of the user at the time of pesticide application.

¹ Registered trademark of American Cyanamid Co.


² Registered trademark of Zeneca

³ Registered trademark of DowElanco

⁴ Registered trademark of Bayer

(Replaces H-63363)

D-011299




For product information call 1-888-6-DUPONT

© 1999 E. I. du Pont de Nemours and Company, Agricultural Products, Wilmington, Delaware 19898



**DuPont Agricultural
Products**

"..... A Growing Partnership With Nature"



5 7 7
SUPPLEMENTAL LABELING

ACCENT® HERBICIDE
FOR CONTROL OF WIRESTEM
MUHLY IN CORN

ACCENT® HERBICIDE

EPA Reg. No. 352-560

FOR CONTROL OF WIRESTEM MUHLY IN CORN

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

DuPont ACCENT Herbicide may be applied in field corn grown grain to control wirestem muhly. Refer to the ACCENT package label for precautions if applying to seed corn or popcorn. ACCENT may be tank mixed with labeled broadleaf products for additional broadleaf weed control.

WHEN TO APPLY

Apply ACCENT in corn when wirestem muhly is 4 to 8 inches in height and is actively growing.

Application to wirestem muhly smaller than 4 inches or taller than 8 inches, or to wirestem muhly that is under stress, may result in unsatisfactory control and require a sequential application or cultivation. A sequential application of ACCENT or cultivation may be necessary to achieve optimum control of wirestem muhly.

ACCENT may be broadcast in corn up to 20 inches tall (free standing) or that has 6 or fewer collars (V6), whichever is more restrictive. For corn 20 to 36 inches tall apply ACCENT with drop nozzles only and avoid spraying into the whorl of corn-stalks. Do not apply to corn that is taller than 36 inches or that exhibits 10 or more collars (V10), whichever is more restrictive.

HOW TO USE

Apply 2/3 ounce per acre of ACCENT (one soluble packet per 4 acres). Add a crop oil concentrate at 1.0% v/v (1 gallon per 100 gallons of spray solution) and an ammonium nitrogen fertilizer such as 28-0-0 (2-4 qts/acre) or 10-34-0 (1 qt/acre). A high quality spray-grade ammonium sulfate (21-0-0) at 2 lb/acre may be used in place of the liquid nitrogen fertilizer.

Refer to the ACCENT label for further instructions on tank mixing for broadleaf weed control and other directions for use. Do not apply more than 1-1/3 ounces of ACCENT per acre/year.

TANK MIX

ACCENT may be tank mixed with ASANA XL Insecticide to aid in the control of insects such as common stalk borer which tend to be present in wirestem muhly.

CULTIVATION OR SEQUENTIAL ACCENT APPLICATION

Subsequent flushes of wirestem muhly, rhizomes not emerged at the time of application, or regrowth may require additional control measures such as sequential application of ACCENT or a cultivation. Do not cultivate during the 10 days prior to application of ACCENT. Optimum timing for cultivation is 7-14 days after application of ACCENT.

IMPORTANT PRECAUTIONS

Do not apply ACCENT through any type of irrigation system.


IMPORTANT

BEFORE USING "ACCENT " HERBICIDE, READ AND FOLLOW ALL APPLICABLE DIRECTIONS, RESTRICTIONS, AND PRECAUTIONS ON THE EPA-REGISTERED LABEL.

This bulletin contains new or supplemental instructions for use of these products which does not appear on the EPA-registered package labels. Follow these instructions carefully.

This label must be in the possession of the user at the time of application.

D-111198




For product information call 1-888-6-DUPONT

© 1998 E. I. du Pont de Nemours and Company, Agricultural Products, Wilmington, Delaware



**DuPont Agricultural
Products**

"..... A Growing Partnership With Nature"



6 7 /
SUPPLEMENTAL LABELING

**ACCENT® HERBICIDE
BASIS® GOLD HERBICIDE
CRABGRASS CONTROL**

ACCENT® HERBICIDE

EPA Reg. No. 352-560

BASIS® GOLD HERBICIDE

EPA Reg. No. 352-585

**TANK MIXES WITH REDUCED RATES OF PREEMERGENCE GRASS
HERBICIDES FOR CRABGRASS CONTROL IN CORN**

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

DuPont ACCENT or BASIS GOLD Herbicides may be used in a tank mix with reduced rates of preemergence grass herbicides for residual activity on later emerging grass flushes, including large crabgrass (*Digitaria sanguinalis*) and smooth crabgrass (*Digitaria ischaemum*).

ACCENT may be used in field corn, seed corn, and popcorn. Refer to the ACCENT label for use instructions in popcorn and seed corn. BASIS GOLD may be used in field corn grown for grain only.

ACCENT or BASIS GOLD may also be tank mixed with labeled broadleaf products for additional broadleaf weed control.

WHEN TO APPLY

Tank mixes of ACCENT or BASIS GOLD plus one of the pre-emergence grass herbicides listed below may be broadcast to corn before the crop exceeds the heights listed below.

Product	Maximum Corn Height
Surpass EC Herbicide ¹	11 inches
Harness Herbicide ¹	11 inches
Dual II Herbicide ³	12 inches
Dual II MAGNUM ³	12 inches
Frontier 6.0 Herbicide ⁴	8 inches
Prowl 3.3 EC Herbicide ⁵	6 leaf stage
LeadOff Herbicide	8 inches

Applications must be made before the crabgrass emerges (pre-emergence control only), and before the other grass weed

species exceed the sizes listed on the ACCENT or BASIS GOLD label. Refer to the ACCENT or BASIS GOLD label for a complete list of grasses controlled and the corn and weed sizes for application.

HOW TO USE

Rate

Apply a tank mix of ACCENT or BASIS GOLD plus a pre-emergence grass herbicide for postemergence control of the grasses listed on the ACCENT or BASIS GOLD label and residual control of large and smooth crabgrass as directed below:
ACCENT at a rate of 2/3 ounce per acre (1 soluble pack per 4 acres)

OR

BASIS GOLD at a rate of 14 ounces per acre (1 soluble pack per 4 acres)
plus -

Soil Type*

	Coarse	Medium	Fine
Surpass EC	1.5 pt	1.5 - 2 pt	1.5 - 2 pt
Harness	1.0 - 1.75 pt	1.25 - 2.25 pt	1.75 - 2.75 pt
Dual II	1.0 - 1.25 pt	1.25 - 1.75 pt	1.5 - 2.0 pt
Dual II MAGNUM	.67 - .8 pt	.8 - 1.2 pt	1.0 - 1.3 pt
Frontier 6.0	8 - 16 fl oz	8 - 16 fl oz	8 - 16 fl oz
Prowl 3.3 EC	1.8 - 2.4 pt	3.0 - 3.6 pt	3.0 - 3.6 pt
LeadOff	2.0-2.5 pt	2.5-3.5 pt	3.5- 4.5 pt

*Use the mid to lower end of the rate range for soils with less than 3% organic matter. Use the mid to upper end of the rate range for soils with 3% or more organic matter content.

For product information call 1-888-6-DUPONT

© 1998 E. I. du Pont de Nemours and Company, Agricultural Products, Wilmington, Delaware 19898

- plus -

A crop oil concentrate at 1.0% v/v (1 gallon per 100 gallons of spray solution) or a nonionic surfactant at 0.25% v/v (1 quart per 100 gallons of spray solution). In addition, an ammonium nitrogen fertilizer such as 28-0-0 (2 quarts per acre), 10-34-0 (1 quart per acre), or ammonium sulfate (21-0-0 at 2 lb. per acre) is recommended to improve postemergence control.

Refer to the ACCENT or BASIS GOLD labels for further instructions on tank mixing for broadleaf weed control.

IMPORTANT PRECAUTIONS

Refer to the ACCENT or BASIS GOLD label for mixing and other use instructions.

Preemergence herbicides rely on activation for control of grasses such as crabgrass that germinate or emerge after application.

Precipitation or overhead sprinkler irrigation is required after application to move the preemergence grass herbicide into the weed germination zone to control unemerged weeds. The amount of precipitation or irrigation required depends on existing soil moisture, soil type, and percent organic matter content, but 1/4 to 3/4 inch is normally adequate. If weeds emerge after treatment, immediately rotary hoe or use a shallow cultivation to improve performance.

Refer to the ACCENT or BASIS GOLD and the respective pre-emergence herbicide labels for additional use directions, use restrictions, rotational crop intervals, and precautions. The most restrictive provision on either label shall apply.

Do not apply ACCENT or BASIS GOLD through any type of irrigation system.


IMPORTANT

BEFORE USING "ACCENT" OR "BASIS GOLD " HERBICIDE, READ AND FOLLOW ALL APPLICABLE DIRECTIONS, RESTRICTIONS, AND PRECAUTIONS ON THE EPA-REGISTERED LABELS.

This bulletin contains new or supplemental instructions for use of these products which does not appear on the EPA-registered package labels. Follow these instructions carefully.

This label must be in the possession of the user at the time of application.

- 1 Registered trademark of Zeneca Inc.
- 2 Registered trademark of Monsanto Company
- 3 Registered trademark of Novartis Crop Protection, Inc.
- 4 Registered trademark of BASF
- 5 Registered trademark of American Cyanamid Company



(Replaces H-63217)

D-111198

For product information call 1-888-6-DUPONT

© 1998 E. I. du Pont de Nemours and Company, Agricultural Products, Wilmington, Delaware 19898