




Accent[®]

herbicide

DRAFT LABEL



“..... A Growing Partnership With Nature”

DUPONT ACCENT® HIGHLIGHTS

- ACCENT® provides selective postemergence grass control in field corn, field corn grown for seed, and popcorn.
- ACCENT® use rate is 2/3 to 1 1/3 oz per acre. [When packaged in water-soluble packets, one packet contains 2 2/3 oz product which treats 4 acres at the 2/3 oz per acre rate.]
- Include an adjuvant as recommended in this label. See **Spray Additives**.
- The use of nitrogen fertilizer is recommended for certain grasses and for certain environmental conditions. See **Spray Additives**.
- ACCENT® may be applied by ground (broadcast or band) or by air.
- For ground application, apply in 10 (light grass pressure) to 15 (heavy grass pressure) gal of water at 20-40 PSI using flat fan nozzles.
- Recommended tank mixes are specified on this label. See **Tank Mix Applications**.
- Apply to actively growing grasses at the recommended sizes. See **Rate**.
- Applications of ACCENT® to grasses of corn under stress may affect the performance of ACCENT® or in some cases reduce crop tolerance. See **Environmental Conditions**.
- Consult label text for complete instructions. Always read and follow label directions for use.

TABLE OF CONTENTS

Precautionary Statements	1
Application Information	2
When to Apply	2
Soil Insecticide Interaction Information	2
Normal Planned Use	2
Rescue Applications	3
Rate	3
Spray Adjuvants	4
Mixing Instructions	4
Tank Mix Applications	5
Sequential Applications	6
Application Information	6
Ground Application	6
Aerial Application	6
Chemigation	6
Environmental Conditions and	
Biological Activity	6
Cultivation	7
Crop Rotation	7
Sprayer Preparation/Cleanup	8
Spray Drift	8
Resistance	9
Integrated Pest Management	9
Important Precautions	9
Storage and Disposal	10
Notice of Warranty	10



Accent®

herbicide

For Use on Field Corn, Field Corn Grown for Seed, and Popcorn.

This product is a water-dispersible granule containing 75% active ingredient by weight. ACCENT® may be packaged in water soluble film packets.

<i>Active Ingredients</i>	<i>By Weight</i>
Nicosulfuron	
2-(((4,6-Dimethoxypyrimidin-2-yl)aminocarbonyl)aminosulfonyl)-N,N-dimethyl-3-pyridinecarboxamide	75.0%
<i>Inert Ingredients</i>	25.0%
TOTAL	100.0%

EPA Reg. No. 352 - 560

ACCEPTED

SEP 13 1999

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. **352-560**

KEEP OUT OF REACH OF CHILDREN

CAUTION

STATEMENT OF PRACTICAL TREATMENT

If in eyes: Flush eyes with plenty of water. Call a physician if irritation persists.

If on skin: Wash with plenty of soap and water.

If swallowed: No specific intervention is indicated as the compound is not likely to be hazardous by ingestion. However consult a physician if necessary.

For medical emergencies involving this product, call toll-free 1-800-441-3637.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Caution! Causes moderate eye irritation. Harmful if absorbed through skin. Avoid contact with skin, eyes or clothing.

PERSONAL PROTECTIVE EQUIPMENT

Applicators and other handlers must wear:
Long-sleeved shirt and long pants.
Waterproof gloves.
Shoes plus socks.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

USERS SHOULD: Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment rinsewater. Do not apply where/when conditions could favor runoff.

NOT REGISTERED

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 4 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls.
- Waterproof gloves.
- Shoes plus socks.

DuPont ACCENT® Herbicide should be used only in accordance with recommendations on this label or in supplemental DuPont publications. DuPont will not be responsible for losses or damage resulting from use of this product in any manner not specifically recommended by DuPont.

APPLICATION INFORMATION

DuPont ACCENT® Herbicide is a water-dispersible granule that readily dissolves in water.

Apply ACCENT® at the rate of 2/3 ounce per acre for selective postemergence grass weed control.

For all application systems, use 50-mesh or larger strainer screens.

Do not apply aurally in California or New York.

WHEN TO APPLY

ACCENT® performs best when applied to actively growing weeds at weed heights no greater than those given in the rate guide. See Rate.

SOIL INSECTICIDE INTERACTION INFORMATION

Before using ACCENT®, ensure that it is compatible with any other insecticides previously applied to the corn crop.

Conventional Field Corn and "IT" Hybrids, Field Corn Grown for Seed, and Popcorn

Soil Insecticides	Application Method	Soil O.M.	ACCENT® Use Precautions
Counter 15G ¹	All	All	Do not use.
Counter 20CR ²	In furrow at planting	All	Do not use.
	Over the row at cultivation	All	Do not use.
	T-band or Surface band	≤4%	May cause unacceptable injury.
	T-band or Surface band	>4%	May result in temporary injury.
Dyfonate ²	All labeled methods	All	May result in temporary injury.
Lorsban ¹	All labeled methods	All	May result in temporary injury.
Thimet ¹	All labeled methods	All	May result in temporary injury.
Fortress ⁴ , Aztec ³ , All and other Non-organophosphates		All	No use precautions.

- In all cases, the use of ACCENT® on popcorn or field corn grown for seed that has been previously treated with Counter insecticide is prohibited.
- For popcorn or field corn grown for seed, contact the seed supplier for full information on the use of ACCENT® and its interaction with previously applied organophosphate insecticides.

Herbicide-Resistant Field Corn

- ACCENT® may be used on fields treated with Counter 15G or Counter 20CR (applied in-furrow, T- or surface-banded) if the field has been planted with an imidazilinone-resistant ("IR") hybrid such as Pioneer 3377 IR, Pioneer 301R, etc.
- For ACCENT® applied to imidazilinone-tolerant ("IT") field corn hybrids, follow directions above for Conventional and IT Field Corn, Popcorn, and Field Corn Grown for Seed.

NORMAL PLANNED USE

- ACCENT® may be used on Field Corn, High Lysine, Waxy, White or other Food Grade corn hybrids.
- ACCENT® may be broadcast to corn up to 20" tall (free standing) or that has 6 or fewer collars (V6), whichever is more restrictive.

Timing to Weeds

- Apply DuPont ACCENT® when grasses are young and actively growing, but before they exceed the sizes indicated in the table below.
- Treat heavy infestations of weeds before they become too competitive with the crop, especially where soil moisture and/or fertility are limited.
- ACCENT® provides weed control via foliar absorption. ACCENT® only controls those weeds that have emerged. For later-emerging weeds, a second application or a timely cultivation is required.
- Applications made to weeds larger than the size indicated on this label or to weeds under stress may result in unsatisfactory control. Refer to LATE OR RESCUE APPLICATIONS.

LATE OR RESCUE APPLICATIONS

ACCENT® may be applied to field corn as a rescue treatment for the control of escaped grasses, or as a directed postemergence application on corn that is taller than 20" or which has more than 6 collars (which ever occurs first).

- For corn 20" to 36" tall, apply ACCENT® with drop nozzles only and avoid spraying into the whorl of cornstalks.
- Do not apply to corn that is taller than 36" or that exhibits 10 or more collars (V10), whichever is most restrictive.

Applications made to weeds larger than those listed on this label may vary from complete control to suppression. Level of control will depend on the weed species, stage of growth, and environmental conditions.

Due to the unplanned nature of rescue applications, choices must be made between the risks that arise from applications made beyond the proper time for ACCENT® use, and the effects of season long grass competition and/or harvest complications. These choices must balance risks from improperly timed ACCENT® use that include, but are not limited to:

- Yield loss due to competition: Research indicates competition from foxtail exceeding 4 inches in height may reduce corn yields. Applications to foxtail and other annual grasses that exceed the sizes stated on the label increases the risk of yield losses due to prolonged competition with the crop even though control may be acceptable.
- Incomplete control of grasses beyond labeled size: Applications to grasses that exceed the labeled sizes can result in reduced control. This incomplete control may reduce corn yield.
- Incomplete grass control due to herbicide stress: Grasses under stress from previous herbicide applications may not be actively growing and susceptible to ACCENT®. This stress may reduce grass control in "rescue" situations.
- Ear malformation: Applications of ACCENT® on corn that has 7 to 10 collars (V7 to V10) increases the potential for ear malformation (pinching). This risk may be greatly reduced, but not eliminated, by using drop nozzles properly adjusted so as to not apply ACCENT® into the corn whorl.

RATE

- Optimum control of the weeds listed can be achieved with 2/3 oz of ACCENT®. Weeds that exceed the listed weed sizes by up to 50% may be partially controlled with rates between 2/3 and 1-1/3 oz of ACCENT® per acre.
- In the WESTERN U.S. and WESTERN AREAS OF THE CORN BELT, to control difficult annual grasses such as green and yellow foxtail, wild proso millet and sandbur:
 - Always add a liquid nitrogen fertilizer or ammonium sulfate to a nonionic surfactant, or use a crop oil concentrate.
 - 1.0 oz of ACCENT® per acre may be required.
- As weeds mature, their sensitivity to ACCENT® decreases. As grassy weeds become mature (more than 3 tillers), they may not reach the size listed below, due to drought or other environmental factors. Grassy weeds that are maturing rapidly should be treated before they reach the stages listed below.

When applied as directed, ACCENT® will control the following weeds:

Grasses	Height (Inches) at Application
Barnyardgrass*	2-4"
Broadleaf signalgrass	1-2"
Foxtails	
Bristly	2-4"
Giant	2-4"
Green	2-4"
Yellow	2-4"
Itchgrass	2-6"
Johnsongrass	
Seedling	4-12"
Rhizome	8-18"
Panicum	
Browntop	1-3"
Fall	2-4"
Texas	1-3"
Quackgrass	4-10"
Ryegrass, Italian	2-6"
Sandbur	
Field*	1-3"
Longspine*	1-3"
Shattercane	4-12"
Sorghum alnum	4-12"
Wild oats	2-4"
Wild proso millet*	1-4"
Woolly Cupgrass**	2-4"

* Size in inches or diameter, whichever is more restrictive.

† Requires the use of COC plus ammonium nitrogen fertilizer.

Broadleaves	Height (Inches) at Application
Burcucumber	1-3"
Jimsonweed	1-3"
Morningglory	
Ivyleaf	1-3"
Pitted	1-2"
Tall	1-2"
Pigweed	
Redroot	1-4"
Smooth	1-4"
Smartweed, annual	1-4"

Popcorn and Field Corn Grown for Seed

- DuPont ACCENT® may be broadcast or applied with drop nozzles to popcorn or field corn grown for seed that is less than 20" tall (free-standing) or that exhibits 6 or fewer collars (V6), whichever is most restrictive.
- Do not apply to popcorn or field corn grown for seed that is taller than 20" or that exhibits 6 collars (V6), whichever is more restrictive.
- Many seed companies have tested seed corn inbreds or yellow popcorn hybrids for sensitivity to ACCENT® and have reported excellent safety.
- Not all seed corn inbreds or popcorn hybrids have been tested, nor does DuPont have access to all seed company data. Consequently, DuPont is not responsible for any crop injury arising from the use of ACCENT® on field corn grown for seed or popcorn.
- Do not apply ACCENT® to any white popcorn inbred, or white popcorn hybrid unless specifically approved by the seed company. This includes "White Dynamite" popcorn.
- Also check the tank mix component label for tolerance and instructions for use on seed corn inbreds.
- See Soil Insecticide Interaction Information regarding the use of ACCENT® on popcorn or field corn grown for seed that has been previously treated with a soil insecticide.

SPRAY ADJUVANTS

Applications of ACCENT® must include either a crop oil concentrate or a nonionic surfactant. The addition of ammonium nitrogen fertilizer is recommended. Additional information on adjuvant selection may be found in the bulletin "Approved Adjuvants For Use With DuPont Row Crop And Cereal Herbicides".

- Corn tolerance to ACCENT® is very high, so adjuvant selection can be based solely on maximizing weed control.
- If another herbicide is tank mixed with ACCENT® to increase the broadleaf weed spectrum, select adjuvants based on the adjuvant limitations of the other herbicide.

Crop Oil Concentrate (COC)

- Apply at a rate (concentration) of 1.0% v/v (1 gal per 100 gal spray solution).

- The crop oil concentrate must be high quality, petroleum- or vegetable-seed oil-based product (methylated seed oil is considered a vegetable seed-based oil). Petroleum based oils must contain at least 14% emulsifiers/surfactants.
- Crop oil concentrate is recommended for use when conditions have been hot and dry prior to ACCENT® application.

Nonionic Surfactant (NIS)

- Apply at a rate (concentration) of 0.25–0.5% v/v (1–2 qt per 100 gal spray solution). Use the higher rate in drought conditions to enhance control.
- At least 50% of the surfactant product must be active nonionic surfactant.
- Avoid products that do not accurately define their ingredients. Products must contain only EPA-exempt ingredients (40 CFR 1001).
- Biodegradable products are encouraged.
- Do not use products that change the pH of the spray tank solution.

Ammonium Nitrogen Fertilizer

An ammonium nitrogen fertilizer may be added to either crop oil concentrate or nonionic surfactant to improve control under adverse conditions.

- Use a high-quality liquid nitrogen fertilizer such as 28-0-0 at a rate of 2-4 qt per acre or 10-34-0 at 1-2 qt per acre.
- A high-quality spray-grade ammonium sulfate (21-0-0) may be applied at a rate of 2-4 lb per acre in place of the liquid nitrogen fertilizer.
- Do not use liquid nitrogen fertilizers without either a crop oil concentrate or nonionic surfactant.
- Liquid nitrogen fertilizers should not be used as the total carrier solution.

MIXING INSTRUCTIONS

1. Fill the tank 1/4 to 1/3 full of water.
2. While agitating, add the required amount of ACCENT®.
3. Continue agitation until the ACCENT® is fully dispersed, at least 5 minutes.
4. Once the ACCENT® is fully dispersed, maintain agitation and continue filling tank with water. ACCENT® should be thoroughly mixed with water before adding any other material.
5. As the tank is filling, add the required spray adjuvants (Crop Oil Concentrate, nonionic surfactant, liquid nitrogen fertilizer, or ammonium sulfate).
6. If the mixture is not continuously agitated, settling will occur. If settling occurs, thoroughly re-agitate before using.
7. Apply ACCENT® spray mixture within 24 hours of mixing to avoid product degradation.
8. If ACCENT® and a tank mix partner are to be applied in multiple loads, pre-slurry the ACCENT® in clean water prior to adding to the tank. This will prevent the tank mix partner from interfering with the dissolution of the ACCENT®.

Additional Use and Handling Information for DuPont ACCENT® When Packaged in Water Soluble Packets

Five Soluble Packets are contained in a waterproof, resealable bag. Three resealable bags are enclosed in each cardboard box. The individual Soluble Packets will dissolve completely in water. Open the outer resealable bag and remove the required number of Soluble Packets and follow the mixing instructions above.

- To apply 2/3 oz product per acre, use one 2-2/3 oz Soluble Packet for every 4 treated acres.
- To apply 1 oz product per acre, use three 2-2/3 oz Soluble Packets for every 8 treated acres.

Precautions

- The outer resealable bag is NOT soluble in water. DO NOT place it in the spray tank.
- Exposure to moisture or excessive handling of the Soluble Packets will cause them to break.
- Do not touch the packets with wet hands or place them on wet surfaces.
- Protect unused Soluble Packets by resealing them in the resealable bag.

TANK MIX APPLICATIONS

For Improved Control of Broadleaf Weeds

- ACCENT® may be tank mixed with other herbicides, listed below, for improved control of broadleaf weeds. See tank mix partner label for additional weeds controlled, weed sizes, and application conditions.
- For weeds listed on both ACCENT® and tank mix partner product labels, follow the size guidelines that are least restrictive.
- Ammonium nitrogen fertilizer should be added to ACCENT® tank mixes to control green and yellow foxtails to minimize or eliminate antagonism.
- An adjuvant must be used in all tank mixes. Consult table below for adjuvant recommendations. See *Spray Adjuvants* for adjuvant rate information.
- If antagonism occurs, complete control can be obtained with either a timely cultivation (see *Cultivation*) or a second application of DuPont ACCENT® (see *Sequential Applications*).

Tank Mixes—Improved Broadleaf Control

Product	Rate per Acre	Application Method	Adjuvant	Comments
Atrazine 4L or Atrazine 90DF	3/4 to 1-1/2 qt 0.83 to 1.66 lb	Broadcast on corn up to 12" tall.	COC	<ul style="list-style-type: none"> • Do not apply to corn taller than 12". • Ammonium nitrogen fertilizer is recommended.
Buctril® Buctril® Gel	1 to 1 1/2 pt 1/2 to 3/4 pt	Broadcast or with drop nozzles on corn up to 24" tall; apply with drops only on corn up to 36" tall.	NIS	<ul style="list-style-type: none"> • Ammonium nitrogen fertilizer is recommended, especially where green or yellow foxtail is present. • Do not apply ACCENT®/ "Buctril" tank mixes on corn taller than 36" or that exhibits 10 collars (V10), whichever is more restrictive. • The substitution of COC for NIS may result in excessive crop injury.
Buctril®+atrazine	1 1/2 to 3 pt	Broadcast or with drop nozzles on corn up to 12" tall.	NIS	<ul style="list-style-type: none"> • Ammonium nitrogen fertilizer is recommended, especially where green or yellow foxtail is present. • NIS is still required. • Do not apply ACCENT®/ "Buctril+ atrazine" tank mixes on corn taller than 12". • The substitution of COC for NIS may result in excessive crop injury.
Banvel® / Clarity®	1/2 to 1 pt 1/2 pt	Broadcast on corn to 8" tall. Apply with drop nozzles only on corn 8-24" tall with "Banvel", 8-12" tall with "Clarity".	NIS	<ul style="list-style-type: none"> • Ammonium nitrogen fertilizer is recommended, especially where green or yellow foxtail is present. • NIS is required. • Do not exceed the maximum "Banvel"/ "Clarity" labeled corn size when tank mixing with ACCENT®. • Do not substitute COC for NIS or excessive crop injury may occur.
Marksman®	2 to 3 1/2 pt	Broadcast on corn up to 8" tall.	NIS	<ul style="list-style-type: none"> • Ammonium nitrogen fertilizer is recommended, especially where green or yellow foxtail is present. • NIS is still required. • Do not apply ACCENT®/ "Marksman" tank mixes on corn taller than 12". • Do not substitute COC for NIS or excessive crop injury may occur.

TANK MIX APPLICATIONS

Additional Information

- Corn plants' predisposition to develop fused tissue emerging from the whorl (rat tail) after the V11 stage may increase when a product containing dicamba (such as Banvel, Clarity, or Marksman) is applied to small corn under early, stressful conditions. Be aware of this when applying tank mixes with dicamba to small corn (V3 stage or smaller) under stressful conditions (See Environmental Conditions for a description of these stressful conditions).
- ACCENT® should not be tank mixed with Basagran®, Laddok®, or Tandem® as severe crop injury may occur.
- ACCENT® should not be tank mixed with 2, 4-D or DuPont Bladex® Herbicide as severe grass control antagonism may occur.
- ACCENT® should not be tank mixed with foliar-applied organophosphate insecticides such as Lorsban, malathion, parathion, etc., as severe crop injury may occur.
- To avoid crop injury or antagonism, apply Basagran, Laddok, Tandem, 2, 4-D, or organophosphate insecticides at least seven days before or 3 days after the application of ACCENT®.
- ACCENT® may be tank mixed with ASANA® XL or LANNATE® insecticides.

SEQUENTIAL APPLICATIONS

Annual grasses may have more than one flush of emerging seedlings. Also, regrowth of treated annual grasses may occur due to adverse environmental conditions following application. Perennial grasses may regrow from underground stems or roots, depending upon environmental conditions.

To control grasses under these conditions, a sequential application of ACCENT® may be necessary. The combined dosage of the sequential applications cannot exceed 1-1/3 oz per acre of ACCENT®.

APPLICATION INFORMATION

Many crops are highly sensitive to ACCENT®. All direct or indirect contact (such as spray drift) with crops other than field corn, popcorn, or field corn grown for seed should be avoided.

GROUND APPLICATION (SEE ALSO SPRAY DRIFT)

Broadcast Application

- Use a minimum of 10 gal of water per acre (GPA) for light, scattered stands of grass. Under heavy weed pressure, dense crop foliage or moisture stress, increase volume to at least 15 GPA.
- Use flat fan nozzles at 20-40 psi.
- Increase both spray volume and pressure as weed density and size increase.
- Do not use flood, hollow cone, rain drop, whirl chamber controlled droplet applicator (CDA) nozzles or air assisted sprayers. Unacceptable crop injury, excessive spray drift, or poor weed control may result.

- For proper spray coverage, adjust the boom and nozzle height according to manufacturers' specifications. For additional information on calibration see DuPont's bulletin, "Application Accuracy for ACCENT® Herbicide."
- Ensure that equipment is set up to avoid applying an excessive rate directly over the rows and into the corn plant whorl.
- Overlaps or starting, stopping, slowing, and turning while spraying may result in crop injury.

Band Application

- For band applications, use proportionately less spray mixture.
- To avoid crop injury, carefully calibrate the band applicator to not exceed the labeled rate.
- Carefully follow the manufacturer's instructions for nozzle type (flat fans), orientation, distance of nozzles from the crop and weeds, spray volumes, calibration and spray pressure.

AERIAL APPLICATION (EXCEPT CA AND NY) (See also Spray Drift)

- Use nozzle types and arrangements that will provide optimum spray distribution and maximum coverage at 3 to 5 GPA.
- Do not apply during a temperature inversion, when winds are gusty, or when conditions favor poor coverage and/or off-target spray movement.

Size of crop for aerial applications

Field Corn	Seed Corn/ Popcorn	Minimum GPA	Target Grass
up to 20"/ 6 collars (V6)	up to 20"/ 6 collars (V6)	3-5	Shattercane Johnsongrass
up to 8" 8-16"	up to 8" 8-16"	3-5 5	Other Labeled Grasses

CHEMIGATION

ACCENT® should not be applied through any type of irrigation system.

ENVIRONMENTAL CONDITIONS AND BIOLOGICAL ACTIVITY

ACCENT® provides best results when applied to young, actively growing weeds. Success is heightened by warm, moist conditions (70 °F or more) and adequate soil moisture both before and after application. The degree and duration of control depend on: application rate, weed spectrum, weed size, growing conditions before and after treatment, soil moisture, precipitation, and adjuvants. Stress affects all weeds, but especially weeds such as field sandbur, woolly cupgrass, green and yellow foxtail, and wild proso millet. If weeds are under stress, delay application of ACCENT® until the stress passes and weeds begin to grow again.

Treating weeds that exceed maximum label height or that are under stress may result in only partial control.

- ACCENT® is rainfast in 4 hours.
- Applications made during or immediately following periods of large day/night temperature fluctuations or

where daytime temperatures do not exceed 50 °F may decrease weed control.

- Ground application of DuPont ACCENT® to dry, dusty fields may reduce weed control in wheel track areas.
- Poor weed control or crop injury may result from applications made to plants under stress from:
 - abnormally hot or cold weather
 - environmental conditions such as drought, water-saturated soils, hail damage, or frost
 - disease, insect, or nematode injury
 - prior herbicide, or carryover from a previous year's herbicide application

Delay application until stress passes and both weeds and corn resume growth. Severe stress from conditions immediately following application may also result in crop injury or poor weed control.

As weeds mature, their sensitivity to ACCENT® decreases. As grassy weeds become mature (more than 3 tillers), they may be smaller than the size listed in **Rate**. When conditions exist where weeds are maturing rapidly, apply ACCENT® to weeds that are smaller than those listed in **Rate**.

ACCENT® rapidly inhibits the growth of susceptible weeds, reducing weed competition within as little as 6 hours after application. Susceptible plants are controlled in 7–21 days.

CULTIVATION

A timely cultivation may be necessary to control suppressed weeds, weeds beyond maximum size at application, or weeds that emerge after an application of ACCENT®.

- Do not cultivate during the 10 days before ACCENT® application.
- Cultivation prior to application may decrease weed control by pruning roots and placing the weed under stress.
- Optimum timing for cultivation is 7–14 days after ACCENT® application.

CROP ROTATION

Rotational crops vary in their response to low concentrations of ACCENT® remaining in the soil. ACCENT® dissipates rapidly in warm, acidic, microbiologically active soils.

The amount of ACCENT® which may be present in the soil depends on application rate, soil pH and organic matter content, elapsed time since application, crop production practices, and environmental factors.

Injury to rotational crops may occur in high-pH, cold soils if dry weather prevails between application and rotational crop planting. Consult your local DuPont representative for additional guidelines.

Soil pH should be determined by laboratory analysis using the 1:1 soil:water suspension method on representative soil samples taken at 0–4" depth. Soil pH varies within fields; therefore, recropping should be based on the highest soil pH within each field. Consult local extension publications for recommended soil sampling procedures.

The following rotational intervals should be observed when using ACCENT®:

ACCENT® Rotational Crop Guideline–1

No soil pH restrictions

Crop Rotational	Interval in Months
Corn (Field, Seed)	Anytime
Corn (Pop, Sweet)*	10
Soybeans	0.5 (15 days)
Wheat (Winter)	4
Wheat (Spring)	8
Barley (Winter)	4
Barley (Spring)	8
Rye (Winter)	4
Dry Beans	10
Oats	8
Cotton	10
Peas, Snap Beans	10
Alfalfa**	12
Red Clover**	12
Other Crops	See Rotational Crop Guidelines 2 and 3

* Except the sweet corn varieties "Merit", "Carnival", and "Sweet Success", for which the minimum time interval is 15 months.

**Except for the state of Kansas east of Highway 75, for Minnesota east and south of the Red River Valley and for the states east of the line formed by the western borders of Iowa, Missouri, Arkansas, and Louisiana, where the minimum time interval is 10 months.

ACCENT® Rotational Crop Guideline–2

With soil pH 7.5 restrictions

Crop	Rotational Interval in Months	
	pH ≤ 7.5	pH > 7.5
Sorghum	10	18*
Sunflowers	11**	18
All other crops not listed	See Rotational Guideline 3 in Rotational Guidelines 1 or 2	

* Except in Texas and Oklahoma east of highway 281, where the rotational interval is 10 months, regardless of pH.

**Precipitation following application must exceed 14" prior to planting sunflowers.

ACCENT® Rotational Crop Guideline–3

With soil pH 6.5 restrictions

Crop	Rotational Interval in Months	
	pH ≤ 6.5	pH > 6.5
Sugar Beets*	10	18**
All other crops not listed	10	18**
in Rotational Guidelines 1 or 2		

* Except on irrigated sites in Colorado, Wyoming, Nebraska, Texas, or in Michigan where precipitation following application must exceed 25" prior to planting beets, where the interval is 10 months on soils with pH < 7.5.

**In North Dakota and northwest Minnesota, the cumulative precipitation in the 18 months following application must exceed 28" in order to rotate to sugarbeets or potatoes.

SPRAYER PREPARATION/CLEANUP

It is important that spray equipment is clean and free of previous pesticide deposits before using DuPont ACCENT® and then properly cleaned out following application. Clean all application equipment before applying ACCENT®. Follow the cleanup procedures specified on the label of the product previously sprayed. If no cleanup procedure is provided, use the procedure that follows. Immediately following applications of ACCENT®, thoroughly clean all mixing and spray equipment to avoid subsequent crop injury.

Notes

- When cleaning spray equipment before applying ACCENT®, read and follow label directions for proper rinsate disposal of the product previously sprayed.
- A steam cleaning of aerial spray tanks is recommended to dislodge any visible pesticide deposits.
- When spraying or mixing equipment will be used over an extended period to apply multiple loads of ACCENT®, partially fill the tank with fresh water at the end of each day of spraying, flush the boom and hoses, and allow to sit overnight.

Cleanup Procedure

1. Drain the tank and thoroughly hose down the interior surfaces. Flush the tank, hoses, and boom with clean water for a minimum of 5 min.
 2. Partially fill the tank with clean water and add one gal of household ammonia* (containing 3% active) for every 100 gal of water. Finish filling the tank with water, then flush the cleaning solution through the hoses, boom, and nozzles. Add more water to completely fill the tank and allow to agitate/recirculate for at least 15 min. Again, flush the hoses, boom, and nozzles with the cleaning solution, then drain the tank.
 3. Repeat Step 2.
 4. Remove the nozzles and screens and clean separately in a bucket containing the cleaning agent and water.
 5. Thoroughly rinse the tank with clean water for a minimum of 5 min, flushing the water through the hoses and boom.
- * Equivalent amounts of an alternate strength ammonia solution or a tank cleaner recommended in the DuPont bulletin "Sulfonylurea Herbicides, A Guide to Equipment Cleanout," may be used.

SPRAY DRIFT MANAGEMENT

The interaction of many equipment and weather-related factors determines the potential for spray drift. The applicator is responsible for considering all these factors when making application decisions.

AVOIDING SPRAY DRIFT IS THE RESPONSIBILITY OF THE APPLICATOR.

IMPORTANCE OF DROPLET SIZE

The most effective way to reduce drift potential is to apply large droplets (>150 - 200 microns). The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. The presence of sensitive species nearby, the environmental conditions, and pest pressure may affect how an applicator balances drift control and coverage. **APPLYING LARGER DROPLETS REDUCES DRIFT POTENTIAL, BUT WILL NOT PREVENT DRIFT IF APPLICATIONS ARE MADE IMPROPERLY OR UNDER UNFAVORABLE ENVIRONMENTAL CONDITIONS!** See **Wind, Temperature and Humidity**, and **Temperature Inversions** sections of this label.

Controlling Droplet Size - General Techniques

- **Volume** - Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets.
- **Pressure** - Use the lower spray pressures recommended for the nozzle. Higher pressure reduces droplet size and does not improve canopy penetration. **WHEN HIGHER FLOW RATES ARE NEEDED, USE A HIGHER-CAPACITY NOZZLE INSTEAD OF INCREASING PRESSURE.**
- **Nozzle Type** - Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles.

Controlling Droplet Size - Aircraft

- **Number of Nozzles** - Use the minimum number of nozzles with the highest flow rate that provide uniform coverage.
- **Nozzle Orientation** - Orienting nozzles so that the spray is emitted backwards, parallel to the airstream will produce larger droplets than other orientations.
- **Nozzle Type** - Solid stream nozzles (such as disc and core with swirl plate removed) oriented straight back produce larger droplets than other nozzle types.
- **Boom Length** - The boom length should not exceed 3/4 of the wing or rotor length - longer booms increase drift potential.
- **Application Height** - Application more than 10 ft above the canopy increases the potential for spray drift.

BOOM HEIGHT

Setting the boom at the lowest labeled height (if specified) which provides uniform coverage reduces the exposure of droplets to evaporation and wind. For ground equipment, the boom should remain level with the crop and have minimal bounce.

WIND

Drift potential increases at wind speeds of less than 3 mph (due to inversion potential) or more than 10 mph. However, many factors, including droplet size and equipment type determine drift potential at any given wind speed. **AVOID GUSTY OR WINDLESS CONDITIONS.**

Note: Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they affect spray drift.

TEMPERATURE AND HUMIDITY

When making applications in hot and dry conditions, set up equipment to produce larger droplets to reduce effects of evaporation.

TEMPERATURE INVERSIONS

Drift potential is high during a temperature inversion. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain close to the ground and move laterally in a concentrated cloud. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

SHIELDED SPRAYERS

Shielding the boom or individual nozzles can reduce the effects of wind. However, it is the responsibility of the applicator to verify that the shields are preventing drift and not interfering with uniform deposition of the product.

RESISTANCE

When herbicides with the same mode of action are used repeatedly over several years to control the same weed species in the same field, naturally-occurring resistant weed biotypes may survive a correctly applied herbicide treatment, propagate, and become dominant in that field. These resistant weed biotypes may not be adequately controlled. Cultural practices such as tillage, preventing weed escapes from going to seed, and using herbicides with different modes of action within and between crop seasons can aid in delaying the proliferation and possible dominance of herbicide resistant weed biotypes.

INTEGRATED PEST MANAGEMENT

DuPont recommends the use of Integrated Pest Management (IPM) programs to control pests. This product may be used as part of an Integrated Pest Management (IPM) program which can include biological, cultural, and genetic practices aimed at preventing economic pest damage. Application of this product should be based on IPM principles and practices including field scouting or other detection methods, correct target pest identification, population monitoring, and treating when target pest populations reach locally determined action thresholds. Consult your state cooperative extension service, professional consultants or other qualified authorities to determine appropriate action treatment threshold levels for treating specific pest/crop systems in your area.

IMPORTANT PRECAUTIONS

- Injury to or loss of desirable trees or vegetation may result from failure to observe the following:
- Do not apply DuPont ACCENT® or drain or flush application equipment on or near desirable trees or other plants, or on areas where their roots may extend, or in locations where the chemical may be washed or moved into contact with their roots.
- Do not use on lawns, walks, driveways, tennis courts, or similar areas.
- Prevent drift of spray to desirable plants.
- Do not contaminate any body of water.
- Thoroughly clean application equipment immediately after use. (See the Sprayer Cleanup section of this label for instructions.)
- Do not graze or feed forage, hay, or straw from treated areas to livestock within 30 days of ACCENT® application.
- Do not apply more than 1 1/3 oz per acre per season.
- In fields infested with Johnsongrass, or fields with a previous history of corn virus infection, a corn hybrid with a high degree of virus tolerance should be used. Consult your local seed corn representative for information on virus-tolerant hybrids.

STORAGE AND DISPOSAL

Store product in original container only. Do not contaminate water, other pesticides, fertilizer, food or feed in storage. Store in a cool, dry place.

PRODUCT DISPOSAL: Do not contaminate water, food, or feed by disposal. Waste resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Triple-rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

CONTAINER DISPOSAL (SOLUBLE PACKETS): Do not reuse the outer box or the resealable plastic bag. When all water-soluble packets are used, the outer packaging should be clean and may be disposed of in a sanitary landfill or by incineration, or if allowed by state and local authorities, by open burning. If burned, stay out of smoke. If the resealable plastic bag contacts the formulated product in any way, the bag must be triple-rinsed with clean water. Add the rinsate to the spray tank and dispose of the outer wrap as described above.

NOTICE TO BUYER: Purchase of this material does not confer any rights under patents of countries outside of the United States.

- 1 Registered trademark of American Cyanamid Co.
- 2 Registered trademark of Zeneca Inc.
- 3 Registered trademark of Dow Agro Sciences
- 4 Registered trademark of Rhone-Poulenc Ag Company
- 5 Registered trademark of BASF Corporation
- 6 Registered trademark of E. I. du Pont de Nemours & Co., Inc.
- 7 Registered trademark of Bayer Corporation

D - 692 072999

LIMITATION OF

WARRANTY AND LIABILITY

NOTICE: Read This Limitation of Warranty and Liability Before Buying or Using This Product. If the Terms Are Not Acceptable, Return the Product at Once, Unopened, and the Purchase Price Will Be Refunded.

It is impossible to eliminate all risks associated with the use of this product. Such risks arise from weather conditions, soil factors, off target movement, unconventional farming techniques, presence of other materials, the manner of use or application, or other unknown factors, all of which are beyond the control of DuPont. These risks can cause: ineffectiveness of the product; crop injury, or; injury to non-target crops or plants.

DuPont does not agree to be an insurer of these risks. **WHEN YOU BUY OR USE THIS PRODUCT, YOU AGREE TO ACCEPT THESE RISKS.**

DuPont warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for the purpose stated in the Directions for Use, subject to the inherent risks described above, when used in accordance with the Directions for Use under normal conditions.

DUPONT MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR OF MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

IN NO EVENT SHALL DUPONT OR SELLER BE LIABLE FOR ANY INCIDENTAL, CONSEQUENTIAL OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. BUYER'S OR USER'S BARGAINED-FOR EXPECTATION IS CROP PROTECTION. THE EXCLUSIVE REMEDY OF THE USER OR BUYER AND THE EXCLUSIVE LIABILITY OF DUPONT OR SELLER, FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY OR CONTRACT, NEGLIGENCE, TORT OR STRICT LIABILITY), WHETHER FROM FAILURE TO PERFORM OR INJURY TO CROPS OR OTHER PLANTS, AND RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT, OR AT THE ELECTION OF DUPONT OR SELLER, THE REPLACEMENT OF THE PRODUCT.

DuPont or its Ag Retailer must have prompt notice of any claim so that an immediate inspection of buyer's or user's growing crops can be made. Buyer and all users shall promptly notify DuPont or a DuPont Ag Retailer of any claims, whether based on contract, negligence, strict liability, other tort or otherwise or be barred from any remedy.

This Limitation of Warranty and Liability may not be amended by any oral or written agreement.

For product information call 1-888-6-DUPONT

© 1999 E. I. du Pont de Nemours and Company, Agricultural Products, Wilmington, Delaware 19898

DRAFT

SUPPLEMENTAL LABELING

DuPont Agricultural Products

ACCENT® HERBICIDE FOR WEED CONTROL IN SWEET CORN FOR PROCESSING IN THE STATES OF DE, MD, NJ AND NY

"..... A Growing Partnership With Nature"

ACCENT® HERBICIDE

EPA REG. NO. 352-560

FOR WEED CONTROL IN SWEET CORN FOR PROCESSING IN THE STATES OF DELAWARE, MARYLAND, NEW JERSEY AND NEW YORK

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

DuPont ACCENT® Herbicide may be applied to sweet corn grown under contract for processing. Consult the ACCENT® label for additional requirements, restrictions and precautions.

WHEN TO APPLY

Applications of ACCENT® may be applied broadcast or with drop nozzles (post-directed) on sweet corn up to 12 inches tall. For sweet corn 12 - 18 inches tall, apply only with drop nozzles. Do not apply to sweet corn taller than 18 inches OR THAT EXHIBITS 6 OR MORE COLLARS (V6), WHICHEVER IS MOST RESTRICTIVE. Only one application of ACCENT® is allowed per year.

HOW TO USE

Apply ACCENT® at 0.66 ounces per acre (or, when packed in water soluble packs, one soluble packet per 4 acres) on processing sweet corn for selective postemergence grass control. See Federal label for weeds controlled.

ACCENT® performs best when applied to actively growing weeds at weed heights no greater than those given in the rate guide. Applications made to weeds larger than the size indicated or to weeds under stress may result in unsatisfactory control.

Adjuvants:

Always add either a crop oil concentrate (1% v/v = 1 gal/100 gal), methylated seed oil (1% v/v = 1 gal/100 gal) or nonionic surfactant (0.25% v/v = 1 qt/100 gal). The addition of an ammonium nitrogen fertilizer such as 28-0-0 (2 - 4 qt per acre), 10-34-0 (1 - 2 qt per acre) or ammonium sulfate (21-0-0) at 2 - 4 lb per acre is also recommended.

Broadcast Application:

Apply ACCENT® using properly calibrated ground equipment only. Use flat fan nozzles at 20 - 40 psi and use a minimum of 10 gal water per acre. Increase both spray volume and pressure as weed density and size increase. Spray equipment must be properly set up to avoid applying an excessive rate directly over the rows and into the corn plant whorl.

Tank Mixes:

Do not tank mix ACCENT® with any other product except those specified on this label. Atrazine can be tank mixed with ACCENT® for improved control of broadleaf weeds. See the Atrazine label for additional information on rates to use and weeds controlled. Do not apply this tank mix to corn taller than 12 inches.

Rotational Crops:


Follow rotational intervals on ACCENT® label.

RESTRICTIONS AND LIMITATIONS

In the state of New York this product is limited to ground application only. Do not apply by air in New York.

ACCENT® may be used only on the following sweet corn hybrids grown under contract for processing:

- Bonus
Excalibur
GG-43
More(GG 55)
Reward
Viking
Zenith



Applying ACCENT® to processing sweet corn under stress may result in crop injury. Stress may be caused by:

- abnormally hot or cold weather
environmental conditions such as drought, water saturated soils, soil compaction, hail damage, or frost
disease, insect, or nematode injury
prior herbicide, or carryover from a previous year's herbicide application.

The user assumes all responsibility associated with applications of ACCENT® to sweet corn.

Do not apply any organophosphate insecticide or any other herbicide within 7 days before or 3 days after an application of ACCENT®.

For product information call 1-888-6-DUPONT

© 1999 E. I. du Pont de Nemours and Company, Agricultural Products, Wilmington, Delaware 19898

Do not apply to sweet corn previously treated with any formulation of Counter¹ soil insecticide. Application of ACCENT® to sweet corn previously treated with Thimet¹, Dyfonate², or Lorsban¹ soil insecticide may cause unacceptable injury and is not recommended. IF A SOIL INSECTICIDE IS REQUIRED, DUPONT FORTRESS SOIL INSECTICIDE, AZTEC³ OR FORCE² ARE THE RECOMMENDED SOIL INSECTICIDES, AS THEY DO NOT INTERACT WITH ACCENT.

Applications within 4 hours of rain may reduce weed control.

A timely cultivation may be necessary to control weeds that emerge after an application of ACCENT®. Do not cultivate within 10 days prior to the application because weed control might be reduced by pruning roots and placing the weed under stress. Optimum timing for cultivation is 7 - 14 days after ACCENT® application.

To avoid subsequent injury to crops other than corn, thoroughly clean all mixing and spray equipment immediately following application. Refer to the label for cleanout procedure.

Do not graze or feed forage, hay, or straw from treated areas to livestock within 30 days of ACCENT® application.

IMPORTANT

BEFORE USING ACCENT®, READ AND FOLLOW ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA-REGISTERED LABEL.

This bulletin contains new or supplemental instructions for use of this product which do not appear on the EPA-registered package label. Follow the instructions carefully.

This labeling must be in the possession of the user at the time of pesticide application.

- ¹ Registered trademark of American Cyanamid Co.
- ² Registered trademark of Zeneca, Inc.
- ³ Registered trademark of Dow Agro Sciences
- ⁴ Registered trademark of Bayer Corporation

(Replaces H-63616 and EPA SLN No. NY-940001)

D-072799

For product information call 1-888-6-DUPONT

© 1999 E. I. du Pont de Nemours and Company, Agricultural Products, Wilmington, Delaware 19898