



AGRICULTURAL PRODUCTS

..... A Growing Partnership With Nature



SUPPLEMENTAL LABELING

LOROX® SP HERBICIDE
TANK MIX WITH 2,4-D (LVE)
FOR USE IN NO-TILL
AND MINIMUM-TILL SOYBEANS

LOROX® SP HERBICIDE

EPA Reg. No. 352-562

TANK MIX WITH 2,4-D (LVE) FOR USE IN NO-TILL AND MINIMUM-TILL SOYBEANS

Du Pont "Lorox" SP Herbicide is recommended for use in a tank mix combination with 2,4-D low volatile ester (LVE) for residual and burndown control of weeds prior to planting soybeans which are directly seeded into a stale seedbed, cover crop, established sod, or in previous crop residues.

Flat fan nozzles that minimize drift are recommended. Use a minimum of 10 gallons total spray volume per acre with ground equipment. Do not apply by air.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Make only one application of 2,4-D (LVE) herbicide per growing season, regardless of the application rate used.

When 2,4-D (LVE) herbicide is applied, do not feed hay, forage, or fodder. Restrict livestock from grazing treated fields. Livestock should be restricted from feeding/grazing of treated cover crops.

HOW TO USE

A tank mix of "Lorox" SP plus 2,4-D (LVE) may be applied to control existing weeds prior to planting soybeans. Refer to the "Lorox" SP and 2,4-D (LVE) labels for use rates, weeds controlled, use instructions, limitations and restrictions.

Refer to the 2,4-D (LVE) herbicide label for precautionary statements, restrictions, and mixing instructions. Read the crop injury statements on the 2,4-D (LVE) herbicide label and use only if these statements are acceptable.

Refer to the "Lorox" SP and the 2,4-D (LVE) labels for additional weeds controlled, precautions, restrictions, limitations, and sprayer cleanup information

Observe all cautions and limitations on labeling of all products used in mixtures.

2,4-D (LVE) USE RATES WHEN TANK MIXED WITH "LOROX" SP FOR PRE-PLANT APPLICATIONS TO SOYBEANS

Table with 2 columns: 2,4-D (LVE) RATE and MINIMUM PREPLANT INTERVAL. Rows show 1/2 p/A (7 Days) and 1 p/A (30 Days).

Note: The application rate of 2,4-D (LVE) herbicide is based on, but not restricted to, a product containing 3.8 pounds acid equivalent (a.e./gallon). The equivalent rates in pounds acid equivalent would be:

1/2 p/A = 0.24 lb a.e./A and 1 p/A = 0.48 lb a.e./A

2,4-D (LVE) herbicide is Isooctyl (2-ethylhexyl) ester of 2,4-Dichlorophenoxyacetic acid.

IMPORTANT

BEFORE USING "LOROX" SP, READ AND CAREFULLY OBSERVE THE CAUTIONARY STATEMENTS AND ALL OTHER INFORMATION APPEARING ON THE PRODUCT LABEL.

This bulletin contains new or supplemental instructions for use of this product which may not appear on the package label. Follow the instructions carefully.

This labeling must be in possession of the user at the time of pesticide application.

D-012293

