

PM 25 352-563 / 1018

U.S. ENVIRONMENTAL PROTECTION AGENCY OFFICE OF PESTICIDES PROGRAMS REGISTRATION DIVISION (WH-567) WASHINGTON, D.C. 20460	EPA REGISTRATION NO.	DATE OF ISSUANCE
	TERM OF ISSUANCE	CFR - 9 1993
NOTICE OF PESTICIDE: <input checked="" type="checkbox"/> REGISTRATION <input type="checkbox"/> REREGISTRATION (Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended)		
NAME OF PESTICIDE PRODUCT DuPont Classic/Pinnacle Herbicide Premix		

NAME AND ADDRESS OF REGISTRANT (Include ZIP code)

Ms. Priscilla L. Friedman  
 E.I. DuPont de Nemours & Company, Inc.  
 Walkers Mill, Barley Mill Plaza  
 P.O. Box 89036  
 Wilmington, DE 19880-0038

**NOTE:** Changes in labeling formula differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above U.S. EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby Registered/Reregistered under the Federal Insecticide, Fungicide, and Rodenticide Act.

A copy of the labeling accepted in connection with this Registration/Reregistration is returned herewith.

Registration is in no way to be construed as an indorsement or approval of this product by this Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA sec. 2(c)(2)(A) provided that you:

1. Submit the data listed below, conducted in accordance with the 40 CFR part 158 test guidelines within six months from the date of this letter.

2. Submit a generalization study with regard to herbicidal activity and mode of action of test material in accordance with the following:

3. Submit data and all data required for the registration or reregistration of your product under FIFRA sec. 2(c)(5) and sec. 4 when the Agency requires all registrants of similar products to submit such data.

4. Make the following label changes listed below before you release the product for shipment:

- a. Add the phrase "EPA Registration No. 352-563"
- b. Add "Get Medical Attention" to the "Caution" Statement of Precautionary Statement.

**BEST AVAILABLE COPY**

VKW

ATTACHMENT IS APPLICABLE

SIGNATURE OF APPROVING OFFICIAL \_\_\_\_\_ DATE \_\_\_\_\_

2078

4. Submit five(5) copies of your final printed labeling before you release the product for shipment. Refer to the A-79 Enclosure for a further description of final printed labeling.

5. If these conditons are not complied with, the registration will be subject to cancelation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

PM 25 352-563 3078



**CLASSIC®/PINNACLE®  
Herbicide Premix**

*Dry flowable*

**For Use on Soybeans Only**

For selective postemergence weed control in soybeans in the state of Illinois.

This product is a water-dispersible granule containing 25% active ingredient by weight contained in a premeasured water soluble pack.

<i>Active Ingredients:</i>	<i>By Weight</i>
Methyl 3-[[[(4-methoxy-6-methyl-1,3,5-triazin-2-yl)amino]carbonyl]amino]sulfonyl]-2-thiophenecarboxylate	12.5%
Chlorimuron Ethyl Ethyl 2-[[[(4-chloro-6-methoxypyrimidin-2-yl) amino]carbonyl]amino]sulfonyl]benzoate	12.5%
<b>INERT INGREDIENTS:</b>	75.0%
<b>TOTAL</b>	<b>100.0%</b>

EPA Reg. No. XXXXXXXX

U.S. Patents 4,547,215, 4,481,029 and 4,394,506

APR 19 1990

FEB - 9 1990

352-563

*Keep Out of Reach of Children*

**CAUTION**

*Precautionary Statements*

*Hazards to Humans and Domestic Animals*

**Caution!** Causes moderate eye irritation. Harmful if absorbed through skin. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling.

*Statement of Practical Treatment*

*If in eyes:*

Flush eyes with plenty of water. Call a physician if irritation persists.

*If on skin:*

Wash with plenty of soap and water.

For medical emergencies involving this product, call toll-free 1-800-441-3637.


*Environmental Hazards*

Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater. Do not apply where/when conditions favor runoff.

**DIRECTIONS FOR USE**

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

**For Use on Soybeans Only**



**BEST AVAILABLE COPY**

**APPLICATION INFORMATION**

Du Pont "Classic"/"Pinnacle" Herbicide Premix is a convenient formulation designed for use by the custom applicator. The dispersible granule formulation readily dissolves in water.

**Rate**  
Apply "Classic"/"Pinnacle" at a rate of 1/2 ounce per acre for selective postemergence broadleaf weed control. Tank mixing with a postemergent grass herbicide such as DuPont Assure® II will provide selective control of most broadleaf weeds and grasses in soybeans.

**When to Apply**

"Classic"/"Pinnacle" performs best when applied to small weeds a few weeks after planting.

**Timing to Crop Stage**

- "Classic"/"Pinnacle" may be applied to soybeans any time after the first trifoliolate has expanded fully.
- Apply no later than 60 days before harvest.

**Timing to Weeds**

- Apply "Classic"/"Pinnacle" when weeds are young and actively growing (after the first true leaves have expanded, but before the weeds exceed the size indicated in the table below).
- Applications made to weeds that are in the cotyledon stage, larger than the size indicated, or to weeds under stress may result in unsatisfactory control.

When applied as directed, "Classic"/"Pinnacle" will control the following weeds:

Height(Inches) at Application

	1	2	3	4	5	6	7	8	9	10	11	12	13
<b>Weeds Controlled</b>													
Annual Smartweeds 2 - 8"													
Coclebar 2 - 4"													
Jimsonweed 2 - 5"													
Lambquarters 2 - 4"													
Pigweed species 2 - 12"													
Velvetleaf 2 - 8"													
Wild Mustard up to 4" in dia.													
Wild Sunflower 2 - 8"													
<b>Weeds Suppressed</b>													
Morningglory species													
Entireleaf 1 - 2"													
Ivyleaf 1 - 2"													
Pitted 1 - 2"													
Smallflower 1 - 2"													
Tall 1 - 2"													
Common Ragweed 1 - 3"													

\* Weed suppression is a visual reduction in weed competition (reduced population, size, and/or vigor) as compared to an untreated area. Degree of control can be increased by treating weeds when actively growing and not under stress.

**Spray Additives**

Applications of "Classic"/"Pinnacle" must include a non-ionic surfactant or crop oil concentrate, and an ammonium nitrogen fertilizer.

**Nonionic Surfactant**

Apply at the rate (concentration) of 0.125% - 0.25% v/v (1-2 pt per 100 gal of spray solution).

- Use of the higher rate of nonionic surfactant, particularly under hot humid conditions, may result in temporary crop injury.
- Use only EPA-approved surfactants containing at least 80% active ingredient authorized for use on food crops.
- Do Not Use Dash<sup>1</sup>.

**Crop Oil Concentrate**

Under dry conditions or during cool weather, a crop oil concentrate at 0.5% v/v (4 pt/100 gal of spray solution) may be used in place of a nonionic surfactant to enhance weed control.

- Use a petroleum-based crop oil concentrate (0.5% v/v) with at least 15% emulsifiers/surfactant.
- The use of crop oil concentrate may result in temporary crop injury.

**Ammonium Nitrogen Fertilizer**

An ammonium nitrogen fertilizer is required in addition to a surfactant or a crop oil concentrate.

Use a high-quality liquid nitrogen fertilizer such as 28-0-0 at a rate of 2-4 quarts per acre, or 10-34-0 at a rate of 1-2 quarts per acre.

- Use the lower rate for spray volumes less than 15 gallons per acre.
- Alternately, a high-quality, sprayable grade of ammonium sulfate (21-0-0) may be used at a rate of 2-4 pounds per acre.

**Mixing Instructions**

One container of "Classic"/"Pinnacle" Herbicide Premix for custom applicators contains 7 1/2 pounds of product. This is enough to treat 240 acre at the rate of 1/2 ounce product per acre (applying the equivalent of 1/4 ounce of "Classic" plus 1/4 ounce of "Pinnacle" per acre). It is designed to be used in combination with Du Pont "Assure" II minibulk system, which contains 15 gallons of "Assure" II Herbicide (sufficient to treat 240 acres at a rate of 8 fluid ounces per acre).

1. Measure out the proper amount of "Classic"/"Pinnacle" Herbicide Premix to apply 1/2 ounce per acre.
2. Fill the tank 1/4 to 1/3 full of water.
3. While agitating, add the required number of ounces of "Classic"/"Pinnacle" Herbicide Premix.

4. The "Classic"/"Pinnacle" Herbicide Premix should dissolve completely within few minutes. Continue adequate agitation.

"Classic"/"Pinnacle" Herbicide Premix should be thoroughly mixed with water in the spray tank before adding any other material (e.g. postemergent grass herbicide, surfactant, crop oil concentrate, or nitrogen fertilizer). As the tank is filling and after the "Classic"/"Pinnacle" Herbicide Premix has dissolved in the needed tank mix products such as "Assure" II and the required spray adjuvants.

6. Apply "Classic"/"Pinnacle" Herbicide Premix spray preparation within 24 hours of mixing to avoid product degradation.

7. If the mixture has settled, thoroughly reagitrate before using.

**Tank Mix Applications**

Do not tank mix "Classic"/"Pinnacle" with any other pesticide or spray adjuvant except as specified on this or other supplemental labeling.

"Classic"/"Pinnacle" may be tank mixed with post grass herbicides such as Du Pont's Assure® II Herbicide.

Under certain conditions, "Classic"/"Pinnacle" may reduce the activity of the post grass herbicide. The broadleaf activity of "Classic"/"Pinnacle" will not be affected. Refer to the grass herbicide label for specific use information and precautions.

Include a nonionic surfactant with the tank mix of "Classic"/"Pinnacle" and post grass herbicides such as "Assure" II.

- Surfactant rate (concentration) should be 0.125%-0.25% v/v (1-2 pints per 100 gallons of spray solution).
- Use of a higher rate of nonionic surfactant, particularly under hot, humid conditions, may result in temporary crop injury.
- Do not use "Dash" or crop oil concentrate when tank mixing "Classic"/"Pinnacle" with postemerge grass herbicides.

The amount of "Assure" II to be used in a tank mix with "Classic"/"Pinnacle" will be determined by the grasses present. When applied as directed, a tank mix of "Classic"/"Pinnacle" and "Assure" II will control the following grasses:

**"Classic"/"Pinnacle" + 5 oz of "Assure" II per acre**

Grass	Size (Height) inches
volunteer corn	6 - 18
shattercane	6 - 12
giant foxtail	2 - 8
seedling johnsongrass	2 - 8

**"Classic"/"Pinnacle" + 8 oz of "Assure" II per acre**

Grass	Size (Height) inches
wild proso millet	2 - 6
fall panicum	2 - 6
green foxtail	2 - 4
bristly foxtail	2 - 4
field sandbur	2 - 6
volunteer cereals	2 - 6

**"Classic"/"Pinnacle" + 10 oz of "Assure" II per acre**

Grass	Size (Height) inches
rhizome johnsongrass	10 - 24

Do not tank mix "Classic"/"Pinnacle" with organophosphate insecticides, or apply "Classic"/"Pinnacle" within 14 days before or after an application of an organophosphate insecticide, as severe crop injury may occur.

**APPLICATION EQUIPMENT**

Many crops are highly sensitive to "Classic"/"Pinnacle". All direct or indirect contact (such as spray drift) with crops other than soybeans should be avoided.

**Ground Application (See also Spray Drift)**

**Broadcast Application**

- Use 10-25 gallons of water per acre.
- Use flat fan nozzles at 25-60 psi.
- Do not use flood, hollow cone, rain drop, whirl chamber, or controlled droplet applicator (CDA) type nozzles. Unacceptable crop injury, excessive spray drift, or poor weed control may result.
- For proper spray coverage, adjust the boom and nozzle height according to the specifications listed by the nozzle manufacturer.

**Band Application**

- For band application, use proportionately less spray mixture.
- To avoid crop injury, carefully calibrate the band applicator not to exceed the labeled rate.
- Carefully follow the manufacturer's instructions for nozzle types (flat fan nozzles preferred), nozzle orientation, distance of the nozzles from the crop and weeds, spray volumes, calibration, and spray pressure.
- For additional information on row banders, see Du Pont's bulletin, "Application Accuracy Row Banders."

**Aerial Application (See also Spray Drift)**

- Use nozzle types and arrangements that will provide for optimum spray distribution and maximum coverage at 5 to 10 GPA.
- Do not apply during a temperature-inversion condition, when winds are gusty, or when other conditions will favor poor coverage and/or off-target spray movement.
- Use a minimum of 5 gallons of water per acre.

**ENVIRONMENTAL CONDITIONS AND BIOLOGICAL ACTIVITY**

"Classic"/"Pinnacle" provides best postemergence results when applied to young, actively growing weeds. Degree of control and duration of effect depend on: application rate, weed spectrum, weed size, growing conditions at and following time of treatment, soil moisture, precipitation, and spray adjuvants. Some weeds, such as lambsquarters, are more affected by stress than others. Delay application until stress passes and weeds start to grow again. Treating weeds under stress or large weeds may result in only partial control. Conditions that are conducive to healthy, actively growing plants optimize the performance of "Classic"/"Pinnacle".

- If rain is expected within 1 hour, do not apply "Classic"/"Pinnacle", or weed control may be decreased.
- Applications made during or immediately following periods of abnormally cold weather for soybeans may result in less than satisfactory weed control.
- Poor weed control or crop injury may result from applications made to plants under stress from
- abnormal hot or cold weather,
- growing conditions such as drought or water-saturated soil,
- disease, or
- nematode, insect, or prior herbicide injury.

Delay application until stress passes and weeds and soybeans resume growth. Severe stress from conditions immediately following application may also result in crop injury or poor weed control.

- Applications during periods of hot and humid weather increase the risk of crop injury

Wilting, temporary leaf yellowing, reddened veins, and/or growth retardation of soybeans may follow application of "Classic"/"Pinnacle". The growth retardation is generally in the form of shortened internode spacing. These effects will generally be most evident 5-7 days after application. The soybeans will recover quickly under favorable growing conditions.

"Classic"/"Pinnacle" rapidly inhibits the growth of susceptible weeds, reducing weed competition hours after application. Leaves turn yellow 5-7 days after postemergence application, followed by death of the weed's growing point. Susceptible plants are controlled in 7-21 days. Death of leaf tissue will follow in some species, while others will remain green but stunted and noncompetitive.

**CULTIVATION**

A timely cultivation may be necessary to control suppressed weeds, weeds that were beyond the maximum size at the time of application, or weeds that emerge after an application of "Classic"/"Pinnacle".

- Do not cultivate before, during, or within 7 days after the application.
- Cultivation may decrease weed control by pruning roots and placing the weed under stress.
- The best time to cultivate is approximately 14 days after application.

**CROP ROTATION**

**Rotational Crop Intervals for "Classic"/"Pinnacle"**

The following rotational intervals must be observed when using "Classic"/"Pinnacle":

<i>Crop</i>	<i>Rotational Interval In Months</i>
<i>Soybeans</i>	<i>Anytime</i>
<i>Cereal Grains</i>	<i>3</i>
<i>Ryegrass</i>	<i>3</i>
<i>Field Corn *</i>	<i>9</i>
<i>Cotton</i>	<i>9</i>
<i>Alfalfa</i>	<i>9</i>
<i>Clover</i>	<i>9</i>
<i>Dry Beans</i>	<i>9</i>
<i>Sorghum</i>	<i>9</i>
<i>Peanut</i>	<i>6</i>
<i>Rice</i>	<i>9</i>
<i>Tobacco (transplant)</i>	<i>9</i>
<i>Tomato (transplant)</i>	<i>9</i>

\* When using "Classic"/"Pinnacle" Herbicide, the term "FIELD CORN" is defined to include only that corn grown for grain or silage, popcorn, and seed corn relative to the Rotational Crop Guidelines section of this label.

**Field Bioassay for Other Crops**

All other crops require a successful field bioassay. The suggested interval before initiating a bioassay is 9 months. If the crop you want to plant does not appear in the table above, you must complete a successful field bioassay during the season before planting. The field bioassay will detect small quantities of

"Classic"/"Pinnacle" that can remain in the soil and could potentially injure rotational crops. A successful field bioassay means growing to maturity a test strip of crop(s) intended for production the following year. The test strip should cross the whole field, including knolls and low areas.

### SPRAYER PREPARATION AND CLEANUP

It is important that spray equipment is clean and free of previous pesticide deposits before using "Classic"/"Pinnacle", and then properly cleaned out following application. Clean all application equipment before applying "Classic"/"Pinnacle". Follow the cleanup procedures specified on the label of the product previously sprayed. If no cleanup procedure is provided, use the procedure that follows. Immediately following applications of "Classic"/"Pinnacle", thoroughly clean all mixing and spray equipment to avoid applications to crops other than soybeans.

#### Notes

- When cleaning spray equipment before applying "Classic"/"Pinnacle", read and follow label directions for proper rinsate disposal of the product previously sprayed.
- A steam cleaning of aerial spray tanks is recommended to dislodge any visible pesticide deposits.
- When spraying or mixing equipment will be used over an extended period to apply multiple loads of "Classic"/"Pinnacle", partially fill the tank with fresh water at the end of each day of spraying, flush the boom and hoses, and allow to sit overnight.

#### Cleanup Procedure

1. Drain the tank and thoroughly hose down the interior surfaces. Flush the tank, hoses, and boom with clean water for a minimum of 5 minutes.
2. Partially fill the tank with clean water and add one gallon of household ammonia\* (containing 3% active) for every 100 gallons of water. Finish filling the tank with water, then flush the cleaning solution through the hoses, boom, and nozzles. Add more water to completely fill the tank and allow to agitate/recirculate for at least 15 minutes. Again, flush the hoses, boom, and nozzles with the cleaning solution, then drain the tank.
3. Repeat Step 2.
4. Remove the nozzles and screens and clean separately in a bucket containing the cleaning agent and water.
5. Thoroughly rinse the tank with clean water for a minimum of 5 minutes, flushing the water through the hoses and boom.

\* Equivalent amounts of an alternate strength ammonia solution may be used, or a tank cleaner recommended in the DuPont bulletin "Sulfonylurea Herbicides. A Guide to Equipment Cleanup."

### SPRAY DRIFT

Do not allow spray from either ground or aerial equipment to drift onto adjacent crops or land, as injury to other plants may occur. The minimization of pesticide drift is the responsibility of the applicator. When spraying adjacent to crops other than soybeans, minimize the spray drift potential by following these practices:

- Stop spraying if wind speed becomes excessive. Spray drift can occur at wind speeds less than 10 MPH. Do not spray if sensitive crops are downwind. Do not spray if winds are gusty.
- High temperatures, drought, and low relative humidity increase the possibility of spray drift. Regardless of wind speed extreme caution must be used when these conditions are present and sensitive crops or plants are nearby.
- Do not apply when a temperature inversion exists. An inversion is characterized by low air movement and an increase in air temperature with an increase in altitude. In humid regions, a fog or mist may form. An inversion may be detected by producing a smoke column and checking for a layering effect. Smoke-producing devices on aircraft are recommended. If not sure whether inversion conditions are present, consult with local weather services before making an application.

To further reduce drift from aerial or ground equipment:

1. Use large-droplet size sprays to minimize drift. Do not use hollow-cone nozzles on ground equipment. Do not use nozzles that produce very small droplets, such as airblast-type nozzles. For ground applications, nozzles should be oriented at an angle between straight down and straight back. For aerial applications, orient nozzles straight back along the windstream. If using flood-type nozzles on aircraft, orient the nozzles so spray is produced in direction of the airstream.
2. Increase volume of spray mix per acre (for example, 10 GPA by air, 25 GPA by ground) by using higher-flow-rate nozzles.
3. Reduce pressure (PSI). Do not exceed 60 psi when applying "Classic"/"Pinnacle". (Vehicle speed must also be reduced to maintain the proper application rate per acre.) Consult manufacturers' catalogs for details on correct calibration.
4. Apply as close to target plants as possible while still maintaining a good spray pattern.

BEST COPY AVAILABLE

8078

### IMPORTANT PRECAUTIONS

- "Classic"/"Pinnacle" Herbicide should be used only in accordance with recommendations on this label or in separately published Du Pont recommendations.
- Injury to or loss of desirable trees or vegetation may result from failure to observe the following:
  - Do not apply or drain or flush equipment on or near desirable trees or other plants, or on areas where their roots may extend, or in locations where the chemical may be washed or moved into contact with their roots.
  - Do not use on lawns, walks, driveways, tennis courts or similar areas.
  - Prevent drift of spray to desirable plants.
- Do not contaminate any body of water.
- Thoroughly clean application equipment immediately after use. (Refer to Sprayer Preparation and Cleanup section of this label.)
- Do not graze or feed forage, hay or straw from treated areas to livestock.
- Do not apply this product through any type of irrigation system.
- Du Pont will not be responsible for losses or damages resulting from the use of this product in any manner not specifically recommended by Du Pont. "Classic"/"Pinnacle" rapidly inhibits the growth of susceptible weeds, reducing weed competition hours after application. Leaves turn yellow 5-7 days after postemergence application, followed by death of the weed's growing point. Susceptible plants are controlled in 7-21 days. Death of leaf tissue will follow in some species, while others will remain green but stunted and noncompetitive.

### STORAGE AND DISPOSAL

**Storage:** Store product only in original container, away from other pesticides, fertilizer, food or feed.

**Product Disposal:** Do not contaminate water, food or feed by storage or disposal. Waste resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**Container Disposal:** Triple rinse (or equivalent), then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

**Notice to Buyer:** Purchase of this material does not confer any rights under patents of countries outside of the United States.

### NOTICE OF WARRANTY

Du Pont warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for purposes stated on such label only when used in accordance with directions under normal use conditions. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Du Pont. In no case shall Du Pont be liable for consequential, special or indirect damages resulting from the use or handling of this product. All such risks shall be assumed by the buyer.

**DU PONT MAKES NO WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESSED OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.**

1, Registered Trademark of BASF

D -090292/092192



© 1992 E.I. du Pont de Nemours and Company  
Agricultural Products, Wilmington, Delaware 19898

