



DuPont Agricultural Products

..... A Growing Partnership With Nature



10/2
SUPPLEMENTAL LABELING

BENLATE® SP FUNGICIDE
EUTYPA DIEBACK OF GRAPES
STATE OF CALIFORNIA

BENLATE® SP FUNGICIDE

(EPA Reg. No. 352 - 564)

FOR DISTRIBUTION AND USE ONLY WITHIN CALIFORNIA

FOR PREVENTION OF EUTYPA DIEBACK ON GRAPES IN THE STATE OF CALIFORNIA

DIRECTIONS FOR USE

) It is a violation of federal law to use this product in a manner inconsistent with its labeling.

DuPont "Benlate" SP Fungicide is recommended for wound protection on grapes in California to reduce Eutypa Dieback.

HOW TO USE

Apply "Benlate" SP at 0.2 lbs. per gallon of water to cut or pruned surface or grapevine immediately after cutting and before rain or dew and spores come into contact with fresh wood. Application may be painted or sprayed onto cut surface area. Results are improved by pruning at a time of year when little or no rain is expected soon after treatment.

Note: Do not contaminate water when disposing of equipment washwaters.

IMPORTANT


BEFORE USING "BENLATE" SP, READ AND FOLLOW ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA-REGISTERED LABEL.

This bulletin contains new or supplemental instructions for use of this product which do not appear on the EPA-registered package label. Follow the instructions carefully.

This labeling must be in the possession of the user at the time of pesticide application.

012694

06/23/94



BEST AVAILABLE COPY



**DuPont Agricultural
Products**

"..... A Growing Partnership With Nature"



SUPPLEMENTAL LABELING

**BENLATE® SP FUNGICIDE
ALMONDS
STATE OF CALIFORNIA**

BENLATE® SP FUNGICIDE

(EPA Reg. No. 352 - 564)

**FOR CONTROL OF SCAB IN ALMONDS
IN THE STATE OF CALIFORNIA**

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

DuPont "Benlate" SP Fungicide is recommended for use on almonds in California for control of scab.

CROP	DISEASE	LIMIT /ACRE /CROP	RATE, MINIMUM GALLONAGE	APPLICATION TIMING	LAST APPLICATION (DAYS TO HARVEST)
Almonds	Scab <i>(Cladosporium carpophilum)</i>	48 oz (3 lb)	16-24 oz/A (1-1.5 lb/A) 10 gal/A air	Apply at pink bud. If needed, repeat at full bloom or 10 days later. Repeat again 2 to 5 weeks later.	80
<ul style="list-style-type: none"> Do not use "Benlate" SP alone. Use only in combination with a nonbenzimidazole fungicide. Under severe disease conditions or on disease susceptible varieties, make the second application during half to full bloom. The first and second applications can be made by ground or air. Make ground applications only for the third application. 					

IMPORTANT

BEFORE USING "BENLATE" SP, READ AND FOLLOW ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA-REGISTERED LABEL.

This bulletin contains new or supplemental instructions for use of this product which do not appear on the EPA-registered package label. Follow the instructions carefully.

This labeling must be in the possession of the user at the time of pesticide application.

D-112294/120694

BEST AVAILABLE COPY

ACCEPTED
with COMMENTS
in EPA Letter Dated

JAN 31 1995

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, this pesticide
registered under EPA Reg. No.

352-564