




PM 21

352-564

6/3/97  
SUPPLEMENTAL LABELING 1/1

DuPont Agricultural  
Products



BENLATE® SP FUNGICIDE  
CONTROL OF WHITE MOLD  
(SCLEROTINIA) ON RADISH  
(SEED CROP)

"..... A Growing Partnership With Nature"

**BENLATE® SP FUNGICIDE**

EPA Reg. No. 352-564

**CONTROL OF WHITE MOLD (SCLEROTINIA) ON RADISH (SEED CROP)**

**DIRECTIONS FOR USE**

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

DuPont "BENLATE" SP Fungicide is recommended for the control of white mold (Sclerotinia) on radish grown for seed.

**HOW TO USE**

Apply 32 oz of "BENLATE" SP per acre. Apply at first petal fall. Repeat at 14 day intervals. Add a spreader sticker to aid in wetting plants. For aerial application use a minimum of 5 gallons per acre.

Do not graze livestock in treated areas.

Do not use treated seed or plant parts for food or feed.

Do not apply more than 96 oz (6 lbs) of "BENLATE" SP per acre per crop.

**IMPORTANT**

**BEFORE USING "BENLATE" SP, READ AND FOLLOW ALL APPLICABLE DIRECTIONS, RESTRICTIONS, AND PRECAUTIONS ON THE EPA REGISTERED LABEL.**

This bulletin contains new or supplemental recommendations for use of this product which do not appear on the EPA registered package label. Follow the instructions carefully.

This bulletin must be in the possession of the user at the time of pesticide application.

D-042997

