

352-569

04-15-2010

1
18



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
Washington, D.C. 20460

OFFICE OF
PREVENTION, PESTICIDES
AND TOXIC SUBSTANCES

Rebecca Ashley
E.I. du Pont de Nemours and Company
1007 Market Street
Wilmington, DE 19898

APR 15 2010

Subject: Label Amendment (container disposal)
DuPont UpBeet Herbicide
EPA Reg. No. 352-569
Application Dated March 15, 2010

Dear Ms. Ashley:

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, is acceptable, provided you make the following changes:

1. On page 1, change the text to "UPBEET is **used** for selective postemergence control..."
2. Change the heading from "Inert Ingredients" to "**Other Ingredients**".
3. Change the heading to "PERSONAL PROTECTIVE EQUIPMENT (PPE)".
4. Change the PPE phrase to "If no such instructions for washables **exist**,"
5. Change the Environmental Hazards phrase to "equipment washwaters or rinsate."
6. It is suggested that the resistance-management grouping symbols be placed on the front panel of the label as described in PR Notice 2001-5.
7. On pages 2 & 6, change the text to read "DuPont UPBEET herbicide **must** be used only in accordance with the **instructions** on this label, or in separately published DuPont **instructions**."
8. On page 2 under Rate, change the text to read "**Make sequential applications** 5 to 10 days apart or as weeds germinate."
9. On page 3, change the text to read "Follow all manufacturer's label **instructions** for the companion product. If these **instructions** conflict with this label, do not tank mix with UPBEET."

- 10. Change the Pesticide Disposal statement to read "Wastes resulting from the use of this product **must** be disposed of on site or at an approved waste disposal facility."
- 11. Add the phrase "TO THE FULLEST EXTENT PERMITTED BY LAW" directly in front of the statements "DUPONT MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY...", "IN NO EVENT SHALL DUPONT OR SELLER BE LIABLE...", and "THE EXCLUSIVE REMEDY OF THE USER OR BUYER..."

The supplemental labels have past their two year grace periods for incorporation into the main label. **You must incorporate the three supplemental labels within ninety (90) days or provide an explanation as to why you believe they are inappropriate for incorporation into the main label.**


A stamped copy of your label is enclosed for your records. This label supercedes all previously accepted labels. You must submit one (1) copy of the final printed label before you release the product for shipment. Products shipped after eighteen (18) months from the date of this letter or the next printing of the label, whichever occurs first, must bear the new revised label. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA §6(e). Your release for shipment of the product constitutes acceptance of these conditions.

Sincerely,


Jimindy Ondish for

Jim Tompkins
Product Manager 25
Herbicide Branch
Registration Division (7505P)

H-



DuPont™ UpBeet®
herbicide



DRAFT LABEL

DUPONT™ UPBEET® HIGHLIGHTS

- For selective postemergence control of broadleaf and grass weeds in sugar beets, including those grown for seed.
- For best results, use UPBEET® with "Betamix" in a tank mix starting with the first application and followed by a minimum of one sequential application separated by 5-10 days. "Betanex" or "Betamix" Progress may also be used. See "Tank Mix Options".
- May be applied by ground (broadcast or band) or air. Use a minimum of 10 gallons of water per acre on a broadcast basis for ground application. For aerial application, use a minimum of 5 gallons of water per acre.
- UPBEET® can be safely applied to sugar beets anytime after planting.
- Apply UPBEET® to actively growing, small seedling weeds. Applications of UPBEET® to weeds under stress may result in less than desirable performance. See "Environmental Conditions".
- Consult label text for complete instructions. Always read and follow label directions for use.

TABLE OF CONTENTS

Precautionary Statements	1
Application Information	2
Rate	2
Broadcast Applications	2
Band Applications	2
Crop Stage at Application	2
Weed Stage at Application	3
Weeds Controlled	3
Specific Weed Problems	3
Tank Mix Options	3
Spray Additives	3
Mixing Instructions	4
Application Equipment	4
Ground	4
Aerial	4
Cultivation	4
Environmental Conditions and Biological Activity	4
Information on Resistant Weeds	4
Integrated Pest Management	5
Crop Rotation	5
Grazing	5
Sprayer Preparation and Cleanup	5
Spray Drift Management	5
Important Precautions	6
Storage and Disposal	7
Notice to Buyer	8
Limitation of Warranty and Liability	8



DuPont™ UpBeet®

herbicide

Dry Flowable

For Weed Control in Sugar Beets

UPBEET® is recommended for selective postemergence control, or partial control, of broadleaf and grass weeds in sugar beets.

This product is a water dispersible granule containing 50% active ingredient by weight.

<i>Active Ingredient</i>	<i>By Weight</i>
Triflurosulfuron methyl: Methyl 2-[[[[[4-(dimethylamino)-6-(2,2,2-trifluoroethoxy)-1,3,5-triazin-2-yl]-amino]carbonyl]amino]sulfonyl]-3-methylbenzoate	50%
<i>Inert Ingredients</i>	50%
TOTAL	100%

EPA Reg. No. 352-569 EPA Est. No. _____

Nonrefillable Container

Net: _____

OR

Refillable Container

Net: _____

ACCEPTED
with COMMENTS
in EPA Letter Dated

APR 15 2010

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

352-569

KEEP OUT OF REACH OF CHILDREN CAUTION

FIRST AID

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-441-3637 for emergency medical treatment information.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION! Avoid contact with skin, eyes and clothing. In case of contact with eyes, immediately flush with plenty of water. Get medical attention if irritation persists.

For medical emergencies involving this product, call toll free 1-800-441-3637.

PERSONAL PROTECTIVE EQUIPMENT

Applicators and other handlers must wear:

Long-sleeve shirt and long pants.

Chemical Resistant Gloves Category A (such as butyl rubber, natural rubber, neoprene rubber or nitrile rubber), all ≥14-mls.

Shoes plus socks.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

USERS SHOULD: Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater.

5
18

6
18

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

DuPont™ UPBEET® herbicide should be used only in accordance with the recommendations on this label, or in separately published DuPont recommendations.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 4 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

Coveralls.

Chemical Resistant Gloves Category A (such as butyl rubber, natural rubber, neoprene rubber or nitrile rubber), all ≥14 mls.

Shoes plus socks.

For best results on the broadest spectrum of weeds, use a minimum of 2 sequential applications of UPBEET® tank mixed with "Betamix", "Betanex" or "Betamix" Progress may also be used. Treat small weeds beginning with the first application. Apply UPBEET® at a broadcast rate of 1/2 - 1 ounce per acre in a tank mix with "Betanex", "Betamix", or "Betamix" Progress for control or partial control of the weeds listed. (See Tank Mix Options.) Use higher rates as weed size or population increases.

Applications should be made 5 to 10 days apart or as weeds germinate. Use a close sequential application if first application was on weeds with 4 leaves.

With an adjuvant, some weeds are controlled with UPBEET® alone (see Weeds Controlled). For best results on these weeds, use a minimum of two sequential applications 5-10 days apart or as weeds germinate.

BAND APPLICATIONS

Dosage Chart for UPBEET® Band Applications

Band width	Band Rates* according to Row Spacing		
	22" rows	30" rows	42" rows
7"	1/6 oz/A	1/8 oz/A	1/12 oz/A
11"	1/4 oz/A	1/5 oz/A	1/7 oz/A

* Equivalent 1/2 oz/Acre Broadcast.

Use proportionately more for higher broadcast rate equivalents.

Wider band widths may result in better overall weed control on fields with high weed populations.

See DuPont guide – "Application Accuracy Row Banders."

Refer to the tank mix partner label for any applicable band rate charts.

See the Broadcast Application section of this label for additional application information.

CROP STAGE AT APPLICATION

UPBEET® can be safely applied to sugar beets any time after planting.

Chlorosis (yellowing) may be observed following an application of UPBEET®. This effect is temporary.

Applications of UPBEET® to crops under stress (such as stress due to lack of moisture, hail damage, flooding, herbicide injury, mechanical injury, or widely fluctuating temperatures) may result in crop injury. Before making applications of UPBEET® to crops previously under stress, or injured from other pesticide applications, the crop needs to be fully recovered and growing vigorously.

Apply no later than 60 days before harvest. Tank mix partners have different preharvest intervals. Always use the most restrictive interval when tank mixing.

APPLICATION INFORMATION

FOR USE ON SUGAR BEETS INCLUDING THOSE GROWN FOR SEED

RATE

BROADCAST APPLICATIONS

The maximum use rate is 2.5 ounces UPBEET® per acre per growing season.

WEED STAGE AT APPLICATION

The growth stage of weeds at application is very important for satisfactory control. For best results, apply to small, emerged weeds between the cotyledon and two true leaf stage.

Weeds should be actively growing and not under stress. Applications made to larger weeds or to weeds under stress may result in unsatisfactory control. See Environmental Conditions and Biological Activity.

Since DuPont™ UPBEET® has little to no soil activity, only weeds that have emerged above the soil surface will be controlled. Use sequential tank mix applications to control new weed flushes.

WEEDS CONTROLLED

with the tankmix of UPBEET® and "Betamix"

Buckwheat, wild	Polygonum convolvulus
Burclover, California (a)	Medicago polymorpha
Chickweed, common	Stellaria media
Dock, curly	Rumex crispus
Fiddleneck, coast	Amsinckia intermedia
Goosefoot, nettleleaf (a)	Chenopodium murale
Groundcherry, Wright (a)	Physalis wrightii
Knotweed, silversheath (b)	..	Polygonum argyrocoleon
Kochia (c)	Kochia scoparia
Lambsquarters, common	Chenopodium album
London rocket (a)	Sisymbrium irio
Mallow, common	Malva neglecta
Mallow, little (a)	Malva parviflora
Mustard, black (a)	Brassica nigra
Mustard, wild (a)	Brassica kaber
Nightshade, black	Solanum nigrum
Nightshade, hairy (seedling only)	Solanum sarrachoides
Radish, wild	Raphanus sativus
Pigweed, prostrate (b)	Amaranthus blitoides
Pigweed, redroot (b)	Amaranthus retroflexus
Ragweed, common	Ambrosia artemisiifolia
Shepherd's-purse (a)	Capsella bursa-pastoris
Smartweed, Pennsylvania	Polygonum pensylvanicum
Sowthistle, annual (a)	Sonchus oleraceus
Velvetleaf (a) (b)	Abutilon theophrasti

(a) In California, UPBEET® + adjuvant will control these weeds, see Spray Additives.

(b) Minimum of 2 applications at higher rates needed for acceptable control in CA.

(c) See "Information on Resistant Weeds".

PARTIAL CONTROL**

with the tankmix of UPBEET® and "Betamix"

Foxtail, green	Setaria viridis
Foxtail, yellow	Setaria lutescens
Junglerice	Echinochloa colonum
Purslane, common	Portulaca oleracea
Smartweed, ladysthumb	Polygonum persicaria

**Partial control: A visual reduction of weed competition (reduced population or size) as well as a significant loss of vigor for individual weed plants.

SPECIFIC WEED PROBLEMS

Wild buckwheat- Apply to cotyledon to 2 leaf stage.

Mallow species- Apply to cotyledon to 1 leaf stage. Three applications may be required to control multiple flushes. Larger sizes may only be suppressed.

Velvetleaf- Apply to cotyledon to 1 leaf stage. Using UPBEET® + adjuvant may give best control. Three applications may be required to control multiple flushes. Larger sizes may only be suppressed.

TANK MIX OPTIONS

UPBEET® may be tank mixed with other suitable registered herbicides to control weeds in addition to those listed.

UPBEET® can also be mixed with other suitable registered fungicides, and insecticides labeled for use on sugarbeets.

Read all label precautions for tank mix partners prior to use. Follow all manufacturer's label recommendations for the companion product. If these recommendations conflict with this label, do not tank mix with UPBEET®.

"Betamix", "Betanex" or "Betamix" Progress- Use with UPBEET® in a minimum of two sequential applications. Local recommendations for product choice, sequence and rate should be followed.

Velvetleaf and Wild radish control may be reduced when UPBEET® is mixed with "Betamix".

"Stinger"- UPBEET® may be mixed in a two way tank mix with "Stinger" for additional broadleaf weed control. Use of an adjuvant is required in this tank mix and crop oil is the preferred adjuvant. (See Spray Additives). Using an UPBEET® + "Betamix" + "Stinger" tank mix does not require an adjuvant.

"H273"- Tank mixes of "H273" and UPBEET® should only be used in areas where crop tolerance to "H273" is acceptable or crop injury may result.

Postemergence Grass herbicides- Tank mixes of UPBEET® with postemergence grass herbicides may result in reduced grass control. If grass control is reduced, an additional application of the grass herbicide may be required. For optimum grass control, apply grass herbicides 24 hours prior to or 5 days after UPBEET® mixtures.

SPRAY ADDITIVES

If UPBEET® is to be applied alone or in a tank mix with "Stinger", an adjuvant must be included. Use a DuPont recommended non-ionic surfactant or crop oil. More information on adjuvant selection may be found in the bulletin "Approved Adjuvants for Use With DuPont Row Crop and Cereal Herbicides".

Sufficient adjuvant is available for UPBEET® if a minimum of 1.5 pts of "Betanex", "Betamix", or "Betamix" Progress is included in the spray mixture.

A spray adjuvant may be added to the tank mixture of UPBEET® plus "Betamix" or "Betamix" Progress in

Michigan and Ohio, after the sugar beets have 4 true leaves (2 leaf pairs).

Non-ionic Surfactant

Apply at the rate (concentration) of 0.25 % v/v (2 pts per 100 gal of spray solution).

Crop Oils

Apply at the rate (concentration) of 1 % v/v (1 gal in 100 gal of spray solution).

MIXING INSTRUCTIONS

SPRAY CARRIER

DuPont™ UPBEET® is to be mixed only in water.

SPRAY PREPARATION

Apply UPBEET® spray preparation within 24 hours to avoid product degradation. When using tank mix partners, follow the most restrictive label.

1. Fill the tank 1/4 to 1/3 full of water.
2. Add UPBEET® with the agitator running. Continue agitation until UPBEET® is fully dispersed, at least 5 minutes. UPBEET® should be thoroughly mixed with water before adding any other material.
3. As the tank is filling, add partners. Add adjuvants last, if needed.
4. Triple rinse all empty containers at this time and add rinsate to spray tank
5. If the mixture is not continuously agitated, settling will occur. If settling occurs, thoroughly re-agitate before using.
6. If UPBEET® and a tank mix partner are to be applied in multiple loads, pre-slurry UPBEET® in clean water prior to adding to the tank. This will prevent the tank mix partner from interfering with the dissolution of UPBEET®.

APPLICATION EQUIPMENT

Some crops are sensitive to UPBEET®. All direct and indirect contact (e.g. spray drift) with crops other than sugar beets should be avoided. See Spray Drift.

Since foliar absorption is the primary means of UPBEET® uptake by plants, thorough spray coverage of weeds is essential. Select a spray volume and delivery system that will ensure thorough coverage and a uniform spray pattern. Weeds shielded from spray by other weeds or sugar beet leaves may not be controlled.

Use 50 mesh or finer strainer and nozzle screens. Make sure screens are clean prior to using UPBEET®. Continuous agitation is required to keep UPBEET® in suspension.

Do not apply UPBEET® through any type of irrigation equipment.

GROUND APPLICATION (SEE SPRAY DRIFT)

Use a minimum of 10 gallons water per acre. Under heavy weed pressure, dense crop foliage, or moisture stress conditions, increase volume an additional 5 gallons per acre. Change nozzle size to increase gallonage.

Broadcast or band applications are recommended. For proper spray coverage, adjust the boom and nozzle height according to equipment manufacturer's specifications. For additional information on row banders, see DuPont's bulletin, "Application Accuracy Row Banders".

Injection systems can be used with UPBEET®. More information is included in the bulletin "Using (liquid) Chemical Injection Systems with UPBEET® herbicide".

AERIAL APPLICATION (SEE SPRAY DRIFT)

Use a minimum of 5 gallons water per acre. Do not apply by air in California except as directed by supplemental labeling.

CULTIVATION

Timely cultivation(s) can be used in addition to UPBEET® tank mixes for optimum weed control in a sugar beet management program.

ENVIRONMENTAL CONDITIONS AND BIOLOGICAL ACTIVITY

UPBEET® provides the best postemergence results when applied to small, actively, growing weeds. UPBEET® rapidly stops growth of susceptible weeds. Weeds turn yellow 7-21 days after postemergence application, followed by death of the growing point.

Conditions that promote the activity of UPBEET® are warm temperatures and adequate soil moisture before, during and immediately after application. Treating large or stressed weeds may result in poor weed control. Delay application until stress passes and weeds begin to grow again. Best results are obtained when applications avoid: injury from previous herbicide applications, cold, dry conditions, stress conditions due to frost, drought or water-saturated soil, disease or insect damage.

Dry, dusty field conditions may reduce weed control in wheel track areas. Higher volumes and wider band widths may improve control in these conditions.

A vigorously growing crop will aid weed control by shading and providing competition for weeds. In areas of thin stand or seeding skips, additional flushes of weeds may occur.

Rainfall within 6 hours may reduce weed control.

INFORMATION ON RESISTANT WEEDS

Naturally occurring weed biotypes* that are resistant to DuPont™ HARMONY® EXTRA herbicide or DuPont™ EXPRESS® herbicide will also be resistant to UPBEET®.

If resistant weed biotypes, such as kochia, are suspected or known to be present, spray UPBEET® in tank mixtures with other broadleaf herbicides having a different mode of action**. Adjust the use rate of the tank mix partner so that it alone will control the resistant biotypes.

Several strategies can delay the development of resistant weed biotypes:

1. Use preplant incorporated or preemergence herbicides for weed control prior to postemergence use of DuPont™ UPBEET®.
2. Application of UPBEET® in a tank mix with another mode of action herbicide, such as "Betamix".
3. The use of cultivation and/or hand weeding to control escapes.
4. Improve control of these species in rotational crops.

* Biotypes are naturally-occurring individuals of the species which have a slightly different genetic makeup. Resistant biotypes may look exactly the same as susceptible biotypes. Herbicide-resistant biotypes are able to survive a use rate several times higher than needed to control susceptible biotypes.

** Mode of action is the chemical interaction that disrupts plant growth and development.

DuPont recommends the use of Integrated Pest Management (IPM) programs to control pests. This product may be used as part of an Integrated Pest Management (IPM) program which can include biological, cultural, and genetic practices aimed at preventing economic pest damage. Application of this product should be based on IPM principles and practices including field scouting or other detection methods, correct target pest identification, population monitoring, and treating when target pest populations reach locally determined action thresholds. Consult your state cooperative extension service, professional consultants or other qualified authorities to determine appropriate action treatment threshold levels for treating specific pest/crop or site systems in your area.

CROP ROTATION

Sugar beets may be replanted anytime after application of UPBEET®.

Any other crop, except corn, may be planted 14 days after the last application of UPBEET®. Corn can be planted 21 days after using UPBEET®.

GRAZING

UPBEET® has no restriction on grazing or feeding of crop residue to livestock. Tank mix partners may have grazing or feeding restrictions, therefore always refer to the label of the tank mix partner and follow the most restrictive label.

SPRAYER PREPARATION AND CLEANUP

It is important that spray equipment is cleaned and free of existing pesticide deposits before using UPBEET®. Follow the cleanup procedures specified on the label of the product previously sprayed. If no cleanout procedure is provided, use the one that follows.

To avoid subsequent injury to other crops, thoroughly clean all mixing and spray equipment immediately following applications of UPBEET®.

AT THE END OF THE DAY

It is recommended when UPBEET® will be applied over several days, at the end of each day, rinse the interior of the tank with fresh water, then partially fill the tank and flush the boom and hoses. This will prevent the buildup of dried pesticide deposits which are difficult to remove from application equipment.

CLEANUP PROCEDURE

1. Drain tank and thoroughly hose down the interior surfaces. Flush the tanks, boom, and hoses with clean water. Loosen and physically remove any visible deposits.
2. Partially fill the tank with clean water and one gallon household ammonia* (contains 3% active ingredient) for every 100 gallons of water. Flush the hoses, boom, and nozzles with the cleaning solution, then add more water to completely fill the tank. Circulate, with agitation, the cleaning solution through the tank and hoses for at least 15 minutes. Flush the hoses, boom, and nozzles again with the cleaning solution and drain the tank.
3. Remove the nozzles and screens and clean separately in a bucket containing cleaning agent and water.
4. Repeat step 2
5. Rinse the tank, boom, and hoses with clean water.
6. The rinsate may be disposed of on site or at an approved waste disposal facility.

* Equivalent amounts of an alternate strength ammonia solutions may be used, or a tank cleaner recommended in the DuPont bulletin- A Guide to Application Equipment Cleanup.

Notes:

1. When UPBEET® is tank mixed with other pesticides, all cleanout procedures should be examined and the most rigorous procedure should be followed.
2. In addition to this cleanout procedure, any pre-cleanout procedures for the next product to be sprayed should be examined and the most rigorous procedure should be followed.
3. Where routine spraying practices include shared equipment frequently being switched between applications of UPBEET® and applications to crops other than sugar beets during the same spray season, it is recommended a sprayer be dedicated to UPBEET® to further reduce the chance of crop injury.

SPRAY DRIFT MANAGEMENT

The interaction of many equipment and weather-related factors determines the potential for spray drift. The applicator is responsible for considering all these factors when making application decisions.

AVOIDING SPRAY DRIFT IS THE RESPONSIBILITY OF THE APPLICATOR.

IMPORTANCE OF DROPLET SIZE

The most effective way to reduce drift potential is to apply large droplets (>150 - 200 microns). The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. The presence of sensitive species nearby, the environmental conditions, and pest pressure may affect how an applicator balances drift control and coverage. **APPLYING LARGER DROPLETS REDUCES DRIFT POTENTIAL, BUT WILL NOT PREVENT DRIFT IF APPLICATIONS ARE MADE IMPROPERLY OR UNDER UNFAVORABLE ENVIRONMENTAL CONDITIONS! See Wind, Temperature and Humidity, and Temperature Inversions sections of this label.**

Controlling Droplet Size - General Techniques

- **Volume** - Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets.
- **Pressure** - Use the lower spray pressures recommended for the nozzle. Higher pressure reduces droplet size and does not improve canopy penetration. WHEN HIGHER FLOW RATES ARE NEEDED, USE A HIGHER-CAPACITY NOZZLE INSTEAD OF INCREASING PRESSURE.
- **Nozzle Type** - Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles.

Controlling Droplet Size - Aircraft

- **Number of Nozzles** - Use the minimum number of nozzles with the highest flow rate that provide uniform coverage.
- **Nozzle Orientation** - Orienting nozzles so that the spray is emitted backwards, parallel to the airstream will produce larger droplets than other orientations.
- **Nozzle Type** - Solid stream nozzles (such as disc and core with swirl plate removed) oriented straight back produce larger droplets than other nozzle types.
- **Boom Length** - The boom length should not exceed 3/4 of the wing or rotor length - longer booms increase drift potential.
- **Application Height** - Application more than 10 ft above the canopy increases the potential for spray drift.

BOOM HEIGHT

Setting the boom at the lowest labeled height (if specified) which provides uniform coverage reduces the exposure of droplets to evaporation and wind. For ground equipment, the boom should remain level with the crop and have minimal bounce.

WIND

Drift potential increases at wind speeds of less than 3 mph (due to inversion potential) or more than 10 mph. However, many factors, including droplet size and equipment type, determine drift potential at any given wind speed. AVOID GUSTY OR WINDLESS CONDITIONS.

Note: Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they affect spray drift.

TEMPERATURE AND HUMIDITY

When making applications in hot and dry conditions, set up equipment to produce larger droplets to reduce effects of evaporation.

TEMPERATURE INVERSIONS

Drift potential is high during a temperature inversion. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain close to the ground and move laterally in a concentrated cloud. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning.

Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

SHIELDED SPRAYERS

Shielding the boom or individual nozzles can reduce the effects of wind. However, it is the responsibility of the applicator to verify that the shields are preventing drift and not interfering with uniform deposition of the product.

AIR ASSISTED (AIR BLAST) FIELD CROP SPRAYERS

Air assisted field crop sprayers carry droplets to the target via a downward directed air stream. Some may reduce the potential for drift, but if a sprayer is unsuitable for the application and/or set up improperly, high drift potential can result. It is the responsibility of the applicator to determine that a sprayer is suitable for the intended application, is configured properly, and that drift is not occurring.

Note: Air assisted field sprayers can affect product performance by affecting spray coverage and canopy penetration. Consult the application equipment section of this label to determine if use of an air assisted sprayer is recommended.

IMPORTANT PRECAUTIONS

DuPont™ UPBEET® should be used only in accordance with recommendations on this label or in separately published DuPont recommendations.

Injury to or loss of desirable trees or vegetation may result from failure to observe the following:

Do not apply, drain or flush equipment on or near desirable trees or other plants, or on areas where their roots may extend, or in locations where the chemical may be washed or moved into contact with their roots. Do not use on lawns, walks, driveways, tennis courts or similar areas. Prevent drift of spray to desirable plants.

Do not contaminate any body of water.

Thoroughly clean application equipment immediately after use (refer to "Sprayer Preparation and Cleanup" section of this label).

UPBEET® is non-corrosive, non-flammable, non-volatile, and does not freeze in storage.

STORAGE AND DISPOSAL

Pesticide Storage: Store product in original container only. Do not contaminate water, other pesticides, fertilizer, food or feed in storage. Store in a cool, dry place.

Pesticide Disposal: Do not contaminate water, food, or feed by disposal. Waste resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Handling: Refer to the Net Contents section of this product's labeling for the applicable "Nonrefillable Container" or "Refillable Container" designation.

Nonrefillable Plastic and Metal Containers (Capacity Equal to or Less Than 50 Pounds):

Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then, for Plastic Containers, offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration. Do not burn, unless allowed by state and local ordinances. For Metal Containers, offer for recycling if available or reconditioning if appropriate, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

Nonrefillable Plastic and Metal Containers (Capacity Greater Than 50 Pounds): Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then, for Plastic Containers, offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration. Do not burn, unless allowed by state and local ordinances. For Metal Containers, offer for recycling if available or reconditioning if appropriate, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

Nonrefillable Plastic and Metal Containers, e.g., Intermediate Bulk Containers [IBC] (Size or Shape Too Large to be Tipped, Rolled or Turned Upside Down): Nonrefillable container. Do not reuse or refill this container. Clean container promptly after emptying the contents from this container into application equipment or mix tank and before final disposal using the following pressure rinsing procedure. Insert a lance fitted with a suitable tank cleaning nozzle into the container and ensure that the water spray thoroughly covers the top, bottom and all sides inside the container. The nozzle manufacturer generally provides instructions for the appropriate spray pressure, spray duration and/or spray volume. If the manufacturer's instructions are not available, pressure rinse the container for at least 60 seconds using a minimum pressure of 30 PSI with a minimum rinse volume of 10% of the container volume. Drain, pour or pump rinsate into application equipment or rinsate collection system. Repeat this pressure rinsing procedure two more times. Then, for Plastic Containers, offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration. For Metal Containers, offer for recycling if available or reconditioning if appropriate, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

Nonrefillable Paper or Plastic Bags, Fiber Sacks including Flexible Intermediate Bulk Containers (FIBC) or Fiber Drums With Liners: Nonrefillable container. Do not reuse or refill this container. Completely empty paper or plastic bag, fiber sack or drum liner by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application or manufacturing equipment. Then offer for recycling if available or dispose of empty paper or plastic bag, fiber sack or fiber drum and liner in a sanitary landfill, or by incineration. Do not burn, unless allowed by state and local ordinances.

Refillable Fiber Drums With Liners: Refillable container (fiber drum only). *Refilling Fiber Drum:* Refill this fiber drum with DuPont™ UPBEET® containing triflurosulfuron methyl only. Do not reuse this fiber drum for any other purpose. Cleaning before refilling is the responsibility of the refiller. Completely empty liner by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application or manufacturing equipment. *Disposing of Fiber Drum and/or Liner:* Do not reuse this fiber drum for any other purpose other than refilling (see preceding). Cleaning the container (liner and/or fiber drum) before final disposal is the responsibility of the person disposing of the container. Offer the liner for recycling if available or dispose of liner in a sanitary landfill, or by incineration. Do not burn, unless allowed by state and local ordinances. If drum is contaminated and cannot be reused, dispose of it in the manner required for its liner. To clean the fiber drum before final disposal, completely empty the fiber drum by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application or manufacturing equipment. Then offer the fiber drum for recycling if available or dispose of in a sanitary landfill, or by incineration. Do not burn, unless allowed by state and local ordinances.

All Other Refillable Containers: Refillable container. *Refilling Container:* Refill this container with DuPont™ UPBEET® containing triflurosulfuron methyl only. Do not reuse this container for any other purpose. Cleaning before refilling is the responsibility of the refiller. Prior to refilling, inspect carefully for damage such as cracks, punctures, abrasions, worn out threads and closure devices. If damage is found, do not use the container, contact DuPont at the number below for instructions. Check for leaks after refilling and before transporting. If leaks are found, do not reuse or transport container, contact DuPont at the number below for instructions. *Disposing of Container:* Do not reuse this container for any other purpose other than refilling (see preceding). Cleaning the container before final disposal is the responsibility of the person disposing of the container. To clean the container before final disposal, use the following pressure rinsing procedure. Insert a lance fitted with a suitable tank cleaning nozzle into the container and ensure that the water spray thoroughly covers the top, bottom and all sides inside the container. The nozzle manufacturer generally provides instructions for the appropriate spray pressure, spray duration and/or spray volume. If the manufacturer's instructions are not available, pressure rinse the container for at least 60 seconds using a minimum pressure of 30 PSI with a minimum rinse volume of 10% of the container volume. Drain, pour or pump rinsate into application equipment or rinsate collection system. Repeat this pressure rinsing procedure two more times. Then, for Plastic Containers, offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration. Do not burn, unless allowed by state and local ordinances. For Metal Containers, offer for recycling if available or reconditioning if appropriate, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

Outer Foil Pouches of Water Soluble Packets (WSP): Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available or, dispose of the empty outer foil pouch in the trash as long as WSP is unbroken. If the outer pouch contacts the formulated product in any way, the pouch must be triple rinsed with clean water. Add the rinsate to the spray tank and dispose of the outer pouch as described previously.

Do not transport if this container is damaged or leaking. If the container is damaged, leaking or obsolete, or in the event of a major spill, fire or other emergency, contact DuPont at 1-800-441-3637, day or night.

NOTICE TO BUYER: Purchase of this material does not confer any rights under patents of countries outside of the United States.

The DuPont Oval Logo, DuPont™, UPBEET®, HARMONY®, and EXPRESS® are trademarks or registered trademarks of E. I. duPont de Nemours & Co. "Betamix", "Betanex" are trademarks of Aventis CropScience "Stinger" is a trademark of Dow Agro Science "H273" manufactured by Elf Atochem North America

D - 1545 030810

LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read This Limitation of Warranty and Liability Before Buying or Using This Product. If the Terms Are Not Acceptable, Return the Product at Once, Unopened, and the Purchase Price Will Be Refunded.

It is impossible to eliminate all risks associated with the use of this product. Such risks arise from weather conditions, soil factors, off target movement, unconventional farming techniques, presence of other materials, the manner of use or application, or other unknown factors, all of which are beyond the control of DuPont. These risks can cause: ineffectiveness of the product; crop injury, or; injury to non-target crops or plants.

DuPont does not agree to be an insurer of these risks. **WHEN YOU BUY OR USE THIS PRODUCT, YOU AGREE TO ACCEPT THESE RISKS.**

DuPont warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for the purpose stated in the Directions for Use, subject to the inherent risks described above, when used in accordance with the Directions for Use under normal conditions.

DUPONT MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR OF MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

IN NO EVENT SHALL DUPONT OR SELLER BE LIABLE FOR ANY INCIDENTAL, CONSEQUENTIAL OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. BUYER'S OR USER'S BARGAINED-FOR EXPECTATION IS CROP PROTECTION. THE EXCLUSIVE REMEDY OF THE USER OR BUYER AND THE EXCLUSIVE LIABILITY OF DUPONT OR SELLER, FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY OR CONTRACT, NEGLIGENCE, TORT OR STRICT LIABILITY), WHETHER FROM FAILURE TO PERFORM OR INJURY TO CROPS OR OTHER PLANTS, AND RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT, OR AT THE ELECTION OF DUPONT OR SELLER, THE REPLACEMENT OF THE PRODUCT.

DuPont or its Ag Retailer must have prompt notice of any claim so that an immediate inspection of buyer's or user's growing crops can be made. Buyer and all users shall promptly notify DuPont or a DuPont Ag Retailer of any claims, whether based on contract, negligence, strict liability, other tort or otherwise or be barred from any remedy.

This Limitation of Warranty and Liability may not be amended by any oral or written agreement.

For product information call: 1-888-6-DUPONT

Internet address: www.dupont.com/ag/us

© 1998-2010- E. I. du Pont de Nemours and Company, Crop Protection, 1007 Market Street, Wilmington, Delaware 19898.

All rights reserved.

13
18

SUPPLEMENTAL LABELING

**DuPont Crop
Protection**

**DUPONT™ UPBEET® HERBICIDE
FOR USE ON
BELGIAN ENDIVE/CHICORY**

DUPONT™ UPBEET® HERBICIDE

EPA Reg. No. 352-569

FOR POSTEMERGENCE WEED CONTROL IN BELGIAN ENDIVE/CHICORY.

DuPont™ UPBEET® herbicide is recommended for selective postemergence control of redroot pigweed, kochia, shepherdspurse, velvetleaf and partial control of prostrate pigweed in Belgian Endive/Chicory.

This product is a water dispersible granule containing 50% active ingredient by weight.

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation. DuPont™ UPBEET® herbicide should be used only in accordance with the recommendations on this label, or in separately published DuPont recommendations.

DIRECTIONS FOR USE

Apply DuPont™ UPBEET® at a broadcast rate of 1/2 ounce per acre, starting when 80% of the Belgian Endive/Chicory plants have at least 1 true leaf developed (2 cotyledons + 1 leaf). For best results on the broadest spectrum of weeds, use a minimum of 2 sequential applications of UpBeet.

Applications should be made 5 to 10 days apart or as weeds germinate. Use a close sequential application if first application is on larger weeds (3-4 leaves).

In some cases, discoloration and even slowed growth may be observed. These symptoms are temporary and have no significant effect on yield.

For best results apply DuPont™ UPBEET® to small actively growing weeds when the temperature is between 40F and 75F. Do not treat when frost is expected in the hours following application. If high temperatures are expected during the day, treatment should be applied early in the morning or in the evening, so that a period of moderate temperatures (lower than 75F) will follow the application.

Apply no later than 60 days before harvest.

The total amount of DuPont™ UPBEET® applied must not exceed 1.5 ounces per acre per growing season.

WEEDS CONTROLLED with DuPont™ UPBEET®

Redroot Pigweed*Amaranthus retroflexus*
Kochia*Kochia scoparia* (L.) Schrad
Shepherdspurse*Capsella bursa-pastoris* (L.) Medicus
Velvetleaf*Abutilon theophrasti* Medicus

PARTIAL CONTROL* with DuPont™ UPBEET®

Prostrate Pigweed*Amaranthus blitoides*

* Partially controlled weeds exhibit a visual reduction in numbers as well as a significant loss of vigor.

WEED STAGE AT APPLICATION

Weeds should be actively growing and not under stress. Applications made to larger weeds or to weeds under stress may result in unsatisfactory control.

14
18

Since DuPont™ UPBEET® has little to no soil activity, only weeds that have emerged above the soil surface will be controlled. Use sequential applications to control new weed flushes. The growth stage of weeds at application is very important for satisfactory control. For best results, apply to small, emerged weeds between the cotyledon and two true leaf stage.

TANK MIX OPTIONS

When applied to Belgian Endive/Chicory, DuPont™ UPBEET® may not be tank mixed with any other herbicide, insecticide, or fungicide unless specified in other labeling.

SPRAY ADDITIVES

Nonionic Surfactant (NIS)

- A tankmix with 0.25 % volume/volume (2 pt per 100 gal of spray solution) of non-ionic surfactant may be added. Addition of surfactant may increase crop response. First time users should limit the use of surfactant to a small test area to determine if crop response is acceptable.
- Surfactant products must contain at least 60% nonionic surfactant with a hydrophilic/lipophilic balance (HLB) greater than 12.

Crop Oils and Other Surfactants

- Use of petroleum crop oil concentrate (COC), methylated seed oils (MSO), silicon surfactants or other wetting agents with DuPont™ UPBEET® used on Belgian Endive/Chicory may result in severe crop injury and are therefore not recommended for use with UpBeet.

IMPORTANT

BEFORE USING THIS PRODUCT, READ AND CAREFULLY NOTE THE CAUTIONARY STATEMENTS AND OTHER PROCEDURAL INFORMATION APPEARING ON THE EPA REGISTERED LABEL OR ON OTHER SUPPLEMENTAL LABELS. FOLLOW ALL APPLICABLE DIRECTIONS, RESTRICTIONS, AND PRECAUTIONS.

This bulletin contains new or supplemental instructions for use of this product which does not appear on the package label. Follow the instructions carefully.

This labeling must be in the possession of the user at the time of pesticide application.

R-290 030603 06-05-03

15
18

SUPPLEMENTAL LABELING

**DuPont Crop
Protection**

**DUPONT™ UPBEET®
HERBICIDE
MICRO-RATE APPLICATIONS
IN SUGARBEETS**

DUPONT™ UPBEET® HERBICIDE

EPA REG. NO. 352-569

MICRO-RATE BROADCAST APPLICATIONS IN SUGARBEETS

GENERAL INFORMATION

Multiple applications of UPBEET® in tank mixture with reduced rates of "Betamix", "Betanex", "Progress", "Stinger", a grass herbicide such as ASSURE® II, and methylated seed oil may be applied by air or ground equipment to sugarbeets to control early germinating weeds.

DIRECTIONS FOR USE

For best results, apply UPBEET® at a minimum rate of 1/8 ounce per acre in a tank mixture with "Betamix", "Betanex" or "Progress", plus 1.5% v/v methylated or modified seed oil adjuvant (1.5 gals of product per 100 gal of spray solution). ASSURE® II or "Stinger" herbicide may be added as needed. ASANA® XL Insecticide may also be added to the mixture as needed.

Make a minimum of three sequential applications, generally at 5-7 day intervals (or as directed by local recommendations). Accurate timing is essential; make initial application immediately after weeds emerge. Broadcast applications are the preferred application method. If weed control is not adequate due to climatic conditions, spray coverage, or other factors, return to conventional application rates as recommended on the EPA approved UPBEET® label.

PRECAUTIONS

• Not all weeds will be adequately controlled, even with favorable climatic conditions. Conventional rates of UPBEET® and/or hand labor may be required if multiple micro-rate applications do not adequately control weeds.

• Plugging of spray nozzles may be encountered due to the potential for formation of a precipitate in the spray solution that is often associated with micro rate applications.

To minimize this precipitate:

- Allow spray water to warm before mixing products.
- Adjust spray tank solution to a pH of 8-9.
- Add a grass control product. Assure II is the preferred product.
- Slurry the UPBEET® in clean, warm water prior to adding to the spray tank. Add UPBEET® first, followed by "Betamix"/"Betanex"/"Progress", then "Stinger", then grass product (such as ASSURE® II), and/or then methylated seed oil.
- Gently agitate the spray solution. Vigorous agitation tends to increase precipitate formation.
- Apply the spray solution as quickly as possible. Start with a clean sprayer, spray until the tank is dry, flush tank and lines between loads with fresh water, especially if any precipitate formation is seen in the tank or on spray tip screens. At the end of the day thoroughly flush tank and lines with fresh water. Don't leave any undiluted spray solution in the tank overnight. Clean tank screens and spray tip screens to remove any precipitate.

RESTRICTIONS

• The addition of methylated seed oils could increase the possibility of crop injury. Refer to the EPA approved "Betamix", "Betanex" or "Progress" label, and any supplemental labeling, for specific rate recommendations and restrictions. If those recommendations conflict with this label, do not follow this micro-rate labeling. Methylated seed oils must not be added if the "Betamix", "Betanex" or "Progress" rate exceeds that specified for each product and specific sugarbeet leaf stage.

• Read and follow all manufacturer's label recommendations (EPA registered labels, supplemental labels, and State local need labels (24C)) for "Stinger", ASSURE® II, ASANA® XL, "Betamix", "Betanex", "Progress", any other pesticides, and adjuvants. If those recommendations conflict with this label, do not follow this microrate labeling.

16
18

**IMPORTANT NOTICE
BEFORE BUYING OR USING DUPONT PRODUCT(S)
COVERED BY THIS LABEL, READ THIS LABEL, AND
THE LABELS FOR ALL PRODUCTS REFERENCED IN
THIS LABEL, IN THEIR ENTIRETY. FOLLOW ALL
APPLICABLE DIRECTIONS, RESTRICTIONS AND
PRECAUTIONS.**

This label contains supplemental instructions for use of the referenced DuPont product. Follow these instructions carefully. This label must be in the possession of the user at the time of application.

Read the Limitation of Warranty and Liability on the Section 3 Federal product label before buying or using THIS product. If terms are not acceptable, return the unopened package at once to Seller for full refund of purchase price paid. Otherwise, use by Buyer or any other User constitutes acceptance of the terms of the Limitation of Warranty and Liability on the Section 3 Federal product label.

“Betamix”, “Betanex” and “Progress” are registered trademarks of Bayer Crop Protection.

“Stinger” is a registered trademark of Dow Agrosciences.

(Replaces H-63622)

R-332 021804 04-22-04

17
18

SUPPLEMENTAL LABELING

DuPont Crop Protection

DUPONT™ UPBEET® HERBICIDE FOR AERIAL APPLICATION TO SUGARBEETS IN THE IMPERIAL VALLEY OF CALIFORNIA

DUPONT™ UPBEET® HERBICIDE

EPA Reg. No. 352-569

FOR AERIAL APPLICATION TO SUGARBEETS IN THE IMPERIAL VALLEY OF CALIFORNIA

DIRECTIONS FOR USE

It is violation of federal law to use this product in a manner inconsistent with its labeling.

DuPont™ UPBEET® herbicide may be applied by air for weed control in sugarbeets grown in Imperial Valley of California, only by aerial applicators certified by DuPont to be in compliance with conditions of the "DuPont UPBEET® Aerial Stewardship Program".

UPBEET® herbicide may be applied by air for weed control in sugarbeets grown in Imperial Valley of California, in accordance with the restrictions described on the label. Consult the Section 3 Federal label for additional requirements, restrictions and precautions.

Before using UPBEET® herbicide by air, be sure that the aircraft spray tank, and all mixing tanks and delivery systems equipment is clean. Apply UPBEET® herbicide at 1/2 to 1 ounce per acre. Add a nonionic surfactant containing at least 80% active ingredient.

Apply in a minimum of 5 gallons of water per acre when weeds are small and actively growing. Weeds that are sheltered from spray by sugarbeet foliage or other larger weeds may not be controlled. UPBEET® can be tank mixed with other suitable sugarbeet herbicides registered for use by air. Consult the EPA registered UPBEET® label for tank mix recommendations.

SPRAYER PREPARATION AND CLEANUP

It is important that spray equipment is cleaned and free of existing pesticide deposits before using UPBEET. Follow the cleanup procedures specified on the label of the product previously sprayed. If no cleanout procedure is provided, use the one that follows.

To avoid subsequent injury to other crops, thoroughly clean all mixing and spray equipment immediately following application of UPBEET®.

CLEANUP PROCEDURE

1. Drain tank and thoroughly hose down the interior surfaces. Flush the tanks, boom, and hoses with water. Loosen and physically remove any visible deposits.
2. Partially fill the tank with clean water and one gallon household ammonia (contains 3% active ingredient) for every 100 gallons of water. Flush the hoses, boom and nozzles with the cleaning solution, then add more water to completely fill the tank. Circulate, with agitation, the cleaning solution through the tank and hoses for at least 15 minutes. Flush the hoses, boom and nozzles again with the cleaning solution and drain the tank.
3. Remove the nozzles and screens and clean separately in a bucket containing cleaning agent and water.
4. Repeat step 2.
5. Rinse tank, boom and hoses with clean water.
6. The rinsate may be disposed of on site or at an approved waste disposal facility.

CROP INJURY THAT RESULTS FROM INADEQUATELY CLEANED SPRAY EQUIPMENT IS THE RESPONSIBILITY OF THE APPLICATOR.

SPRAY DRIFT MANAGEMENT

Avoid Spray drift to adjacent crops as injury may occur. The interaction of many equipment and weather-related factors determines the potential for spray drift. The applicator is responsible for considering all these factors when making application decisions.

Aerial applicators applying UPBEET® herbicide must position at least one person at the application site, with a wind indicator visible to the pilot of the aircraft, at all times during the application. In addition, the personnel at the application site must be able to communicate with the aircraft.

© 2006 E. I. du Pont de Nemours and Company, Crop Protection, Wilmington, Delaware 19898

18
18

Crops which are sensitive to UPBEET® include, but are not limited to Lettuce (Head and Leaf varieties), Broccoli, Cauliflower, Cabbage, Onions, and seedling alfalfa. Applications to fields adjacent to these and all other crops must be made only under conditions which are not likely to result in drift off the treated site.

AVOIDING SPRAY DRIFT IS THE RESPONSIBILITY OF THE APPLICATOR

IMPORTANT

BEFORE USING THIS PRODUCT HERBICIDE, READ AND CAREFULLY NOTE THE CAUTIONARY STATEMENTS AND OTHER PROCEDURAL INFORMATION APPEARING ON THE EPA REGISTERED LABEL OR ON OTHER SUPPLEMENTAL LABELS.

This label contains supplemental instructions for use of the referenced DuPont product which do not appear on the package label. Follow the instructions carefully.

This labeling must be in the possession of the user at the time of pesticide application.

Read the Limitation of Warranty and Liability on the Section 3 Federal product label before buying or using THIS product. If terms are not acceptable, return the unopened package at once to Seller for full refund of purchase price paid. Otherwise, use by Buyer or any other User constitutes acceptance of the terms of the Limitation of Warranty and Liability on the Section 3 Federal product label.

R-630 051006-2 06-12-06