

BASIS HERBICIDE

EPA Reg No. 352-571

FOR USE ON FOOD GRADE AND WAXY CORN HYBRIDS

DIRECTIONS FOR USE

It is a violation fo Federal law to use this product in a manner inconsistent with its labeling.

Basis may be applied to the following types of field corn of greater than 88 day Relative Maturity (RM): "food grade" (yellow dent field corn with hard endosperm) and waxy.

Not all "food grade" or waxy field corn hybrids of less than 88 day RM have been tested, nor does DuPont have access to all seed company data. Consequently, injury arising from the use of Basis on "food grade" or waxy hybrids with a maturity rating of 88 day RM or less is the responsibility of the user.

IMPORTANT

BEFORE USING "BASIS" HERBICIDE, READ AND FOLLOW ALL APPLICABLE DIREC-TIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA-REGISTERED LABEL.

This bulletin contains new or supplemental instructions for use of this product which does not appear on the EPA-registered package label. Follow these instructions carefully. This label must be in the possession of the user at the time of application.

5

APR

Ъ

2 2

μį

WHEN TO APPLY

Apply Basis alone or with its labeled tank mix partners on 1-2 inch foxtail anytime after corn emergence through 4-leaf (2 collar) stage (approximately 1/2" to 6" tall).

Risk of injury increases as the crop grows beyond the 3 collar stage.

HOW TO USE

Apply Basis at 1/3 ounce per acre. Each soluble packet of Basis will treat 4 acres.

ACCEPTED APR 16 1997 Under the Foderal Insocticido. Fungitide. and Redenticide Act. as ansoeded, for the pesticide registered ander UNA Rep. No. 351-571

© 1997 E. I. du Pont de Nemours and Company, DuPont Agricultural Products, Wilmington, Delaware 19898

D-033197