

352-571

4/9/2009

1/16



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

AND

OFFICE OF
PREVENTION, PESTICIDES
TOXIC SUBSTANCES

APR -9 2009

Mr. Tim McPherson
E.I. DuPont de Nemours & Company
DuPont Crop Protection
Stine-Haskell Research Center
P. O. Box 30
Newark, DE 19714-0030

Dear Mr. McPherson:

Subject: DuPont Basis Herbicide (Add Crop Rotation Interval for STS Soybeans)
EPA Registration No. 352-571
Application Dated January 27, 2009

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended is acceptable provided you make the following changes before you release the product for shipment.

1. On page 2, revise the first sentence beneath the Agricultural Use Requirements box to read "DuPont Basis herbicide **must** be used only in accordance with **instructions** on this label or in supplemental DuPont publications."
2. On page 2, revise the second sentence beneath the Agricultural Use Requirements box to read "DuPont will not be responsible for losses or damage resulting from the use of this product in any manner not **specified** by DuPont."
3. On page 6, revise the heading "Additional Recommendations And/Or Recommendations for Specific Weed Problems" to read "Additional **Instructions** And/Or **Instructions** for Specific Weed Problems".
4. On page 7, revise the third bullet in the highlighted box to read "Do not tank mix BASIS with other acetolacate synthase (ALS) inhibiting herbicides (such as "Spirit", "Exceed", "Permit", and "Northstar") unless the mixture is specified on BASIS labels or fact sheets, as severe crop injury may occur."
5. On page 7, revise the second sentence in the first paragraph below the highlighted box to read "Weed Control and crop response with tank mixtures not **specified** on this label are the responsibility of the user and the manufacturer of the tank mix product."

2/16

Page 2

EPA Registration No. 352-571

6. Update the Storage and Disposal section as per PR Notice 2007-4.
7. Update the Limitation of Warranty and Liability based on Agency guidance entitled "Guidance On Warranty Statements" dated October 17, 2006.

Please submit one (1) copy of your final printed labeling incorporating the above changes before you release the product for shipment. Amended labeling supersedes all previously approved ones. A stamped copy of labeling is enclosed for your records.

Sincerely,


Vike KWalley for
James A. Tompkins
Product Manager 25
Herbicide Branch
Registration Division (7505P)



DuPont™ Basis®

herbicide

DRAFT LABEL



“..... A Growing Partnership With Nature”

DUPONT™ BASIS® HIGHLIGHTS

- BASIS® provides selective preplant, preemergence, and postemergence grass and broadleaf weed control in field corn.
- BASIS® preplant and preemergence use rate is 1/3 - 1 oz. product per acre. (When packaged in water-soluble packets, one packet contains 1 1/3 oz. product which treats 1.33 - 4 acres.)
- BASIS® postemergence use rate is 1/3 oz. product per acre. (When packaged in water-soluble packets, one packet contains 1 1/3 oz. product which treats 4 acres.)
- BASIS® may be applied by ground (broadcast or band) or by air.
- For best postemergence results apply in at least 15 gal of water per acre at 20-40 PSI with flat fan nozzles.
- Always include the use of a crop oil concentrate or non-ionic surfactant and an ammonium nitrogen fertilizer in the spray mixture.
- BASIS® may be tank mixed with certain grass and broadleaf herbicides. See **Tank Mix Applications**.
- Applications of BASIS® to grasses or corn under stress may result in less than desirable performance. See **Environmental Conditions**.
- Consult label text for complete instructions. Always read and follow label directions for use.

Table of Contents

	Page
Precautionary Statements	1
Directions For Use	2
General Information	2
Preemergence & Burndown	
Applications in the Fall	2
Rate	2
Adjuvants	2
Tank Mix Applications	2
Application Timing and Weeds Controlled ..	2
Important Precautions	3
Early Preplant, Preemergence, and	
Burndown Applications in the Spring	3
Rate	3
Adjuvants	3
Tank Mixes	3
Weeds Controlled	3
Residual Activity	4
Sequential Applications	4
Application Equipment	4
Mixing Instructions	4
Postemergence Applications	5
Timing to Crop	5
Timing to Weeds	5
Rate	5
Spray Adjuvants	5
Tank Mixes	6
Mixing Instructions	7
Chemigation	7
Cultivation	7
Environmental Conditions and	
Biological Activity	7
Sequential DuPont™ ACCENT®	
Applications	8
Soil Insecticide Interactions	8
Crop Rotation	8
Application Information	8
Broadcast	8
Band	9
Aerial	9
Sprayer Preparation/Cleanup	9
Spray Drift Management	9
Resistance	10
Important Precautions	10
Storage and Disposal	11
Notice of Warranty	12



DuPont™

Basis®

herbicide

ACCEPTED
with COMMENTS
In EPA Letter Dated:

APR -9 2009

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under EPA Reg. No.
352-571

For use in Field Corn

This product is a water-dispersible granule containing
75% active ingredient by weight.

Active Ingredient	By weight
Rimsulfuron	
<i>N</i> [(4,6-dimethoxy-2-pyrimidin-2-yl)aminocarbonyl]-3-(ethylsulfonyl)-2-pyridinesulfonamide	50%
Thifensulfuron methyl	
<i>Methyl</i> 3-[[[(4-methoxy-6-methyl-1,3,5-triazin-2-yl)amino]carbonyl]amino]sulfonyl]-2-thiophenecarboxylate	25%
Inert Ingredients	25%
TOTAL	100%

EPA Reg. No. 352-571

KEEP OUT OF REACH OF CHILDREN

CAUTION

FIRST AID

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.

IF SWALLOWED: No specific intervention is indicated as the compound is not likely to be hazardous by ingestion. However, consult a poison control center or doctor if necessary.

FIRST AID (cont'd)

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-441-3637 for emergency medical treatment information.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Caution! Causes eye irritation. Harmful if absorbed through skin. Avoid contact with skin, eyes or clothing. Avoid breathing dust or spray mist.

PERSONAL PROTECTIVE EQUIPMENT

Some of the materials that are chemical resistant to this product are listed below. If you want more options follow the instructions for category A on an EPA chemical-resistant category selection chart.

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants.
- Chemical Resistant Gloves Category A (such as butyl rubber, natural rubber, neoprene rubber or nitrile rubber), all ≥14 mils.
- Shoes plus socks.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

USERS SHOULD: Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water by cleaning of equipment or disposal of wastes.

5/16

6/16

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 4 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

Coveralls.

Chemical Resistant Gloves Category A (such as butyl rubber, natural rubber, neoprene rubber or nitrile rubber), all ≥ 14 mils.

Shoes plus socks.

DuPont™ BASIS® herbicide should be used only in accordance with recommendations on this label or in supplemental DuPont publications. DuPont will not be responsible for losses or damage resulting from use of this product in any manner not specifically recommended by DuPont.

GENERAL INFORMATION

BASIS® is a selective herbicide for control of certain annual grass and broadleaf weeds in corn.

BASIS® is a water dispersible granule containing 75% active ingredient by weight.

Make only one application of BASIS® per season.

Do not apply to field corn grown for seed, to popcorn, or to sweet corn.

BASIS® may be applied to "food grade" (yellow dent field corn with hard endosperm) and waxy field corn of greater than 88 day Relative Maturity (RM). Not all "food grade" or waxy field corn hybrids of less than 88 day RM have been tested, nor does DuPont have access to all seed company data. Consequently, injury arising from the use of BASIS® on "food grade" or waxy hybrids with a maturity rating of 88 day RM or less is the responsibility of the user. Not all Hi-Lysine, or high oil corn have been fully tested for crop safety, nor does DuPont have

access to all seed company data. Consequently, DuPont is not responsible for any corn injury arising from the use of BASIS® on these specialty corn. Consult with your seed supplier before applying BASIS® to any of the corn types.

PREEMERGENCE AND BURNDOWN APPLICATIONS IN THE FALL

BASIS® herbicide may be applied after harvest but before the ground freezes for fall burndown of certain existing winter annual weeds with limited preemergence control of some early-emerging spring annual grass and broadleaf weeds the following spring, providing a cleaner seedbed for spring planting. Field corn may be planted in the spring following a fall application of BASIS®. A cleaner seedbed is less attractive to cutworm moths and stink bugs. Some weeds such as deadnettle and henbit may be alternate hosts for soybean cyst nematodes. Reducing weed populations may help reduce pressure from these pests in subsequent crops.

The following spring, weed control programs in field corn may include products containing rimsulfuron or nicosulfuron such as DuPont™ ACCENT®, DuPont™ ACCENT GOLD® WDG, DuPont™ BASIS GOLD®, or DuPont™ STEADFAST®. Follow the crop rotation intervals and directions provided on the labels for each of these products.

RATE

Apply 1/3 - 1/2 oz. BASIS® per acre alone or in combination with other herbicides registered for fall application. Do not apply to coarse soils (sand, loamy sandy or sandy loam) with less than 1% organic matter. Make only one application of BASIS® in the fall.

ADJUVANTS

For burndown control of emerged weeds, always add crop oil. See POSTEMERGENCE APPLICATIONS - SPRAY ADJUVANTS.

TANK MIX APPLICATIONS

For expanded weed control, BASIS® may be tank mixed with full or reduced rates of other herbicides labeled for fall application, such as DuPont™ EXPRESS®, simazine, or 2,4-D ester. Refer to the other product's label for rotational crop intervals and other directions for use. When tank mixing, the most restrictive label applies.

APPLICATION TIMING AND WEEDS CONTROLLED

BASIS® may be applied after harvest but before the ground freezes to control the following emerged weeds and to provide limited residual control of early-emerging spring weeds.

Apply 0.33 oz BASIS® for burndown of smaller emerged weeds where little residual activity is desired. Reduced rates of EXPRESS® herbicide (as low as 0.1 oz/acre) may be tank mixed for improved burndown activity on weeds such as dandelion and chickweed. Read and follow all label instructions on rotational intervals, precautions, and restrictions before using this tank mixture. Use 0.5 oz BASIS® for burndown of larger emerged weeds and for added soil residual activity.

7/16

For best control of emerged weeds, apply 0.5 oz DuPont™ BASIS® to the following weeds less than 3" tall and prior to flowering:

- | | |
|-------------------------|-------------------|
| Bittercress | Hemlock, poison |
| Brome, downy | (up to 12" dia.) |
| Bushy wallflower | Henbit |
| Buttercup, smallflower | Parsnip, wild |
| Butterweed | Pennycress, field |
| Catchweed bedstraw | Shepherdspurse |
| Chickweed, common | Speedwell, corn |
| and mouse ear | Wheat, volunteer |
| Dandelion, 6" diameter* | Yellow rocket |
| Deadnettle, purple | |

* for best results add 1 pt 2,4-D LVE (4 lbs formulation)

For best limited residual control of the following weeds, apply 0.5 oz BASIS® to contribute to a clean seedbed at planting:

- | | |
|-------------------------|---------------------|
| Bittercress | Ragweed, common and |
| Dandelion | giant |
| Foxtail, giant, yellow, | Smartweed, PA |
| green | Speedwell, corn |
| Lambsquarter (including | Sunflower |
| triazine resistant) | |

IMPORTANT PRECAUTIONS

Do not apply more than 1 oz of BASIS® in any 12-month period.

Do not apply to coarse soils (sand, loamy sand or sandy loam) with less than 1% organic matter.

Do not apply BASIS® in the fall if DuPont™ MATRIX® or DuPont™ SHADEOUT® has been applied during the preceding season. If fall-applied BASIS® is followed by a spring application of BASIS®, do not exceed 1 oz. of BASIS® per acre total (adding the rate applied in the fall to the rate applied in the spring) and observe the crop rotation intervals listed in the "Crop Rotation" section of this label.

Do not apply "Counter 15G" insecticide the following spring. Follow the most restrictive guidance for insecticides and other precautions provided on this and on the labels of any products applied sequentially.

EARLY PREPLANT, PREEMERGENCE, AND BURNDOWN APPLICATIONS IN THE SPRING

BASIS® herbicide may be applied alone or in a tank mixture with other registered corn herbicides for burndown control of emerged weeds and to provide early season residual weed control in corn. Do not apply to popcorn, sweet corn, or to field corn grown for seed.

RATE

For burndown of emerged weeds and early season residual weed control, broadcast applications of BASIS® may be applied pre-emergence anytime before corn plants emerge. For maximum residual control of grass and broadleaf weeds, tank mix BASIS® with atrazine and make applications preemergence, at planting, or up to 7 days before planting.

Apply BASIS® at a rate of 1/3 to 1 ounce per acre alone or in combination with one or more herbicides registered for use on field corn (See "Tank Mixes" section). Do not apply less than 1/3 ounce of BASIS® per acre. Observe the crop rotation intervals listed in the "Crop Rotation" section of this label.

Preemergence applications of BASIS® work best on soils having an organic content of 1.0% but not more than 3.5%. Apply the higher rate to heavier soil types. Applications on soils with organic matter greater than 3.5% may not provide the desired length of residual activity.

ADJUVANTS

For burndown control of emerged weeds, always add crop oil concentrate. See POSTEMERGENCE APPLICATIONS - SPRAY ADJUVANTS.

TANK MIXES

For broader spectrum of control, early preplant to preemergence applications of BASIS® may be tank mixed with full or reduced rates of one or more herbicides registered for application in corn, such as atrazine, atrazine-containing preemergence herbicides, simazine, dicamba, or 2,4-D LVE. Select a rate appropriate for the weeds present and the desired length of residual control. Refer to the tank mix product labels for additional use directions, use restrictions, rotational crop intervals and precautions. The most restrictive provision on either label shall apply.

Do not tank mix BASIS® with products containing flumetsulam such as "Hornet", "Python" or "Broadstrike"

WEEDS CONTROLLED - BURNDOWN

For best control of emerged weeds, apply BASIS® to the following grasses <2" tall, broadleaves <3" tall and winter annual/biennials <6" tall. For enhanced burndown control tank mix BASIS® with 2,4-D LVE plus atrazine or a preemergence herbicide containing atrazine.

BASIS® applied alone at 1/3 - 1/2 oz. —

Weeds controlled (burndown)

- | | |
|-----------------------------------|----------------------------|
| Barnyardgrass | Lambsquarters |
| Bluegrass, annual | Mustard, wild |
| Brome, downy* | Panicum, fall |
| Butterweed | Parsnip, wild |
| Catchweed bedstraw | Pigweeds |
| Chickweed | Smartweeds, annual |
| Crabgrass, large (< 1/2") | Sunflower |
| Dandelion (6" diameter) | Velvetleaf |
| Fescue, tall* | Wheat, volunteer winter* |
| Foxtail, bristly, Carolina, giant | Woolly cupgrass (up to 1") |
| green and yellow | Wild Oats (<3 leaf) |
| Hemlock, poison | |
| (up to 12" dia.) | |

(*)suppression only

8/16

**DuPont™ BASIS® applied alone at 1/2 - 1 oz. —
Additional weeds controlled**

- | | |
|------------------|------------------|
| Crabgrass, large | Ragweed, common |
| False chamomile | Shepherdspurse |
| Henbit | Volunteer barley |
| Quackgrass | Wild radish |

1/3 - 1 oz BASIS® + 1 pint 2,4-D LVE + 1 pound active ingredient atrazine --Weeds controlled (burndown)

- | | |
|--|------------------------------|
| Barley, volunteer | Mustard, tansy and wild |
| Barnyardgrass | Nightshade |
| Bluegrass, annual | Panicum, fall |
| Brome, downy* | Pennycress, field |
| Burdock | Prickly lettuce |
| Buttercup, smallflower | Quackgrass (4-8" tall) |
| Catchweed bedstraw | Ragweed, common and giant |
| Chickweed | Shepherdspurse |
| Clover, red | Smartweed, annual |
| Crabgrass, large | Speedwell, corn and purslane |
| Dandelion | Thistle, bull and Canada* |
| Deadnettle, purple | Velvetleaf |
| False chamomile | Venice mallow |
| Fescue, tall* | Waterhemp (up to 4") |
| Fleabane, daisy | Wild garlic* |
| Foxtail, bristly, Carolina, giant,
green and yellow | Wild oats (< 3 leaf) |
| Goldenrod | Wild radish |
| Henbit | Wild strawberry |
| Lambsquarters | Winter wheat, volunteer* |
| Marestail | Woolly cupgrass (up to 1") |
| | Yellow rocket |

(* suppression only)

RESIDUAL ACTIVITY

BASIS® applied at 1/3 oz may deliver 1-2 weeks of residual activity depending on time of activating rain, soil organic matter, and soil temperatures.

BASIS® at a rate of 1/2 - 1 ounce per acre applied up to 30 days early preplant until emergence of corn will provide control or suppression of the weeds listed below. An activating rainfall or irrigation (>1/2 inch) 5 to 7 days after application will be required for herbicide activation and best residual activity. For maximum duration of "in-season" residual control, apply BASIS® or BASIS® tank mixes preemergence, at-planting, or up to 7 days before planting.

Residual control or suppression in soils with 1-3.5% organic matter

- | | |
|---------------------------------|--|
| Barnyardgrass | Palmer amaranth* |
| Carpetweed* | Panicum, fall* |
| Cocklebur* | Pigweed, (prostrate, redroot,
smooth) |
| Crabgrass, large* | Purslane, common |
| Foxtail, (giant, green, yellow) | Ragweed, common* |
| Jimsonweed* | Russian thistle, seedling |
| Kochia | Smartweed, PA* |
| Lambsquarters | Velvetleaf* |
| Morningglory, ivyleaf* | |
| Nightshade, (black*, hairy*) | |

(* suppression only)

SEQUENTIAL APPLICATIONS

**DuPont™ ACCENT®, DuPont™ BASIS GOLD®,
DuPont™ ACCENT GOLD® WDG or DuPont™
STEADFAST®**

For season-long control of grass and broadleaf weeds, a sequential application of ACCENT® and/or a broadleaf herbicide may be needed. Make applications of ACCENT® 14 or more days after the BASIS® application. Refer to the ACCENT® label for product use information.

BASIS GOLD®, ACCENT GOLD® WDG or STEADFAST® may be used as a sequential application in a planned postemergence weed control program in corn following application of a reduced rate of BASIS®. Apply 0.3 to 0.62 ounces of BASIS® preemergence to corn and allow at least 3 weeks before making sequential applications of BASIS GOLD®, ACCENT GOLD® WDG or STEADFAST® at the proper timing for weeds and corn. Use the higher rates of BASIS® on heavier soil types or where weed pressures are expected to be heavy.

Allow at least 3 weeks between applications of BASIS® and BASIS GOLD®, ACCENT GOLD® WDG or STEADFAST®. Make sequential applications after the corn has reached the 2-collar stage but before the corn exceeds the maximum application height listed on the respective product labels. Crop injury may result from applications of BASIS GOLD®, ACCENT GOLD® WDG or STEADFAST® made to plants under stress from prior herbicide application. Refer to the BASIS GOLD®, ACCENT GOLD® WDG or STEADFAST® labels for product use information.

Do not apply BASIS GOLD®, ACCENT GOLD® WDG or STEADFAST® to fields treated with more than 0.62 ounce of BASIS®.

APPLICATION EQUIPMENT AND SPRAY VOLUMES

Apply uniformly by ground equipment using a properly calibrated fixed boom sprayer with a minimum of 10 gallons per acre spray volume. Use a minimum of 15 gallons for best performance. Select a spray nozzle, spray volume, and spray pressure configuration to deliver uniform spray coverage onto weeds. Heavy crop residues may reduce burndown control of emerged weeds if residues impede spray coverage. Higher spray volumes and pressures can improve burndown control in heavy crop residue situations.

MIXING INSTRUCTIONS

BASIS® may be mixed with water or pre-slurried in water and added to liquid fertilizer for application. When using liquid fertilizer as the carrier, always pre-slurry BASIS® in water before adding fertilizer solutions. Add the BASIS® slurry to the final complete liquid fertilizer mixture--Do not add BASIS® during the fertilizer mixing process. Always use good agitation while adding the BASIS® slurry to liquid fertilizers and maintain good agitation until sprayed.

When using liquid fertilizer as the carrier, conduct a compatibility test with all components prior to mixing.

9/16

Do not use with spray additives or liquid fertilizer carriers that alter the pH of the spray solution below pH 5.0 or above pH 9.0 as rapid product degradation can occur. Spray solutions of pH 6.0 - 8.0 allow for optimum stability

RATE

Apply BASIS® at 1/3 oz per acre for control of the grasses listed above.

POSTEMERGENCE APPLICATIONS

Do not apply to field corn grown for seed, to popcorn, or to sweet corn. See "APPLICATION INFORMATION" section of this label for use restrictions on other types of corn.

SPRAY ADJUVANTS

Applications of BASIS® must include either a crop oil concentrate or a nonionic surfactant. In addition, an ammonium nitrogen fertilizer must be used. If another herbicide is tank mixed with BASIS®, select adjuvants authorized for use with both products. Products must contain only EPA-exempt ingredients (40 CFR 1001).

TIMING TO CROP STAGE

BASIS® may be applied to field corn in the spike through 4-leaf (2 collar) stage (approximately 1/2" to 6" tall). Do not apply to corn having 3 fully emerged collars or over 6" tall.

Petroleum Crop Oil Concentrate (COC) or Modified Seed Oil (MSO)

- Apply at 1% v/v (1 gal per 100 gal spray solution) or 2% under arid conditions.
- Oil adjuvants must contain at least 80% high quality, petroleum (mineral) or modified vegetable seed oil with at least 15% surfactant emulsifiers.

TIMING TO WEEDS

Apply BASIS® when grasses are young and actively growing, but before they exceed the size listed in the table below.

Applications made to weeds at growth stages greater than those listed below may result in incomplete control. Grass competition due to incomplete control may reduce yields.

Nonionic Surfactant (NIS)

- Apply at 0.25% v/v (1 qt per 100 gal spray solution) or 0.5% under arid conditions.
- Surfactant products must contain at least 60% nonionic surfactant with a hydrophilic/lipophilic balance (HLB) greater than 12.

Grasses	Maximum Height (Inches)
Barnyardgrass	2"
Crabgrass, large	<1/2"
Foxtails	
bristly	2"
giant	2"
green	2"
yellow	2"
Panicum, fall	2"
Shattercane	4"
Woolly cupgrass*	1"
Wild oats	<3 leaf
Broadleaves**	
Lambsquarters, common	3"
Mustard, wild	3"
Pigweed, redroot	3"
Smartweed, annual	3"
Sunflower, wild	3"
Velvetleaf	3"

Ammonium Nitrogen Fertilizer

- Use 2 qt/acre of a high-quality urea ammonium nitrate (UAN), such as 28%N or 32%N, or 2 lb/acre of a spray-grade ammonium sulfate (AMS). Use 4 qt/acre UAN or 4 lb/acre AMS under arid conditions.

- Do not use liquid nitrogen fertilizer as the total carrier solution.

Special Adjuvant Types

- Combination adjuvant products may be used at doses that provide the required amount of NIS, COC, MSO and/or ammonium nitrogen fertilizer. Consult product literature for use rates and restrictions.
- In addition to the adjuvants specified above, other adjuvant types may be used if they provide the same functionality and have been evaluated and approved by DuPont Product Management. Consult separate DuPont technical bulletins for detailed information before using adjuvant types not specified on this label.

* Requires the use of crop oil concentrate.

** Applications to broadleaf weeds in the cotyledon stage may result in less than satisfactory control. Broadleaf weeds that emerge after application may not be controlled by BASIS®. A properly timed cultivation or follow-up application of a broadleaf herbicide may be required.

Do not use with spray additives that alter the pH of the spray solution below pH 5.0 or above pH 9.0 as rapid product degradation can occur. Spray solutions of pH 6.0 - 8.0 allow for optimum stability of BASIS®.

10/16

TANK MIX APPLICATIONS

For Additional Control of Broadleaf Weeds

DuPont™ BASIS® may be tank mixed with the herbicides below for additional control of broadleaf weeds. See the tank mix partner label for weeds controlled, precautions, use restrictions and crop rotation information.

	Rate/A	Adjuvant + Nitrogen Fertilizer
Atrazine 90 DF	1/2 - 1 lb	COC
"Callisto"	2-3 oz	COC*
"Hornet" WDG	1.25 - 2.0 oz.	COC

*Do not use MSO adjuvants with "Callisto" tank mixtures.

Rates listed are for the specific products noted in the table. If other brands or formulations are used, rates of active ingredients should be adjusted to correspond to the products indicated. Formulations of products other than those listed may not have been tested with BASIS®. Check with the manufacturer for information on tank mix compatibility prior to using (see TANK MIX COMPATIBILITY TESTING).

ADDITIONAL RECOMMENDATIONS AND/OR RECOMMENDATIONS FOR SPECIFIC WEED PROBLEMS

For Additional Control of Grass Weeds

BASIS® herbicide may be tank mixed with DuPont™ ACCENT® herbicide for early postemergence control of emerged grasses.

Apply 1/3 oz BASIS® + 1/4 oz ACCENT® per acre alone or in combination with other broadleaf herbicides listed on this label. If a sequential application of ACCENT® is applied, do not apply more than 1 oz ACCENT® per acre per season. Add crop oil concentrate and ammonium nitrogen fertilizer adjuvants.

GRASSES CONTROLLED

	Maximum Weed Height Grasses At Application (inches)
Foxtails, (bristly, giant, green)	3
yellow	2
Barnyardgrass	3
Crabgrass, large**	1
Panicum, fall	3
Johnsongrass, seedling*	8
Shattercane*	6
Sandbur, field	2
Wild proso millet*	2
Wild oats	2

* Controls emerged plants only. Subsequent flushes may require sequential application of ACCENT® herbicide.

** Large crabgrass will be controlled only if size is "mouse ear" (1 leaf emerging) at application.

For Additional Control of Crabgrass and Later Emerging Grasses

BASIS® may be tank mixed with 2/3 to full labeled rates of preemergence grass herbicides labeled for early postemergence application to field corn (such as "Prowl", "Surpass" EC, "Dual II Magnum", and "Outlook") for increased residual activity of later-emerging flushes of grasses such as smooth and large crabgrass. Application must be made before the crabgrass emerges and before other grass weeds on the BASIS® label exceed their labeled sizes.

Tank mixes of BASIS® and preemergence grass herbicides must be broadcast applied postemergence to field corn before the crop exceeds the heights listed on the preemergence grass herbicide label.

When applying BASIS® in tank mixture with preemergence grass herbicides such as "Prowl", and applications are made early postemergence to small grass weeds; use a non-ionic surfactant (NIS) adjuvant in place of crop oil concentrate.

Refer to POSTEMERGENCE APPLICATIONS and the preemergence grass herbicide label for complete postemergence application information, rates, and restrictions.

Tank Mixtures with Dicamba

BASIS® may be tank mixed with 2.0 - 4.0 oz a.i./acre of dicamba (e.g. 4.0 - 8.0 fl oz/acre of "Clarity") for additional control of many broadleaf weeds.

A corn plant's predisposition to develop fused tissue emerging from the whorl (rattail) after the V-11 stage may increase when a product containing dicamba or growth regulator herbicides (i.e. "Clarity", "Marksman") is applied to small corn under early stressful conditions. Be aware of this when applying tank mixes with dicamba to small corn (V-3 stage or smaller) under stressful conditions. See ENVIRONMENTAL CONDITIONS for a description of these stressful conditions.

Tank Mixtures with Insecticides

BASIS® may be tank mixed with pyrethroid or carbamate insecticides such as DuPont™ ASANA® XL or DuPont™ LANNATE® insecticides.

11/16

ADDITIONAL TANK MIX INFORMATION

- Do not tank mix DuPont™ BASIS® with "Basagran" and "Laddok" or severe crop injury may occur.
- Do not tank mix BASIS® with foliar-applied organophosphate insecticides such as "Lorsban", malathion, parathion, etc., as severe crop injury may occur. To avoid crop injury or antagonism, apply these products at least seven days before or 3 days after the application of BASIS®.
- Do not tank mix BASIS® with other acetolactate synthase (ALS) inhibiting herbicides (such as "Spirit", "Exceed", "Permit", and "Northstar") unless the mixture is specifically recommended on BASIS® labels or fact sheets, as severe crop injury may occur.
- Do not exceed labeled application rates. Do not tank mix BASIS® with other products that contain the same active ingredients as BASIS® (rimsulfuron and thifensulfuron) unless the label of either tank mix partner specifies the maximum rate that may be used.

Other than the exceptions noted, and in addition to the tank mix partners and rates indicated in the tables above, BASIS® may be tank mixed or followed with sequential applications of full or reduced rates of other products registered for use in field corn. Weed control and crop response with tank mixtures not specifically recommended in this label are the responsibility of the user and manufacturer of the tank mix product. Read and follow all applicable use directions, precautions, and limitations specified on the respective product labels and fact sheets. BASIS® may be applied in tank mix combinations with other products provided:

- The tank mix product is labeled for the same timing, method of application, adjuvants, and use restrictions as BASIS®.
- The tank mixture is not specifically prohibited on the label of the tank mix product.
- The tank mix combination is compatible as determined by a "jar test" described in the TANK MIX COMPATIBILITY TESTING section below.

TANK MIX COMPATIBILITY TESTING

Perform a jar test prior to tank mixing to ensure compatibility of BASIS® and other pesticides. Use a clear glass quart jar with lid and mix the tank mix ingredients in their relative proportions. Invert the jar containing the mixture several times and observe the mixture for approximately 1/2 hour. If the mixture balls-up, forms flakes, sludges, gels, forms an oily film or layers, or forms precipitates, it is not compatible and the tank mix combination should not be used.

MIXING INSTRUCTIONS

1. Fill the tank 1/4 to 1/3 full of water.
2. While agitating, add the required amount of BASIS®.
3. Continue agitation until the BASIS® is fully dispersed, at least 5 minutes.
4. Once the BASIS® is fully dispersed, maintain agitation and continue filling tank with water. BASIS® should be thoroughly mixed with water before adding any other material.

5. As the tank is filling, add the required spray adjuvants (Crop Oil Concentrate, nonionic surfactant, liquid nitrogen fertilizer, or ammonium sulfate).
6. If the mixture is not continuously agitated, settling will occur. If settling occurs, thoroughly re-agitate before using.
7. Apply BASIS® spray mixture within 24 hours of mixing to avoid product degradation.
8. If BASIS® and a tank mix partner are to be applied in multiple loads, pre-slurry the BASIS® in clean water prior to adding to the tank. This will prevent the tank mix partner from interfering with the dissolution of the BASIS®.

CHEMIGATION

BASIS® should not be applied through any type of irrigation system.

CULTIVATION

For best results, cultivate when BASIS® burndown and residual activity is complete or if new grasses have emerged. The best time for cultivation is usually 10 to 15 days after application.

ENVIRONMENTAL CONDITIONS AND BIOLOGICAL ACTIVITY

BASIS® provides the best results when applied to young, actively growing grasses. Applications made during warm, moist conditions (70 °F or more) with adequate soil moisture both before and after application maximize performance.

The degree and duration of control depend on spray coverage, weed spectrum, weed size, growing conditions both before and after application, soil moisture, precipitation, and adjuvant selection.

Thorough coverage is important for best performance. Excessively cloddy or rough field conditions may reduce coverage and affect performance.

Poor grass control or crop injury may result from applications made to plants under stress from:

- abnormally hot or cold weather
- environmental conditions such as drought, water-saturated soils, hail damage, or frost
- disease, insect or nematode damage
- prior herbicide applications, including carryover from a previous year's herbicide application

If the corn or grass weeds are under stress, delay application of BASIS® until the stress passes and plants begin to grow again. If the corn stage or grass height exceed the sizes indicated on this label, DuPont™ ACCENT® may be used for grass control. Consult the ACCENT® labels for timing information.

Applications made during or immediately following periods of large day/night temperature fluctuations may decrease control and increase the potential for crop injury. Applications should be made when minimum nighttime temperatures are above 40 °F and the maximum daytime temperatures are below 92 °F.

12/16

Applications of DuPont™ BASIS® to dry, dusty fields may reduce grass control in the wheel tracks.

BASIS® rapidly inhibits the growth of susceptible grasses, reducing competition within as little as 6 hours after application. Susceptible grasses are controlled in 7-14 days. BASIS® is rainfast in 4 hours.

SEQUENTIAL DUPONT™ ACCENT® APPLICATIONS

To control grasses under adverse environmental conditions, a sequential application of ACCENT® herbicide may be necessary. An application of ACCENT® may be made 14 or more days after BASIS® applications. Refer to ACCENT® labels for grass species controlled, proper size of weeds, rates and other information.

SOIL INSECTICIDE INTERACTIONS

Before using BASIS®, ensure that it is compatible with any insecticides applied to the corn crop.

Soil Insecticides	Application Method	Soil O.M.	BASIS® Use Precautions
“Counter 15G”	All	All	Do not use.
“Counter 20CR”	In furrow at planting	All	Do not use.
	Over the row at cultivation	All	Do not use.
	T-band or Surface band	All	May cause unacceptable injury.
“Lorsban”	All labeled methods	All	May cause unacceptable injury.
“Thimet”	All labeled methods	All	May cause unacceptable injury.
“Fortress”	All	All	No use precautions.
“Force”	All	All	No use precautions
“Aztec”	All	All	No use precautions

CROP ROTATION

The following rotational intervals should be observed when using BASIS® at 1/3 oz. :

Crop	Interval in Months
Alfalfa	10
Cereals, Spring (wheat, oats, barley)	8
Cereals, Winter (wheat, barley, rye)	4
Corn (field)	Anytime
Corn (pop or sweet)	10
Cotton	10
Grain sorghum	10
Potatoes	Anytime
Snap beans, dry beans, peas	8
Soybeans*	1/2 (15 days)
Sugarbeets	10
Sunflowers	10
Sweet potatoes	10**
Tomatoes	1
Crops not listed	18

*Crop rotation interval listed for soybeans are intended to support the ability to re-plant soybeans following a failed corn crop where BASIS® had been previously applied. BASIS® should not be used as a part of a planned soybean herbicide program.

**On soils with pH <6.5.

The following rotational intervals should be observed when using BASIS® at between 1/3 and 1 oz. :

Crop	Interval in Months
Cereals, Spring (wheat, oats, barley)	9
Cereals, Winter (wheat, barley, rye)	4
Corn (field)	Anytime
Corn (pop or sweet)	10
Potatoes	Anytime
Snap beans, dry beans	10
Soybeans	10
STS soybeans*	4
Sunflowers	10
Tomatoes	1
Crops not listed	18

* Rotational Interval for STS (sulfonyurea tolerant soybean) is 1 month if using 0.50 oz per acre or less

**APPLICATION INFORMATION
BROADCAST GROUND APPLICATION**

Use a minimum of 15 gal of water per acre (GPA) applied at 20-40 PSI for best performance. Ensure that equipment is set up to avoid applying an excessive rate directly over the rows and into the corn plant whorl.

- Use 50 mesh screens with all applications.
- Use flat fan nozzles only. Do not use hollow cone, raindrop, whirl chamber or controlled droplet applicator (CDA) nozzles, or air-assisted sprayers. Unacceptable crop injury, excessive spray drift, or poor weed control may result.
- Overlaps, or starting, stopping, slowing, and turning while spraying may result in crop injury.

- For proper spray coverage, adjust the boom and nozzle height to manufacturers' specifications.

BAND APPLICATION

- For band applications, use proportionately less spray mixture.
- To avoid crop injury, carefully calibrate the band applicator so as not to exceed the labeled rate.
- Carefully follow the manufacturer's instructions for flat fan nozzle orientation, distance of nozzles from the crop and weeds, spray volumes, calibration and spray pressure.

AERIAL APPLICATION (SEE ALSO SPRAY DRIFT MANAGEMENT)

- Use nozzle types and arrangements that will provide optimum spray distribution and maximum coverage at a minimum of 5 GPA.
- Do not apply during a temperature inversion, when winds are gusty, or when conditions favor poor coverage and/or off-target spray movement.
- Do not apply by air in the state of New York.

SPRAYER PREPARATION/CLEANUP

It is important that spray equipment is clean and free of previous pesticide deposits before using DuPont™ BASIS®. Follow the cleanup procedures specified on the label of the product previously sprayed including directions for rinsate disposal. If no cleanup procedure is provided, use the procedure that follows. Immediately following applications of BASIS®, thoroughly clean all mixing and spray equipment to avoid subsequent crop injury.

Note: When applying multiple loads of BASIS®, do not allow empty sprayer or mixing equipment to stand overnight. Partially fill the empty equipment with fresh water at the end of each day of spraying, flush the boom, hoses and other equipment, and allow to sit overnight.

Cleanup Procedure

1. Drain the tank and thoroughly hose down the interior surfaces. Flush the tank, hoses, and boom with clean water for a minimum of 5 min.
2. Partially fill the tank with clean water and add one gal of household ammonia* (containing 3% active) for every 100 gal of water. Finish filling the tank with water, then flush the cleaning solution through the hoses, boom, and nozzles. Add more water to completely fill the tank and allow to agitate/recirculate for at least 15 min. Again, flush the hoses, boom, and nozzles with the cleaning solution, then drain the tank.
3. Repeat Step 2.
4. Remove the nozzles and screens and clean separately in a bucket containing the cleaning agent and water.
5. Thoroughly rinse the tank with clean water for a minimum of 5 min, flushing the water through the hoses and boom.
6. Rinsate may be applied to areas planted to field corn. Do not exceed the labeled concentration of BASIS®. Read and follow all label directions for BASIS®.

* Equivalent amounts of an alternate strength ammonia solution or a tank cleaner recommended in the DuPont bulletin "Sulfonylurea Herbicides, A Guide to Equipment Cleanout," may be used.

SPRAY DRIFT MANAGEMENT

The interaction of many equipment and weather-related factors determines the potential for spray drift. The applicator is responsible for considering all these factors when making application decisions.

AVOIDING SPRAY DRIFT IS THE RESPONSIBILITY OF THE APPLICATOR.

Importance of Droplet Size

The most effective way to reduce drift potential is to apply large droplets (>150 - 200 microns). The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. The presence of sensitive species nearby, the environmental conditions, and pest pressure may affect how an applicator balances drift control and coverage. **APPLYING LARGER DROPLETS REDUCES DRIFT POTENTIAL, BUT WILL NOT PREVENT DRIFT IF APPLICATIONS ARE MADE IMPROPERLY OR UNDER UNFAVORABLE ENVIRONMENTAL CONDITIONS!** See **Wind, Temperature and Humidity, and Temperature Inversions** sections of this label.

Controlling Droplet Size - General Techniques

- **Volume** - Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets.
- **Pressure** - Use the lower spray pressures recommended for the nozzle. Higher pressure reduces droplet size and does not improve canopy penetration. **WHEN HIGHER FLOW RATES ARE NEEDED, USE A HIGHER-CAPACITY NOZZLE INSTEAD OF INCREASING PRESSURE.**
- **Nozzle Type** - Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles.

Controlling Droplet Size - Aircraft

- **Number of Nozzles** - Use the minimum number of nozzles with the highest flow rate that provide uniform coverage.
- **Nozzle Orientation** - Orienting nozzles so that the spray is emitted backwards, parallel to the airstream will produce larger droplets than other orientations.
- **Nozzle Type** - Solid stream nozzles (such as disc and core with swirl plate removed) oriented straight back produce larger droplets than other nozzle types.
- **Boom Length** - The boom length should not exceed 3/4 of the wing or rotor length - longer booms increase drift potential.
- **Application Height** - Application more than 10 ft above the canopy increases the potential for spray drift.

Boom Height

Setting the boom at the lowest labeled height (if specified) which provides uniform coverage reduces the exposure of droplets to evaporation and wind. For ground equipment, the boom should remain level with the crop and have minimal bounce.

13/116

14/16

Wind

Drift potential increases at wind speeds of less than 3 mph (due to inversion potential) or more than 10 mph. However, many factors, including droplet size and equipment type determine drift potential at any given wind speed. **AVOID GUSTY OR WINDLESS CONDITIONS.**

Note: Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they affect spray drift.

Temperature and Humidity

When making applications in hot and dry conditions, set up equipment to produce larger droplets to reduce effects of evaporation.

Temperature Inversions

Drift potential is high during a temperature inversion. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain close to the ground and move laterally in a concentrated cloud. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

Shielded Sprayers

Shielding the boom or individual nozzles can reduce the effects of wind. However, it is the responsibility of the applicator to verify that the shields are preventing drift and not interfering with uniform deposition of the product.

RESISTANCE

When herbicides with the same mode of action are used repeatedly over several years to control the same weed species in the same field, naturally-occurring resistant weed biotypes may survive a correctly applied herbicide treatment, propagate, and become dominant in that field. These resistant weed biotypes may not be adequately controlled. Cultural practices such as tillage, preventing weed escapes from going to seed, and using herbicides with different modes of action within and between crop seasons can aid in delaying the proliferation and possible dominance of herbicide resistant weed biotypes.

IMPORTANT PRECAUTIONS

Injury or loss of desirable trees or vegetation may result from the failure to observe the following:

- Do not apply DuPont™ BASIS® or drain or flush application equipment on or near desirable trees or other plants, on areas where their roots may extend, or in locations where the chemical may be washed or moved into contact with their roots.
- Do not use on lawns, walks, driveways, tennis courts, or in similar areas.
- Prevent drift of spray onto desirable plants (See SPRAY DRIFT).
- Do not contaminate any body of water with BASIS®.
- Thoroughly clean application equipment before and after use. See SPRAYER PREPARATION/CLEANUP section of this label for instructions.
- Do not graze or feed forage, grain, or stover from treated areas to livestock within 30 days of BASIS® application.

15/
16

STORAGE AND DISPOSAL

Storage: Store product in original container only. Do not contaminate water, other pesticides, fertilizer, food or feed in storage. Store in a cool, dry place.

Product Disposal: Do not contaminate water, food, or feed by disposal. Waste resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: For Plastic Containers: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke. **For Fiber Sacks:** Completely empty fiber sack by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into manufacturing or application equipment. Then dispose of sack in a sanitary landfill or by incineration if allowed by State and local authorities. **For Fiber Drums With Liners:** Completely empty liner by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application equipment. Then dispose of liner in a sanitary landfill or by incineration if allowed by State and local authorities. If drum is contaminated and cannot be reused, dispose of in the same manner. **For Bags Containing Water Soluble Packets:** Do not reuse the outer box or the resealable plastic bag. When all water-soluble packets are used, the outer packaging should be clean and may be disposed of in a sanitary landfill or by incineration, or if allowed by State and local authorities, by open burning. If burned, stay out of smoke. If the resealable plastic bag contacts the formulated product in any way, the bag must be triple-rinsed with clean water. Add the rinsate to the spray tank and dispose of the outer wrap as described above. **For Metal Containers (non aerosol):** Triple rinse (or equivalent) the container. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities. **For Paper and Plastic Bags:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

"Dual II Magnum", "Exceed", "Force", "Northstar", "Spirit", are trademarks of Syngenta Crop Protection Inc.

"Broadstrike", "Hornet", "Lorsban", "Python", "Surpass" are trademarks of Dow AgroSciences

"Permit" is a trademark of Monsanto Company

"Basagran", "Clarity", "Counter", "Frontier", "Marksman", "Outlook", "Prowl", "Thimet" are trademarks of BASF Corporation

"Laddok" is a trademark of Sipcarn Agro USA

"Fortress" is a trademark of Amvac Chemical Corp.

D - 1430 012609

NOTICE TO BUYER: Purchase of this material does not confer any rights under patents of countries outside of the United States.

The DuPont Oval Logo, DuPont™, BASIS®, ACCENT®, ACCENT GOLD®, BASIS GOLD®, EXPRESS®, MATRIX®, SHADEOUT®, STEADFAST®, ASANA® and LANNATE® are trademarks or registered trademarks of E.I. duPont de Nemours and Company

16/
16

**LIMITATION OF
WARRANTY AND LIABILITY**

NOTICE: Read This Limitation of Warranty and Liability Before Buying or Using This Product. If the Terms Are Not Acceptable, Return the Product at Once, Unopened, and the Purchase Price Will Be Refunded.

It is impossible to eliminate all risks associated with the use of this product. Such risks arise from weather conditions, soil factors, off target movement, unconventional farming techniques, presence of other materials, the manner of use or application, or other unknown factors, all of which are beyond the control of DuPont. These risks can cause: ineffectiveness of the product; crop injury, or; injury to non-target crops or plants.

DuPont does not agree to be an insurer of these risks. **WHEN YOU BUY OR USE THIS PRODUCT, YOU AGREE TO ACCEPT THESE RISKS.**

DuPont warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for the purpose stated in the Directions for Use, subject to the inherent risks described above, when used in accordance with the Directions for Use under normal conditions.

DUPONT MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR OF MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

IN NO EVENT SHALL DUPONT OR SELLER BE LIABLE FOR ANY INCIDENTAL, CONSEQUENTIAL OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. BUYER'S OR USER'S BARGAINED-FOR EXPECTATION IS CROP PROTECTION. THE EXCLUSIVE REMEDY OF THE USER OR BUYER AND THE EXCLUSIVE LIABILITY OF DUPONT OR SELLER, FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY OR CONTRACT, NEGLIGENCE, TORT OR STRICT LIABILITY), WHETHER FROM FAILURE TO PERFORM OR INJURY TO CROPS OR OTHER PLANTS, AND RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT, OR AT THE ELECTION OF DUPONT OR SELLER, THE REPLACEMENT OF THE PRODUCT.

DuPont or its Ag Retailer must have prompt notice of any claim so that an immediate inspection of buyer's or user's growing crops can be made. Buyer and all users shall promptly notify DuPont or a DuPont Ag Retailer of any claims, whether based on contract, negligence, strict liability, other tort or otherwise or be barred from any remedy.

This Limitation of Warranty and Liability may not be amended by any oral or written agreement.

For product information call: 1-888-6-DUPONT

Internet address: <http://cropprotection.dupont.com/>

© 1997-2009 E. I. du Pont de Nemours and Company, Wilmington, Delaware 19898.

All rights reserved.