

PLEASE NOTE

This image contains more than one label approved for this product on this date.

352-572

4-25-2001

1/14



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY

WASHINGTON, D.C. 20460

APR 25 2001

OFFICE OF
PREVENTION, PESTICIDES
AND TOXIC SUBSTANCES

Mr. Tony E. Catka
DuPont Crop Protection
Stine-Haskell Research Center
1090 Elkton Road
Newark, DE 19714-0030

Dear Mr. Catka:

Subject: DuPont DPX79406 75 DF Herbicide (Revise and Update Label)
EPA Registration No. 352-572
Your Application Dated January 30, 2001

The labeling submitted above in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended is acceptable provided you make the following changes.

1. The heading "Statement of Practical Treatment" must be changed to "First Aid".
2. Refer to PR Notice 2001-1 for preferred First Aid statements.
3. Under Personal Protection Equipment add the following statement.

--Some materials that are chemical resistant to this product are listed below. If you want more options follow the instructions for Category A on the EPA chemical resistance category selection chart.--

4. Under Personal Protective Equipment and in the Agricultural Requirements Use box, "waterproof gloves" must be replaced with a statement similar to "Chemical Resistant Gloves Category A (such as butyl rubber, natural rubber, neoprene rubber, or nitrile rubber) \geq 14 mls.

Please submit three (3) copies of your final printed labeling bearing the above changes before you release the product for shipment. Amended labeling will supercede all previously accepted ones. A stamped copy of labeling is enclosed for your records.

2/14

-2-

For future submissions, labeling requiring revisions will not be stamped "accepted".
Labeling requiring revisions will be rejected pending submission of corrected labeling.

Sincerely,


Debbie K. Walters for
Dan Rosenblatt
Acting Product Manager 25
Herbicide Branch
Registration Division (7505C)



DPX-79406 75DF

herbicide

DRAFT LABEL



“..... A Growing Partnership With Nature”

DUPONT DPX-79406 75DF HIGHLIGHTS

- DPX-79406 75DF provides selective postemergence grass and broadleaf weed control in field corn.
- Apply at a rate of 1/2 ounce product per acre.
- Include an adjuvant as recommended in this label. The use of nitrogen fertilizer is required in addition to the use of a crop oil concentrate or nonionic surfactant. See SPRAY ADJUVANT.
- DPX-79406 75DF may be applied by ground (broadcast or band) or by air.
- For ground application, apply in a minimum of 15 gal of water at 20-40 PSI using flat fan nozzles for best results. For aerial application, use nozzle types and arrangements that provide optimum spray distribution and maximum coverage at a minimum of 5 GPA.
- Recommended tank mixes are specified on this label. See TANK MIX APPLICATIONS.
- Apply to actively growing grass and broadleaf weeds at the recommended sizes. See RATE.
- Applications made to grasses or corn under stress may affect the performance or in some cases reduce crop tolerance. See ENVIRONMENTAL CONDITIONS.
- Consult label text for complete instructions. Always read and follow label directions for use.

TABLE OF CONTENTS

Directions for Use	2
General Information	2
When to apply - postemergence	2
Timing to crop	2
Timing to weeds	2
Rate	3
Weeds controlled	3
Spray adjuvants	3
Mixing instructions	3
When to apply - sequential applications	4
Tank mix applications	4
Broadleaf weeds	4
Crabgrass	5
Additional information	5
Tank mix compatibility testing	6
Sequential ACCENT® applications	6
Cultivation	6
Environmental conditions and biological activity	6
Soil insecticide interactions	6
Crop rotation	7
Application information	7
Ground application	7
Aerial application	7
Sprayer preparation and cleanout	7
Spray drift management	8
Weed Resistance	8
Integrated Pest Management	9
Important precautions	9
Storage and Disposal	9
Warranty	10



DPX-79406 75DF

herbicide

For use in Field Corn

This product is a water-dispersible granule containing 75% active ingredient by weight.

<u>Active Ingredients</u>	<u>By Weight</u>
Nicosulfuron 2-[[[(4,6-dimethoxypyrimidin-2-yl)aminocarbonyl]aminosulfonyl]-N,N-dimethyl-3-pyridinecarboxamide	37.5%
Rimsulfuron N((4,6-dimethoxypyrimidin-2-yl) aminocarbonyl)-3-(ethylsulfonyl)-2-pyridinesulfonamide	37.5%
<u>Inert Ingredients</u>	25%
TOTAL	100%

EPA Reg. No. 352-572

ACCEPTED
APR 25 2001

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No 352-572

KEEP OUT OF REACH OF CHILDREN

CAUTION

STATEMENT OF PRACTICAL TREATMENT

If in eyes: Flush eyes with plenty of water. Call a physician if irritation persists.

If on skin: Wash with plenty of soap and water.

If swallowed: No specific intervention is indicated as the compound is not likely to be hazardous by ingestion. However consult a physician if necessary.

For medical emergencies involving this product, call toll-free 1-800-441-3637.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Caution! Causes moderate eye irritation. Harmful if absorbed through skin. Avoid contact with skin, eyes or clothing.

PERSONAL PROTECTIVE EQUIPMENT

Applicators and other handlers must wear:

Long-sleeved shirt and long pants.

Waterproof gloves.

Shoes plus socks.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

USERS SHOULD: Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment rinsewater. Do not apply where/when conditions could favor runoff. Do not apply if a severe storm is expected within 24 hours.

5/14

6/14

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 4 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls.
- Waterproof gloves.
- Shoes plus socks.

DuPont DPX-79406 75DF Herbicide should be used only in accordance with recommendations on this label or in supplemental DuPont publications. DuPont will not be responsible for losses or damage resulting from use of this product in any manner not specifically recommended by DuPont.

GENERAL INFORMATION

DPX-79406 75DF Herbicide is a water-dispersible granule used at the rate of 1/2 ounce per acre for selective postemergence grass and broadleaf weed control in field corn.

WHEN TO APPLY - POSTEMERGENCE

TIMING TO CROP STAGE

Apply DPX-79406 75DF to corn that is up to 12 inches tall. Do not apply to corn taller than 12 inches or exhibiting 6 collars, whichever is more restrictive.

Apply DPX-79406 75DF to field corn hybrids with a relative maturity (RM) rating of 88 days or more, including "food grade" (yellow dent, hard endosperm), waxy and DuPont Optimum High-Oil corn. Not all field corn hybrids of less than 88 days RM, not all white corn hybrids nor Hi-Lysine hybrids have been tested for crop safety, nor does DuPont have access to all seed company data. Consequently, injury arising from the use of DPX-79406 75DF on these types of corn is the responsibility of the user. Consult with your seed supplier before applying DPX-79406 75DF to any of these corn types. Consult with your local DuPont representative for any additional supplemental labeling information.

Do not apply to field corn grown for seed, to popcorn or to sweet corn.

TIMING TO WEEDS

Apply DPX-79406 75DF when grasses are young and actively growing, but before they exceed the sizes listed on this label.

- Applications made to weeds at growth stages greater than those listed below may result in incomplete control. Grass competition due to incomplete control may reduce yields.
- Adequate soil moisture is required for optimum activity. Rainfall within 5 to 7 days after application will enhance DPX-79406 75DF residual activity. If an activating rainfall or sprinkler irrigation (>0.5 inch) is not received within 5-7 days after application, follow with a cultivation or with a sequential application of DuPont ACCENT® Herbicide, if needed. See CULTIVATION or SEQUENTIAL ACCENT® APPLICATIONS.

RATE

- Apply DPX-79406 75DF at a rate of 1/2 ounce per acre for season-long control of grass and broadleaf weeds listed below.

WEEDS CONTROLLED

Grasses	Height (Inches) at Application
Barnyardgrass	3"
Cereals, volunteer	2"
Crabgrass, large	1"
Cupgrass, woolly	1"
Foxtails	
bristly	3"
giant	3"
green	3"
yellow	3"
Johnsongrass, seedling	8"
Panicum, fall	3"
Millet, wild proso	2"
Quackgrass	8"
Ryegrass, Italian	4"
Sandbur, field	2"
Shattercane	6"
Signalgrass, broadleaf	2"
Oats, wild	2"

*Size in inches or diameter, whichever is more restrictive. Retreatment of later emerging weeds may be required.

Broadleaf Weeds	Height (Inches) at Application
Control:	
Amaranth, powell	4"
Jimsonweed	4"
Morningglory, annual	4"
Mustard, wild	4"
Pigweed, redroot	4"
Smartweed, PA	3"
Sunflower, common	4"
Suppression:	
Cocklebur, common	4"
Ladysthumb	3"
Lambsquarters, common	2"
Hemp dogbane	4"
Nutsedge, yellow	2"
Thistle, Canada	4"
Velvetleaf	4"
Waterhemp, common	2"
Waterhemp, tall	2"

Do not use less than 1/2 ounce per acre of DuPont DPX-79406 75DF.

- As weeds mature, their sensitivity to DPX-79406 75DF decreases. Grassy weeds growing under stress due to drought or other environmental factors may become mature (more than 3 tillers) before they reach the size listed, in which case their susceptibility to DPX-79406 75DF may be reduced.

SPRAY ADJUVANTS

Applications of DPX-79406 75DF must include either a crop oil concentrate or a nonionic surfactant. In addition, an ammonium nitrogen fertilizer must be used. If another herbicide is tank mixed with DPX-79406 75DF to increase the broadleaf weed spectrum, select adjuvants based on the adjuvant limitations of the other herbicide. Crop oil concentrate plus ammonium nitrogen fertilizer is the preferred adjuvant system for DPX-79406 75DF. Consult local DuPont fact sheets, technical bulletins, and service policies prior to using other adjuvant systems.

Crop Oil Concentrate (COC)

- Apply at rate (concentration) of 1% v/v (1 gal per 100 gal spray solution).
- Under arid conditions such as those found in the High Plains, apply at rate (concentration) of 2% v/v (2 gal/100 gal spray solution).
- The crop oil concentrate must be high quality, petroleum or vegetable seed oil-based product (methylated seed oil is considered a vegetable seed-based oil). Petroleum based oils must contain at least 14% emulsifiers/surfactants.

Nonionic Surfactant (NIS)

- Apply at a rate (concentration) of 0.25–0.5% v/v (1–2 qt per 100 gal spray solution). Use the higher rate in drought conditions to enhance control.
- At least 50% of the surfactant product must be active nonionic surfactant.
- Avoid products that do not accurately define their ingredients. Products must contain only EPA-exempt ingredients (40 CFR 1001).

Ammonium Nitrogen Fertilizer

An ammonium nitrogen fertilizer is required in addition to a crop oil concentrate or nonionic surfactant.

- Use a high-quality liquid nitrogen fertilizer such as 28–0–0 at a rate of 2 qt per acre or 10–34–0 at 1 qt per acre.
- A high-quality spray-grade ammonium sulfate (21–0–0) may be applied at a rate of 2 lb per acre in place of the liquid nitrogen fertilizer.
- Under arid conditions such as those found in the High Plains, use 28–0–0 at 4 qt per acre, 10–34–0 at 2 qt per acre or spray-grade ammonium sulfate (21–0–0) at 4 lb per acre.
- Do not use liquid nitrogen fertilizer as the total carrier solution.

MIXING INSTRUCTIONS

1. Fill the tank 1/4 to 1/3 full of water.
2. While agitating, add the required amount of DPX-79406 75DF.
3. Continue agitation until the DPX-79406 75DF is fully dispersed, at least 5 minutes.
4. Once the DPX-79406 75DF is fully dispersed, maintain agitation and continue filling tank with water. Thoroughly mix DPX-79406 75DF with water before adding any other material.

8/14

5. As the tank is filling, add the required spray adjuvants (crop oil concentrate, nonionic surfactant, or ammonium nitrogen fertilizer).
6. If the mixture is not continuously agitated, settling will occur. If settling occurs, thoroughly re-agitate before using.
7. Apply DPX-79406 75DF spray mixture within 24 hours of mixing to avoid product degradation.
8. Apply DPX-79406 75DF + "Hornet" tank mixture within 4 hours of mixing to avoid product degradation. If spray mixture will not be applied within 4 hours and either 28-0-0 or ammonium sulfate was used as the nitrogen source, either add 1 qt/acre of 10-34-0 or 11-37-0, or a commercial tank buffer that will raise the tank pH to 7.0. If 10-34-0 or 11-37-0 was used as the liquid nitrogen source, do not add additional fertilizer or buffer solution to the spray tank.
9. If DPX-79406 75DF and a tank mix partner are to be applied in multiple loads, pre-slurry the DPX-79406 75DF in clean water prior to adding to the tank. This will prevent the tank mix partner from interfering with the dissolution of the DPX-79406 75DF.

WHEN TO APPLY - SEQUENTIAL APPLICATIONS FOLLOWING REDUCED RATES OF PREEMERGENCE HERBICIDES

DuPont DPX-79406 75DF may be used as a sequential application in a planned postemergence weed control program in corn following a reduced rate of a preemergence herbicide.

Apply a reduced rate of a preemergence grass herbicide prior to corn emergence and then follow with a postemergence application of DPX-79406 75DF. Apply products such as DuPont LEADOFF®, "Balance PRO", "Axiom", "Dual II Magnum", "Surpass", "Frontier" and "Harness Xtra" at as low as 1/4 to 1/2 of the full labeled use rate and follow with a sequential postemergence application of DPX-79406 75DF. Refer to WHEN TO APPLY - POSTEMERGENCE and ENVIRONMENTAL CONDITIONS AND BIOLOGICAL ACTIVITY sections for complete application information and precautions. Refer to the preemergence grass herbicide label for use restrictions, application information, rotational crop guidelines, and cautionary statements prior to applying DPX-79406 75DF.

Do not apply DPX-79406 75DF to corn that exhibits herbicide injury from previous applications made to the current or preceding crop.

TANK MIX APPLICATIONS

Refer to the table below for weeds controlled using preferred tank mixtures.

For Additional Control of Broadleaf Weeds

DPX-79406 75DF may be tank mixed with the herbicides below for additional control of broadleaf weeds. See the tank mix partner label for weeds controlled, precautions, use restrictions and crop rotation information.

Crop oil concentrate must be used in the tank mixtures specified below. The use of nonionic surfactant is permitted in place of crop oil concentrate for tank mixtures with preemergence grass herbicides such as "Prowl" where applications are made early postemergence to small grass weeds (see FOR ADDITIONAL CONTROL OF CRABGRASS AND LATER EMERGING GRASS WEEDS). See SPRAY ADJUVANTS for adjuvant rate recommendations.

	Rate/A
atrazine 90DF	1/4 - 1 lb
dicamba (i.e. "Clarity" - 4lb/gal dicamba)	1/8 - 1/4 pt
dicamba + atrazine (i.e. "Marksman" - 1.1 lb/gal dicamba)	1/2 - 1 pt
"Hornet"	1.6 - 2.4 oz
"Hornet WDG"	2.0 - 3.0 oz

Rates listed are for the specific products noted in the table. If other brands or formulations are used, rates of active ingredients should be adjusted to correspond to the products indicated. Formulations of products other than those listed may not have been tested with DPX-79406 75DF. Check with the manufacturer for information on tank mix compatibility prior to using (see TANK MIX COMPATIBILITY TESTING).

The table below indicates weeds controlled using preferred tank mixtures.

Broadleaf Weeds	DPX-79406 75DF Alone	+ 4 oz. "Clarity"	+ 1 pt "Marksman"	+ 1.6 oz. "Hornet"	+ 1/2 lb atrazine
Amaranth, powell	4"	4"	4"	4"	4"
Cocklebur, common	4**	3"	3"	3"	3"
Hemp, dogbane†	4**	4**	4**	4**	4**
Jimsonweed	4"	4"	4"	4"	4"
Kochia		3**	3**		
Ladysthumb	3**	3**	3**	3"	3**
Lambsquarters, common	2**	3"	3"	2**	3"
Mallow, venice				3"	
Morningglory, annual	4"	4"	4"	4"	4"
Mustard, wild	4"	4"	4"	4"	4"
Nightshade, eastern black		2"	3"	2**	2"
Nutsedge, yellow†	2**	2**	2**	2**	2**
Pigweed, redroot	4"	4"	4"	4"	4"
Ragweed, common		3"	3"	3"	3"
Ragweed, giant		3**	3**	3"	3**
Smartweed, PA	3"	4"	4"	4"	4"
Sunflower, common	4"	4"	4"	4"	6"
Thistle, Canada†	4**	4**	4**	4**	4**
Velvetleaf	4**	3"	3"	3"	2"
Waterhemp, common, tall	2**	2"	3"	2**	2"

*Suppression
†Above ground portions only.

All tank mixtures require the addition of crop oil concentrate and ammonium nitrogen fertilizer as noted in SPRAY ADJUVANTS.

9/14

For Additional Control of Grass Weeds

DuPont DPX-79406 75DF may be tankmixed with a reduced rate of DuPont ACCENT® herbicide for enhanced control of many grass weeds listed in WEEDS CONTROLLED. Mix DPX-79406 75DF with 1/4 - 1/3 oz of ACCENT® for control of the following weeds:

WEED	MAXIMUM SIZE CONTROLLED	
	DPX-79406 75DF Alone	DPX-79406 75DF PLUS 1/4 - 1/3 oz ACCENT®
Cupgrass, woolly	<2"	4"
Foxtail: bristly, giant, green, yellow	3"	4"
Johnsongrass, seedling	8"	12"
Panicum, fall	3"	4"
Proso millet, wild	2"	4"
Sandbur, field	2"	2"

For Additional Control of Crabgrass and Later Emerging Grasses

Tank mix DPX-79406 75DF with preemergence grass herbicides labeled for early postemergence application to field corn (such as DuPont LEADOFF®, "Prowl", "Surpass EC", "Dual II Magnum", "Harness", and "Frontier") for increased residual activity of later-emerging flushes of grasses such as smooth and large crabgrass. Application must be made before the crabgrass emerges and before other grass weeds on the DPX-79406 75DF label exceed their labeled sizes.

Tank mixes of DPX-79406 75DF and preemergence grass herbicides must be broadcast applied postemergence to field corn before the crop exceeds the heights listed on the preemergence grass herbicide label. Refer to WHEN TO APPLY-POSTEMERGENCE and the preemergence grass herbicide label for complete postemergence application information, rates, and restrictions.

If preemergence herbicides containing atrazine are mixed with DPX-79406 75DF, use less than a total of 1.0 lb atrazine in the tank mixture.

Add the required amount of preemergence grass herbicide to DPX-79406 75DF. Use the lower end of the rate ranges for soils with less than 3% organic matter. Use the mid-upper end of the rate ranges for soils with >3% organic matter. One of the broadleaf products listed on this label may also be tank mixed for additional broadleaf weed control.

Preemergence herbicides rely on activation from either rainfall or overhead sprinkler irrigation to move the herbicide into the grass weed germination zone and provide control. The amount of precipitation or irrigation required is dependent upon existing soil moisture, soil type and organic matter content. Normally, 1/2-3/4 inch is sufficient.

Additional Information

DuPont DPX-79406 75DF may be tank mixed with DuPont ASANA® XL or DuPont LANNATE® Insecticides.

Do not tank mix DPX-79406 75DF with DuPont BEADEX® Herbicide, "Basagran" and "Laddok" or severe crop injury may occur.

Do not tank mix DPX-79406 75DF with 2,4-D-containing products as severe grass control antagonism may occur.

Do not tank mix DPX-79406 75DF with foliar-applied organophosphate insecticides such as "Lorsban", malathion, parathion, etc., as severe crop injury may occur.

To avoid crop injury or antagonism, apply these products at least seven days before or 3 days after the application of DPX-79406 75DF.

Do not tank mix DPX-79406 75DF with other acetolactate synthase (ALS) inhibiting herbicides (such as "Spirit", "Exceed", "Permit", and "Northstar") unless the mixture is specifically recommended on DPX-79406 75DF labels or fact sheets, as severe crop injury may occur.

• A corn plant's predisposition to develop fused tissue emerging from the whorl (rattail) after the V-11 stage may increase when a product containing dicamba ("Banvel", "Clarity", "Marksman") is applied to small corn under early stressful conditions. Be aware of this when applying tank mixes with dicamba to small corn (V-3 stage or smaller) under stressful conditions. See ENVIRONMENTAL CONDITIONS for a description of these stressful conditions

• Apply DuPont DPX-79406 75DF/"Hornet" tank mixes within four hours or product degradation may occur. If the tank mix cannot be applied within four hours, add 1 quart per acre of 10-34-0 or 11-37-0 or a commercial buffer which will raise the tank pH to 7.0. If 10-34-0 or 11-37-0 was used as the ammonium nitrogen fertilizer source, do not add additional fertilizer or buffer.

Other than the exceptions noted, and in addition to the tank mix partners and rates indicated in the table above, DPX-79406 75DF may be tank mixed or followed with sequential applications of other products registered for use in field corn. DPX-79406 75DF may be applied in tank mix combinations with labeled rates of other products provided:

- The tank mix product is labeled for the same timing, method of application, adjuvants, and use restrictions as DPX-79406 75DF.
- The tank mixture is not specifically prohibited on the label of the tank mix product.
- The tank mix combination is compatible as determined by a "jar test" described in the TANK MIX COMPATIBILITY TESTING section below.

Tank Mixing Precautions:

- Weed control and crop response with tank mixtures not specifically recommended in this label are the responsibility of the user and manufacturer of the tank mix product.

- Read and follow all applicable use directions, precautions, and limitations specified on the respective product labels and fact sheets.
- Do not exceed labeled application rates. Do not tank mix DPX-79406 75DF with other products that contain the same active ingredients as DPX-79406 75DF (nicosulfuron and rimsulfuron) unless the label of either tank mix partner specifies the maximum rate that may be used.

TANK MIX COMPATIBILITY TESTING

Perform a jar test prior to tank mixing to ensure compatibility of DPX-79406 75DF and other pesticides. Use a clear glass quart jar with lid and mix the tank mix ingredients in their relative proportions. Invert the jar containing the mixture several times and observe the mixture for approximately 1/2 hour. If the mixture ball-up, forms flakes, sludges, gels, oily film or layers, or other precipitates, it is not compatible and the tank mix combination should not be used.

SEQUENTIAL ACCENT® APPLICATIONS

Apply DuPont ACCENT® Herbicide 14 or more days after DPX-79406 75DF applications to control grasses that may emerge later in the season. Refer to the ACCENT® label for grass species controlled, proper size of weeds, rates, corn sizes, and other information. When following a DPX-79406 75DF application, do not use more than 1 oz ACCENT® per acre.

A sequential application of ACCENT® will effect crop rotation intervals to certain sensitive crops, such as sugarbeets. For maximum crop rotation flexibility, consult the CROP ROTATION section before applying ACCENT® to fields previously treated with DPX-79406 75DF.

CULTIVATION

A timely cultivation may be necessary to control suppressed weeds, or weeds that emerge after an application of DPX-79406 75DF in the absence of an activating rainfall.

- Optimum timing for cultivation is 7-14 days after DPX-79406 75DF application or upon seeing the establishment of new weeds.

ENVIRONMENTAL CONDITIONS AND BIOLOGICAL ACTIVITY

DPX-79406 75DF provides best results when applied to young, actively growing weeds. Applications made during warm, moist conditions (70°F or more) and adequate soil moisture both before and after application maximizes performance.

The degree and duration of control depend on spray coverage, activating rainfall, weed spectrum, weed size, growing conditions before and after treatment, soil moisture, and adjuvant selection.

Adequate soil moisture is required for optimum activity. Rainfall within 5-7 days will enhance DPX-79406 75DF residual activity. A timely cultivation may be required for maximum weed control without an activating rain.

DPX-79406 75DF is rainfast in 4 hours.

Treating weeds that exceed maximum label height or that are under stress may result in incomplete control. Poor weed control or crop injury may result from applications made to plants under stress from:

- abnormally hot or cold weather
- environmental conditions such as drought, water-saturated soils, hail damage, or frost
- disease, insect, or nematode injury
- prior herbicide, or carryover from a previous year's herbicide application

Severe stress from conditions immediately following application may also result in crop injury or poor weed control. Stress affects all weeds, but especially weeds such as woolly cupgrass, green and yellow foxtail, and wild proso millet.

If the corn or grass weeds are under stress, delay application until stress passes and both weeds and corn resume active growth.

Apply DuPont DPX-79406 75DF when minimum nighttime temperatures are above 40°F and the maximum daytime temperatures are below 92°F to maximize performance and minimize the potential for crop injury.

Applications made during or immediately following periods of large day/night temperature fluctuations or where daytime temperatures do not exceed 50°F may decrease weed control and increase the potential for crop injury.

DPX-79406 75DF rapidly inhibits the growth of susceptible weeds, reducing weed competition within as little as 6 hours after application. Susceptible plants are controlled in 7-21 days.

Ground application of DPX-79406 75DF to dry, dusty fields may reduce weed control in wheel track areas.

SOIL INSECTICIDE INTERACTION INFORMATION

Before using DPX-79406 75DF, ensure that it is compatible with any other insecticides previously applied to the corn crop.

Conventional Field Corn and "IT" Hybrids

Soil Insecticides	Application Method	Soil O.M.	DPX-79406 75DF Use Precautions
"Counter 15G"	All	All	Do not use.
"Counter 20CR"	In furrow at planting	All	Do not use.
	Over the row at cultivation	All	Do not use.
	T-band or Surface band	≤4%	May cause unacceptable injury.
	T-band or Surface band	>4%	May result in temporary injury.
"Thimet"	All labeled methods	All	May cause unacceptable injury.
Dyfonate	All labeled methods	All	May result in temporary injury.
"Lorsban"	All labeled methods	All	May result in temporary injury.
"Aztec"	All	All	No use precautions
"Force"	All	All	No use precautions
"Fortress"	All	All	No use precautions.

Herbicide-Resistant Field Corn

- DPX-79406 75DF may be used on fields treated with "Counter 15G" or "Counter 20CR" (applied in-furrow, T- or surface-banded) if the field has been planted with an imidazilone-resistant ("IR") hybrid such as Pioneer 35A19, 3417IR, or 3395IR, etc.
- For DPX-79406 75DF applied to "IMI" or "ClearFields" corn hybrids, not designated as "IR", follow directions above for Conventional and "IT" Field Corn.

CROP ROTATION

Rotational crops vary in their response to low concentrations of DPX-79406 75DF remaining in the soil. DPX-79406 75DF dissipates rapidly in warm, acidic, microbiologically active soils. The amount of DPX-79406 75DF which may be present in the soil depends on soil pH and organic matter content, elapsed time since application, crop production practices, and environmental factors.

Injury to rotational crops may occur in high-pH, cold soils if dry weather prevails between application and rotational crop planting.

For fields treated with sequential applications of DPX-79406 75DF and ACCENT® herbicide, consult the crop rotation intervals listed on the ACCENT® and DPX-79406 75DF labels. Use the most restrictive re-crop interval from either label.

The following rotational intervals must be observed when using DPX-79406 75DF:

DPX-79406 75DF ROTATIONAL CROP GUIDELINE

No soil pH restrictions

Crop	Rotational Interval in Months
Alfalfa	12
Cereals, Spring (wheat, barley, oats & rye)	8
Cereals, Winter (wheat, barley, oats & rye)	4
Corn (Field)	Anytime
Corn (Pop, Sweet, Seed)	10
Cotton	10
Dry Beans, Peas, Snap Beans	10
Oats	8
Sorghum	10
Soybeans	1/2 (15 days)
Sugar Beets	10
Sunflowers	11
Other Crops	18

APPLICATION INFORMATION

Many crops are highly sensitive to DPX-79406 75DF. All direct or indirect contact (such as spray drift) with crops other than field corn should be avoided (see also SPRAY DRIFT MANAGEMENT).

For all application systems, use 50-mesh or larger strainer screens.

Do not apply DPX-79406 75DF through any type of irrigation system.

GROUND APPLICATION

Broadcast Application

- Use a minimum of 15 gal of water per acre (15 GPA) for best performance. Use a minimum of 10 gal of water per acre (GPA) for light, scattered stands of grass.
- Use flat fan nozzles at 20–40 psi.
- Increase both spray volume and pressure as weed density and size increase.
- Do not use flood, hollow cone, rain drop, whirl chamber controlled droplet applicator (CDA) nozzles or air assisted sprayers. Unacceptable crop injury, excessive spray drift, or poor weed control may result.
- For proper spray coverage, adjust the boom and nozzle height according to manufacturers' specifications.
- Ensure that equipment is set up to avoid applying an excessive rate directly over the rows and into the corn plant whorl.
- Overlaps or starting, stopping, and turning while spraying may result in crop injury.

Band Application

- For band applications, use proportionately less spray mixture.
- To avoid crop injury, carefully calibrate the band applicator to not exceed the labeled rate.
- Carefully follow the manufacturer's instructions for nozzle type (flat fans), orientation, distance of nozzles from the crop and weeds, spray volumes, calibration and spray pressure.

AERIAL APPLICATION

- Use nozzle types and arrangements that will provide optimum spray distribution and maximum coverage at a minimum of 5 GPA.
- Do not apply during a temperature inversion, when winds are gusty, or when conditions favor poor coverage and/or off-target spray movement.

SPRAYER PREPARATION/CLEANUP

It is important that spray equipment is clean and free of previous pesticide deposits before using DPX-79406 75DF and then properly cleaned out following application. Clean all application equipment before applying DPX-79406 75DF. Follow the cleanup procedures specified on the label of the product previously sprayed. If no cleanup procedure is provided, use the procedure that follows. Immediately following applications of

DPX-79406 75DF, thoroughly clean all mixing and spray equipment to avoid subsequent crop injury.

Note:

- When cleaning spray equipment before applying DPX-79406 75DF, read and follow label directions for proper rinsate disposal of the product previously sprayed.
- Steam cleaning of aerial spray tanks will help to dislodge any visible pesticide deposits.
- When spraying or mixing equipment will be used over an extended period to apply multiple loads of DPX-79406 75DF, partially fill the tank with fresh water at the end of each day of spraying, flush the boom and hoses, and allow to sit overnight.

Cleanup Procedure

1. Drain the tank and thoroughly hose down the interior surfaces. Flush the tank, hoses, and boom with clean water for a minimum of 5 min.
2. Partially fill the tank with clean water and add one gal of household ammonia* (containing 3% active) for every 100 gal of water. Finish filling the tank with water, then flush the cleaning solution through the hoses, boom, and nozzles. Add more water to completely fill the tank and allow to agitate/recirculate for at least 15 min. Again, flush the hoses, boom, and nozzles with the cleaning solution, then drain the tank.
3. Repeat Step 2.
4. Remove the nozzles and screens and clean separately in a bucket containing the cleaning agent and water.
5. Thoroughly rinse the tank with clean water for a minimum of 5 min, flushing the water through the hoses and boom.

* Equivalent amounts of an alternate strength ammonia solution or a tank cleaner recommended in the DuPont bulletin "Sulfonylurea Herbicides, A Guide to Equipment Cleanout," may be used.

SPRAY DRIFT MANAGEMENT

The interaction of many equipment and weather-related factors determines the potential for spray drift. The applicator is responsible for considering all these factors when making application decisions.

AVOIDING SPRAY DRIFT IS THE RESPONSIBILITY OF THE APPLICATOR.

IMPORTANCE OF DROPLET SIZE

The most effective way to reduce drift potential is to apply large droplets (>150 - 200 microns). The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. The presence of sensitive species nearby, the environmental conditions, and pest pressure may affect how an applicator balances drift control and coverage. **APPLYING LARGER DROPLETS REDUCES DRIFT POTENTIAL, BUT WILL NOT PREVENT DRIFT IF APPLICATIONS ARE MADE IMPROPERLY OR UNDER UNFAVORABLE ENVIRONMENTAL CONDITIONS!** See **Wind, Temperature and Humidity, and Temperature Inversions** sections of this label.

Controlling Droplet Size - General Techniques

- **Volume** - Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets.
- **Pressure** - Use the lower spray pressures recommended for the nozzle. Higher pressure reduces droplet size and does not improve canopy penetration. **WHEN HIGHER FLOW RATES ARE NEEDED, USE A HIGHER-CAPACITY NOZZLE INSTEAD OF INCREASING PRESSURE.**
- **Nozzle Type** - Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles.

Controlling Droplet Size - Aircraft

- **Number of Nozzles** - Use the minimum number of nozzles with the highest flow rate that provide uniform coverage.
- **Nozzle Orientation** - Orienting nozzles so that the spray is emitted backwards, parallel to the airstream will produce larger droplets than other orientations.
- **Nozzle Type** - Solid stream nozzles (such as disc and core with swirl plate removed) oriented straight back produce larger droplets than other nozzle types.
- **Boom Length** - The boom length should not exceed 3/4 of the wing or rotor length - longer booms increase drift potential.
- **Application Height** - Application more than 10 ft above the canopy increases the potential for spray drift.

BOOM HEIGHT

Setting the boom at the lowest labeled height (if specified) which provides uniform coverage reduces the exposure of droplets to evaporation and wind. For ground equipment, the boom should remain level with the crop and have minimal bounce.

WIND

Drift potential increases at wind speeds of less than 3 mph (due to inversion potential) or more than 10 mph. However, many factors, including droplet size and equipment type determine drift potential at any given wind speed. **AVOID GUSTY OR WINDLESS CONDITIONS.**

Note: Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they affect spray drift.

TEMPERATURE AND HUMIDITY

When making applications in hot and dry conditions, set up equipment to produce larger droplets to reduce effects of evaporation.

TEMPERATURE INVERSIONS

Drift potential is high during a temperature inversion. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain close to the ground and move laterally in a concentrated cloud. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning.

13/14

Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

SHIELDED SPRAYERS

Shielding the boom or individual nozzles can reduce the effects of wind. However, it is the responsibility of the applicator to verify that the shields are preventing drift and not interfering with uniform deposition of the product.

WEED RESISTANCE

When herbicides with the same mode of action are used repeatedly over several years to control the same weed species in the same field, naturally-occurring resistant weed biotypes may survive a correctly applied herbicide treatment, propagate, and become dominant in that field. These resistant weed biotypes may not be adequately controlled. Cultural practices such as tillage, preventing weed escapes from going to seed, and using herbicides with different modes of action within and between crop seasons can aid in delaying the proliferation and possible dominance of herbicide resistant weed biotypes.

INTEGRATED PEST MANAGEMENT

DuPont recommends the use of Integrated Pest Management (IPM) programs to control pests. This product may be used as part of an Integrated Pest Management (IPM) program which can include biological, cultural, and genetic practices aimed at preventing economic pest damage. Application of this product should be based on IPM principles and practices including field scouting or other detection methods, correct target pest identification, population monitoring, and treating when target pest populations reach locally determined action thresholds. Consult your state cooperative extension service, professional consultants or other qualified authorities to determine appropriate action treatment threshold levels for treating specific pest/crop systems in your area.

IMPORTANT PRECAUTIONS

Injury to or loss of desirable vegetation may result from failure to observe the following:

- Do not apply DuPont DPX-79406 75DF or drain or flush application equipment on or near desirable trees or other plants, or on areas where their roots may extend, or in locations where the chemical may be washed or moved into contact with their roots.
- Do not use on lawns, walks, driveways, tennis courts, or similar areas.
- Prevent drift of spray to desirable plants.
- Do not contaminate any body of water.
- Thoroughly clean application equipment immediately after use. (See the Sprayer Cleanup section of this label for instructions.)

- Do not graze or feed forage, hay, or straw from treated areas to livestock within 30 days of DPX-79406 75DF application.
- Do not apply more than 3/4 oz per acre per season.

STORAGE AND DISPOSAL

Storage: Store product in original container only. Do not contaminate water, other pesticides, fertilizer, food or feed in storage. Store in a cool, dry place.

Product Disposal: Do not contaminate water, food, or feed by disposal. Waste resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: For Plastic Containers: Triple rinse (or equivalent) the container. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke. **For Fiber Sacks:** Completely empty fiber sack by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application equipment. Dispose of bags at an approved waste disposal facility, in accordance with Federal, state and local regulations. **For Fiber Drums With Liners:** Completely empty liner by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application equipment. Then dispose of liner in a sanitary landfill or by incineration if allowed by state and local authorities. If drum is contaminated and cannot be reused, dispose of in the same manner. **For Bags Containing Water Soluble Packets:** Do not reuse the outer box or the resealable plastic bag. When all water-soluble packets are used, the outer packaging should be clean and may be disposed of in a sanitary landfill or by incineration, or if allowed by state and local authorities, by open burning. If burned, stay out of smoke. If the resealable plastic bag contacts the formulated product in any way, the bag must be triple-rinsed with clean water. Add the rinsate to the spray tank and dispose of the outer wrap as described above. **For Metal Containers (non aerosol):** Triple rinse (or equivalent) the container. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

NOTICE TO BUYER: Purchase of this material does not confer any rights under patents of countries outside of the United States.

"Axiom", "Aztec" are registered trademarks of Bayer Crop Protection

"Balance PRO", "Regent" are registered trademarks of Aventis Crop Science

"Banvel", "Basagran", "Clarity", "Counter", "Frontier", "Laddok", "Marksman", "Prowl" are registered trademarks of BASF Corp.

"ClearField", "IMI", "Thimet" are registered trademarks of American Cyanamid

"Dual II", "Dual II Magnum", "Exceed", "Northstar", "Spirit" are registered trademarks of Novartis Crop Protection Inc.

"Force", "Surpass EC", "Topnotch" are registered trademarks of Zeneca Ag Products

"Fortress" is a registered trademark of Amvac Chemical Co.

"Harness", "Permit" are registered trademarks of Monsanto Co.

"Hornet", "Hornet WDG", "Lorsban" are registered trademarks of Dow Agro Sciences

D - 727 122000

14/14

**LIMITATION OF
WARRANTY AND LIABILITY**

NOTICE: Read This Limitation of Warranty and Liability Before Buying or Using This Product. If the Terms Are Not Acceptable, Return the Product at Once, Unopened, and the Purchase Price Will Be Refunded.

It is impossible to eliminate all risks associated with the use of this product. Such risks arise from weather conditions, soil factors, off target movement, unconventional farming techniques, presence of other materials, the manner of use or application, or other unknown factors, all of which are beyond the control of DuPont. These risks can cause: ineffectiveness of the product; crop injury, or; injury to non-target crops or plants.

DuPont does not agree to be an insurer of these risks. **WHEN YOU BUY OR USE THIS PRODUCT, YOU AGREE TO ACCEPT THESE RISKS.**

DuPont warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for the purpose stated in the Directions for Use, subject to the inherent risks described above, when used in accordance with the Directions for Use under normal conditions.

DUPONT MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR OF MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

IN NO EVENT SHALL DUPONT OR SELLER BE LIABLE FOR ANY INCIDENTAL, CONSEQUENTIAL OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. BUYER'S OR USER'S BARGAINED-FOR EXPECTATION IS CROP PROTECTION. THE EXCLUSIVE REMEDY OF THE USER OR BUYER AND THE EXCLUSIVE LIABILITY OF DUPONT OR SELLER, FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY OR CONTRACT, NEGLIGENCE, TORT OR STRICT LIABILITY), WHETHER FROM FAILURE TO PERFORM OR INJURY TO CROPS OR OTHER PLANTS, AND RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT, OR AT THE ELECTION OF DUPONT OR SELLER, THE REPLACEMENT OF THE PRODUCT.

DuPont or its Ag Retailer must have prompt notice of any claim so that an immediate inspection of buyer's or user's growing crops can be made. Buyer and all users shall promptly notify DuPont or a DuPont Ag Retailer of any claims, whether based on contract, negligence, strict liability, other tort or otherwise or be barred from any remedy.

This Limitation of Warranty and Liability may not be amended by any oral or written agreement.

For product information call: 1-888-6-DUPONT

Internet address: www.dupont.com/ag/us

© 2000 E. I. du Pont de Nemours and Company, Agricultural Products, Wilmington, Delaware 19898

NEXT

LABEL

SUPPLEMENTAL LABELING

**DuPont Agricultural
Products**

DRAFT

**DPX-79406 75 DF HERBICIDE
PLUS DISTINCT¹ HERBICIDE,
TOUGH² HERBICIDE OR
STINGER³ HERBICIDE AND COC**

"..... A Growing Partnership With Nature"

DPX-79406 75 DF HERBICIDE

EPA Reg. No. 352-572

TANK MIXTURES WITH DISTINCT, TOUGH OR STINGER HERBICIDES PLUS CROP OIL CONCENTRATE FOR WEED CONTROL IN FIELD CORN

DIRECTIONS FOR USE

DuPont DPX-79406 75 DF Herbicide may be tank mixed with 1-2 oz of Distinct, 1 pt of Tough, or 2-4 oz of Stinger for additional control of broadleaf weeds when applied to field corn. Do not apply to field corn grown for seed, to popcorn or to sweet corn.

WHEN TO APPLY

Postemergence applications of DPX-79406 75 DF plus Distinct, Tough or Stinger should be made to corn that is 4-12" tall and is exhibiting fewer than 6 leaf-collars. See the DPX-79406 75 DF label for sizes of grasses and other broadleaf weeds controlled.

HOW TO USE

DPX-79406 75 DF performs best when applied to young, actively growing grasses and weeds. An activating rainfall (>1/2") is required shortly after application for maximum DPX-79406 75 DF residual activity. Use a minimum of 15 gallons of water per acre (15 GPA) for best results. Do not apply less than 1/2 ounce of DPX-79406 75 DF per acre. Apply DPX-79406 75 DF plus Distinct, Tough or Stinger at the following rates:

Tank Mix Herbicide	Rate	Adjuvant
Distinct	1-2 oz	COC + N Fertilizer
Tough	1 pt	COC + N Fertilizer
Stinger	2-4 oz	COC + N Fertilizer

SPRAY ADDITIVES

Add an ammonium nitrogen fertilizer such as 28-0-0 (2-4 qts/acre), 10-34-0 (1-2 qts/acre) or spray grade ammonium sulfate (21-0-0) at 2-4 lbs/acre plus crop oil concentrate (COC) at 1% v/v (1 gal/100 gal spray solution).

IMPORTANT PRECAUTIONS

Refer to the DPX-79406 75 DF and Distinct, Tough or Stinger labels for additional use directions, restrictions, rotational crop intervals and precautions. The most restrictive provision on either label shall apply.

Do not apply DPX-79406 75 DF through any type of irrigation system.

IMPORTANT

BEFORE USING THESE PRODUCTS, READ AND FOLLOW ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA-REGISTERED LABELS.

This bulletin contains new or supplemental instructions for use of this product which do not appear on the EPA-registered package label and may contain recommendations which are not appropriate for all geographies and weed control situations. Consult local DuPont fact sheets, technical bulletins, and service policies prior to following the recommendations contained in this label.


This labeling must be in the possession of the user at the time of pesticide application.

¹ Registered trademark of BASF

² Registered trademark of Novartis

³ Registered trademark of Dow AgroSciences

DR-058 011501



NEXT

LABEL

SUPPLEMENTAL LABELING

**DuPont Agricultural
Products**

DRAFT

**DPX-79406 75 DF HERBICIDE
PLUS DISTINCT¹ HERBICIDE OR
CLARITY¹ HERBICIDE AND NIS**

"..... A Growing Partnership With Nature"

DPX-79406 75 DF HERBICIDE

EPA Reg. No. 352-572

TANK MIXTURES WITH DISTINCT HERBICIDE OR CLARITY HERBICIDE AND NON-IONIC SURFACTANT FOR WEED CONTROL IN FIELD CORN

DIRECTIONS FOR USE

DuPont DPX-79406 75 DF Herbicide may be tank mixed with 2 oz of Distinct Herbicide or 4-8 oz Clarity Herbicide for additional control of broadleaf weeds when applied to field corn. Do not apply to field corn grown for seed, to popcorn or to sweet corn.

WHEN TO APPLY

Postemergence applications of DPX-79406 75 DF plus Distinct or Clarity should be made to corn that is 4-12" tall and is exhibiting fewer than 6 leaf-collars. See the DPX-79406 75 DF label for sizes of grasses and other broadleaf weeds controlled.

HOW TO USE

DPX-79406 75 DF performs best when applied to young, actively growing grasses and weeds. An activating rainfall (>1/2") is required shortly after application for maximum DPX-79406 75 DF residual activity. Use a minimum of 15 gallons of water per acre (15 GPA) for best results. Do not apply less than 1/2 ounce of DPX-79406 75 DF per acre. Apply DPX-79406 75 DF plus Distinct or Clarity at the following rates:

Tank Mix Herbicide	Rate	Adjuvant
Distinct	2 oz	NIS + N Fertilizer
Clarity	4-8 oz	NIS + N Fertilizer

SPRAY ADDITIVES

Add an ammonium nitrogen fertilizer such as 28-0-0 (2-4 qts/acre), 10-34-0 (1-2 qts/acre) or spray grade ammonium sulfate (21-0-0) at 2-4 lbs/acre plus non-ionic surfactant (NIS) at 0.25% v/v (1 qt/100 gal spray solution).

IMPORTANT PRECAUTIONS

Refer to the DPX-79406 75 DF and Distinct or Clarity labels for additional use directions, restrictions, rotational crop intervals and precautions. The most restrictive provision on either label shall apply.

Do not apply DPX-79406 75 DF through any type of irrigation system.

IMPORTANT BEFORE USING THESE PRODUCTS, READ AND FOLLOW ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA-REGISTERED LABELS.

This bulletin contains new or supplemental instructions for use of this product which do not appear on the EPA-registered package label and may contain recommendations which are not appropriate for all geographies and weed control situations. Consult local DuPont fact sheets, technical bulletins, and service policies prior to following the recommendations contained in this label.

This labeling must be in the possession of the user at the time of pesticide application.

¹ Registered trademark of BASF.

DR-060 011501

