

Best copy available

352-574

07/14/95

H-62755

1/10




Synchrony[®] STS[®] SP

herbicide

ACCEPTED

JUL 14 1995

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 352-574



“..... A Growing Partnership With Nature”

2/10




TABLE OF CONTENTS

- Directions For Use 2
- Application Information 2
 - Timing To Crop Stage 2
 - Timing To Weeds 2
 - Rate and Weeds Controlled 2
 - Split Applications 3
- Where SYNCHRONY STS SP
 - May Be Used 3
 - No-Till/Conservation Till 3
 - Spray Additives 3
 - Tank Mix Applications 3
- Mixing Instructions 4
- Application Equipment 5
 - Ground Application 5
 - Aerial Application 5
- Environmental Conditions and
 - Biological Activity 5
- Rotational Crop Guidelines 5
- Sprayer Preparation and Cleanup 6
- Spray Drift Management 6
- Important Precautions 7
- Information on Resistant Weeds 7
- Storage and Disposal 8
- Notice to Buyer 8
- Notice of Warranty 8



Synchrony® STS® SP

herbicide


Dispersible Granules

Active Ingredients	By Weight
Chlorimuron Ethyl	
Ethyl 2-[[[(4-chloro-6-methoxypyrimidin-2-yl)amino]carbonyl]amino]sulfonyl]benzoate	31.8%
Thifensulfuron methyl	
Methyl 3-[[[(4-methoxy-6-methyl-1,3,5-triazin-2-yl)amino]carbonyl]amino]sulfonyl]-2-thiophenecarboxylate	10.2%
INERT INGREDIENTS	58.0%
TOTAL	100.0%

EPA Reg. No. 352 - 574

U.S. Patent No. 4,394,506, 4,547,215 & 4,481,029

FOR USE ONLY ON SOYBEAN VARIETIES DESIGNATED AS "STS".



KEEP OUT OF REACH OF CHILDREN

CAUTION

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Caution! May be harmful if absorbed through skin. Causes (moderate) eye injury (irritation). Avoid contact with skin, eyes, and clothing. Wash thoroughly with soap and water after handling.

STATEMENT OF PRACTICAL TREATMENT

IF IN EYES: Flush eyes with plenty of water. Call a physician if irritation persists.

IF ON SKIN: Wash with plenty of soap and water. Get medical attention.

For medical emergencies involving this product, call toll-free 1-800-441-3637.

PERSONAL PROTECTIVE EQUIPMENT

Applicators and other handlers must wear:

Long-sleeved shirt and long pants.

Waterproof gloves.

Shoes plus socks.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENGINEERING CONTROL STATEMENTS

When handlers use closed systems, enclosed cabs or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR part 170 Section 170.240 (d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

USERS SHOULD: Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls.
- Waterproof gloves.
- Shoes plus socks.

Do not apply this product through any type of irrigation system.

FOR USE ONLY ON SOYBEAN VARIETIES DESIGNATED AS "STS".

These "STS" varieties must be purchased from an authorized seed supplier.

The "STS" designation indicates the soybean variety contains a proprietary trait that enhances the soybeans natural tolerance to DuPont soybean sulfonylurea herbicides. Information on "STS" soybean varieties may be obtained from your seed supplier or DuPont representative.

THIS PRODUCT IS NOT A PESTICIDE FOR ALL PURPOSES. IT IS A HERBICIDE FOR SOYBEAN VARIETIES DESIGNATED AS "STS".

This product is a water-dispersible granule containing 25% active ingredient in a premeasured water soluble pack.

APPLICATION INFORMATION

Du Pont's SYNCHRONY STS SP herbicide is a dispersible granule formulation premeasured in 2.0 ounce packs which readily dissolve in water.

Timing To Crop Stage

SYNCHRONY STS SP performs best when applied to small weeds a few weeks after planting.

Timing To Weeds

- Apply SYNCHRONY STS SP when weeds are young and actively growing (after the first true leaves have expanded, but before the weeds exceed the size indicated below).
- Applications made to weeds larger than the sizes indicated below, or to weeds under stress, may result in unsatisfactory control.
- Applications made after July 15 may result in less than satisfactory weed control due to the large sizes of the weeds and increased weather stress.

Rate

When applied as directed, SYNCHRONY STS SP will control the following weeds:

Weed	Height (in inches)
Beggarticks (Bidens sp)	2-6
Bristly Starbur	2-3
Burcucumber*	2-3
Cocklebur	2-8
Common Milkweed (above ground portion)	2-6
Cowpea	2-5
Florida Beggardweed	2-5
Florida Pusley	1-3
Herp Sesbania	2-5
Lambquarters	2-4
Jerusalem Artichoke (above ground portion)	2-6
Jimsonweed	2-5
Marestail	2-5
Morningglory* (annual)	
Entireleaf	1-3
Ivyleaf	1-3
Pitted	1-3
Smallflower	1-3
Tall	1-3
Mustard	5**
Pigweeds	
Redroot (Rough)	2-8
Others	2-8
Ragweed	
Common	2-4
Giant*	2-4
Sicklepod*	1-3
Smartweed	
Ladysthumb	2-8
Pennsylvania	2-8
Sunflower	2-8
Tall Waterhemp	2-8
Wild Poinsettia	1-2
Yellow Nutsedge	2-3
Velvetleaf	2-8

* See Split Applications section.

** Diameter

5/10


When applied as directed, SYNCHRONY STS SP will suppress the following weeds:

Weed	Height (in inches)
Buffalograss	2-6
Canada Thistle	2-4
Kochia	1-3
Spurred Anoda	1-3
Purple Nutsedge	2-4
Venice Mallow	1-3

* See Split Applications section.

** Diameter

Split Applications



A follow up application of DuPont Classic Herbicide may be made 2-3 weeks after a SYNCHRONY STS SP application to control weeds with multiple germination flushes or suppressed weeds such as burcucumber, cowpea, giant ragweed, morningglory, and sicklepod. See Rotational Crop Guidelines for intervals following sequential applications.

Cultivation

Cultivation approximately 14 days after application will help control suppressed weeds.

Where SYNCHRONY STS SP May Be Used

SYNCHRONY STS SP may be used without any soil pH restrictions in the states of Alabama, Arkansas, Delaware, Florida, Georgia, Illinois, Indiana, Iowa (except in the Clarion-Nicollet-Webster and Hamburg-Ida-Monona soil associations and the historic floodplain of the Missouri River), Kansas, Kentucky, Maryland, Mississippi (except in the Blackbelt soils), Missouri, New Jersey, North Carolina, Ohio, Oklahoma, Pennsylvania, South Carolina, Tennessee, Virginia, and West Virginia.


In the states of Michigan, Nebraska, and in Iowa on Clarion-Nicollet-Webster and Hamburg-Ida-Monona soils and in the historic flood plain of the Missouri River, and on the Blackbelt soils of Mississippi, use SYNCHRONY STS SP on soils having a pH of 7.0 or less.

Soil pH may vary greatly within a single field. It may be affected by soil amendments or disturbed soils. Determine soil pH by laboratory analysis using a 1:1 soil:water suspension. SYNCHRONY STS SP should not be used on soils with a history of nutrient deficiency (such as iron chlorosis). Crop injury may occur.

No-Till/Conservation Till

SYNCHRONY STS SP may be used for postemergence weed control in no-till/conservation till operations. It may be used alone, in a tank mix for postemergence broadleaf weed control, or tank mixed with postemergence grass herbicides such as Du Pont's ASSURE II Herbicide for total postemergence weed control. SYNCHRONY STS SP may be used in combination with premerge applications of Du Pont's CANOPY® herbicide for a pre-post No-Till herbicide program.

Spray Additives



Crop Oil Concentrate

- Apply crop oil concentrate at 8 pt per 100 gal of spray solution (1.0% v/v).
- Use a good-quality, petroleum-based or methylated seed oil-based crop oil concentrate with at least 14% emulsifiers and 80% oil.

Ammonium Nitrogen Fertilizer


In addition to a crop oil concentrate, an ammonium nitrogen fertilizer is recommended, and required where velvetleaf is present.

- Use a high-quality, liquid nitrogen fertilizer such as 28-0-0 at a rate of 4-8 pt per acre, or a 10-34-0 at a rate of 2-4 pt per acre.
- Alternatively, a high-quality, sprayable grade of ammonium sulfate (21-0-0) may be used at a rate of 2-4 lb per acre.
- Use the lower rate of fertilizer for spray volumes of less than 15 gal per acre.

TANK MIX APPLICATIONS

Do not tank mix SYNCHRONY STS SP with any other pesticide or spray adjuvant except as specified on this or other supplemental labeling.

SYNCHRONY STS SP AND POSTEMERGENCE GRASS HERBICIDES



- For best results apply SYNCHRONY STS SP 7 days before or 1 day after the grass herbicide. Refer to the grass herbicide label for precautions and specific use information.
- Include a crop oil concentrate with the tank mix of SYNCHRONY STS SP and postemergence grass herbicides such as ASSURE II. Use the rate listed in the Soybean Spray Additives section.
- Under certain conditions SYNCHRONY STS SP may reduce the activity of the grass herbicide. The broadleaf activity of SYNCHRONY STS SP will not be affected.
- SYNCHRONY STS SP may be tank mixed with other grass herbicides such as Poast Plus®, Select®, Fusion®, or Fusilade® 2000. Read and follow label directions for any tank mix product.

The types of grass present determine the amount of ASSURE II to be tank mixed with SYNCHRONY STS SP. When applied as directed, a tank mix of SYNCHRONY STS SP and ASSURE II will control the following grasses:

SYNCHRONY STS SP + 5 oz of ASSURE II per acre

Grass	Height (Inches)
Volunteer Corn	6 - 18
Shattercane	6 - 12
Giant Foxtail	2 - 4 (pretillcr)
Seedling Johnsongrass	2 - 8

6/10

SYNCHRONY STS SP + 7 oz of ASSURE II per acre

Grass	Height (Inches)
Giant Foxtail	4 - 8
Wild Proso Millet	2 - 6

SYNCHRONY STS SP + 8 oz of ASSURE II per acre

Grass	Height (Inches)
Crowfoot Grass	2 - 6
Fall Panicum	2 - 6
Green Foxtail	2 - 4
Bristly Foxtail	2 - 4
Goosegrass	2 - 4
Ichgrass	2 - 8
Field Sandbur	2 - 6
Sprangletop	2 - 6
Volunteer Cereals	2 - 6
Wild Oats	2 - 6
Witchgrass	2 - 6

SYNCHRONY STS SP + 10 oz of ASSURE II per acre

Grass	Height (Inches)
Junglerice	2 - 6
Rhizome Johnsongrass	10 - 24

SYNCHRONY STS SP and 2,4-DB


- Applications to morningglory species must be made before the weeds are 4" tall.



- Applications of SYNCHRONY STS SP and 2,4-DB must include a nonionic surfactant at the rate of 2 pints per 100 gal (minimum of 0.125% v/v of actual nonionic surfactant) or crop oil concentrate at 8 pints per 100 gal of spray solution (1.0% v/v). See Spray Additives.
- Apply a tank mix of SYNCHRONY STS SP and 2,4-DB by ground only.
- In Kansas and Missouri (except the bootheel area), when conditions are excessively hot and dry (> 90°F and < 30% relative humidity), make applications at the rate of 2 fl oz of 2,4-DB and SYNCHRONY STS SP.
- The use of crop oil concentrate may increase temporary injury to soybeans.

Some crop response may occur 5-7 days after application of SYNCHRONY STS SP and 2,4-DB to soybeans under stress. Temporary yellowing, leaf crinkling, and/or soybean growth retardation may occur following application of SYNCHRONY STS SP and 2,4-DB. Under favorable growing conditions, the crop will quickly recover.


Consult the 2,4-DB label for use precautions.

SYNCHRONY STS SP AND COBRA¹

Weed	Height (Inches)
Prickly Sida (teaweed)	up to 1 inch
Cocklebur	2 - 6
Hemp Scabania	2 - 4
Morningglory	
Pitted	2 - 3
Ivyleaf	2 - 3
Entireleaf	2 - 3

SYNCHRONY STS SP and "Cobra" Application Information


- Include a nonionic surfactant at 2 pt per 100 gal of spray solution (minimum of 0.125% v/v actual surfactant).



- Tank mix applications of SYNCHRONY STS SP and "Cobra" may not control weeds listed on the SYNCHRONY STS SP label as completely as applications of SYNCHRONY STS SP alone.
- Applications with less than 20 gal per acre may result in erratic weed control.

MIXING INSTRUCTIONS

Soluble Packs are contained in waterproof, resealable plastic bags, with plastic bags enclosed in a cardboard box. The individual Soluble Packs will dissolve completely in water. Open the outer resealable plastic bag, remove the number of required Soluble Packs for the application rate of one 2.0 ounce Soluble Pack per 4 treated acres, and drop them into the spray tank as directed below.

Soluble Pack Handling Precautions:

Excessive handling of the packs, or exposure to moisture, will cause breakage.

Do not touch the packs with wet hands or place them on wet surfaces.

Protect unused Soluble Packs by resealing them in the plastic bag.

To Use the Soluble Packs:

1. Fill the tank 1/4 to 1/3 full of water.
2. While agitating, add the required number of SYNCHRONY STS SP Soluble Packs (one 2.0 ounce Soluble Pack per 4 acres).
3. The packs should dissolve completely within 5 minutes. Continue adequate agitation.
4. SYNCHRONY STS SP should be thoroughly mixed with water in the spray tank before adding any other material. As the tank is filling and after the Soluble Packs have dissolved, add (in order): other herbicide(s), the required spray adjuvant, the required nitrogen fertilizer.

7/10

5. Apply SYNCHRONY STS SP spray preparation within 24 hours of mixing to avoid product degradation.

6. If the mixture has settled, thoroughly reagitrate before using.

APPLICATION EQUIPMENT

Many crops are sensitive to SYNCHRONY STS SP. All direct or indirect contact (such as spray drift) with crops other than soybeans should be avoided.

Ground Application

(See Also Spray Drift Management)

Broadcast Application

- Use a minimum of 10 gal water per acre.
- Under heavy weed pressure or dense crop foliage, increase minimum spray volume to 15-25 gal per acre.
- Use flat fan nozzles at 25-60 psi for SYNCHRONY STS SP applications.
- Do not use flood, rain drop, whirl chamber, or controlled droplet applicator (CDA) type nozzles. Unacceptable crop injury, excessive spray drift, or poor weed control may result.
- For proper spray coverage adjust the boom and nozzle height according to the specifications listed by the nozzle manufacturer.

Band Application

- Because band applicators spray a narrower area than broadcast applicators, use proportionately less spray solution for band applications.
- Carefully calibrate the band applicator to not exceed the labeled rate.
- Flat fan nozzles are preferred.
- Carefully follow the nozzle manufacturer's instructions for nozzle orientation, distance of the nozzles from the crop and weeds, spray volumes, calibration, and spray pressure for band applications.
- For additional information on row banders, see Du Pont's bulletin, "Application Accuracy - Row Banders."

Aerial Application (See Also Spray Drift)

- Use nozzle types and arrangements that will provide optimum spray distribution and maximum coverage at 3-5 gal per acre.
- Use a minimum of 3 gal water per acre. Under heavy weed pressure or dense crop foliage, increase the minimum spray volume to 5 gal per acre.
- Do not apply during a temperature inversion, when winds are gusty, or when other conditions could produce poor coverage and/or off-target spray movement.

CULTIVATION


Do not cultivate within 7 days of application. Cultivation may put weeds under stress by pruning roots, thus diminishing control.

The best time to cultivate is approximately 14 days after application.

ENVIRONMENTAL CONDITIONS AND BIOLOGICAL ACTIVITY

SYNCHRONY STS SP rapidly inhibits the growth of susceptible weeds. Leaves of susceptible plants yellow 3-5 days after application, followed, in controlled plants, by the death of the growing point. SYNCHRONY STS SP will provide complete

control of susceptible weeds in 7-21 days. Suppressed plants may remain green but will be stunted and noncompetitive.




Stress affects some weeds, such as pigweed, more than others. Delay application until stress passes and weeds start to grow again.

Severe stress (drought, disease, insect damage, or nutrient deficiency such as iron chlorosis) following application may also result in poor weed control.

Do not apply SYNCHRONY STS SP if rain is expected within 1 hour or weed control may decrease.

ROTATIONAL CROP GUIDELINES



Rotational Intervals Following the Use of SYNCHRONY STS SP

Group A States: AL, AR, FL, GA, LA, MS*, NC, SC, TN, TX.

Group B States: DE, IA*, IL, IN, KS, KY, MD, MI, MO, NE, NJ, OH, OK, PA, VA, W.VA.

* Refer to "Where SYNCHRONY STS SP may be used" section of label.

INTERVAL 1

Following SYNCHRONY STS SP Application

- One application of SYNCHRONY STS SP in Group A or Group B states.
OR
- A single application of SYNCHRONY STS SP followed by a sequential application of Classic in Group B States having a soil pH not greater than 6.8.
OR
- A single application of SYNCHRONY STS SP following CANOPY*** (not more than 6 oz) or GEMINI*** (not more than 10 oz) or PREVIEW*** (not more than 8 oz) or NEW LOROX PLUS*** (not more than 16 oz) on a soil pH of 6.8 or less.
OR
- A single application of SYNCHRONY STS SP following any rate of CANOPY*** or GEMINI*** in Group A states.
OR

8/10

- A single application of SYNCHRONY STS SP following not more than 22 oz of GEMINI in the states of DE, MD, NJ, PA on soils with a pH of 6.8 or less.

INTERVAL 2

- A single application of SYNCHRONY STS SP followed by a sequential application of 1/2 oz Classic in Group B states with a soil pH greater than 6.8.

INTERVAL 3

- A single application of SYNCHRONY STS SP following CANOPY (7 oz or more), PREVIEW (9 oz or more), NEW LOROX PLUS (17 oz or more), or GEMINI (11 oz or more) in Group B states.
- OR
- Where a single application of SYNCHRONY STS SP follows CANOPY or GEMINI used on soils with a pH of 6.9 to 7.5 (where allowed).

Rotational Intervals following the use of SYNCHRONY STS SP*

Crop	Interval 1*	Interval 2*	Interval 3*
Soybeans	Anytime	Anytime	Anytime
Cereal Grains	3	3	3
Ryegrass	3	3	3
Field Corn**	9	9	15
IR Field Corn	8	8	8
Peanut	6	15	15
(if following Canopy)	9		
Alfalfa	11	15	15
Clover	11	15	15
Cotton	9	15	15
Dry Beans	9	15	15
Rice	9	15	15
Sorghum	11	15	15
Tobacco (transplant)	9	15	15
Tomato (transplant)	9	15	15

- * If SYNCHRONY STS SP is applied after August 1, extend recrop interval 2 months on corn, cotton, rice, and sorghum or rotational crop may be injured.
- ** For the Rotational Crop Guidelines, "Field Corn" refers only to popcorn and corn grown for grain, silage, or seed.
- *** Consult the rotational crop interval sections of the CANOPY, PREVIEW, and GEMINI, NEW LOROX PLUS labels. Some crop rotations may be longer than indicated because of metribuzin or linuron restrictions. Use the longest rotational crop interval.

The season before planting any crop not listed, a successful field bioassay must be completed. The field bioassay detects small quantities of SYNCHRONY STS SP that can remain in the soil and injure rotational crops. A successful field bioassay means growing to maturity a test strip of crop(s) intended for production the following year. The test strip should cross the whole field, including knolls and low areas. Suggested interval before initiating bioassay is 9 months for Group A and 15 months for Group B states.

SPRAYER PREPARATION AND CLEANUP

Prior to application of SYNCHRONY STS SP, start with clean, well maintained application equipment. Immediately

following application, thoroughly clean all application equipment. Postponing action, even for a few hours, only makes effective cleanup more difficult. Failure to clean spraying equipment thoroughly may result in injury to subsequently sprayed crops.

When spraying multiple loads of SYNCHRONY STS SP over an extended period of time, rinse the equipment with clean water at the end of the day. Leave water in the equipment overnight to prevent deposits from drying on surfaces.

When applications of SYNCHRONY STS SP are completed and prior to using the sprayer and associated equipment for other products or for crops other than soybeans, thoroughly clean the equipment using the procedure below.

- STEP 1. Drain spray equipment. Thoroughly rinse sprayer, and flush hoses, boom and nozzles with clean water. Loosen and physically remove visible deposits.
- STEP 2. Fill the sprayer with clean water and add household ammonia (one gallon of 3% active for every 100 gallons of water) or correct amount of a DuPont approved cleaner*. Flush hoses, boom and nozzles. Turn off the boom and top off the tank with clean water. Circulate through the spraying system for 15 minutes. Flush the hoses, boom and nozzles with the cleaning solution. Drain the tank.
- STEP 3. Remove and clean nozzle, screens and strainers in a bucket of fresh cleaner and water.
- STEP 4. Repeat STEP 2.
- STEP 5. Thoroughly rinse the sprayer, hoses, boom and nozzles with clean water, several times.

Clean all other associated application equipment. Take all necessary safety precautions when cleaning equipment. Do not clean near wells, water sources or near desirable vegetation. Dispose of waste rinse water in accordance with local regulations.

* For additional information on sprayer cleanup and a listing of DuPont-approved cleaners, see DuPont Bulletin "A Guide To Application Equipment Cleanout For DuPont Sulfonylurea Herbicides".

SPRAY DRIFT MANAGEMENT

The interaction of many equipment- and weather-related factors determines the potential for spray drift. The applicator is responsible for considering all these factors when making application decisions.

AVOIDING SPRAY DRIFT IS THE RESPONSIBILITY OF THE APPLICATOR.

IMPORTANCE OF DROPLET SIZE

The most effective way to reduce drift potential is to apply large droplets (>150 - 200 microns). The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. The presence of sensitive species nearby, the environmental conditions, and pest pressure may affect how an applicator balances drift control and coverage. **APPLYING LARGER DROPLETS REDUCES DRIFT POTENTIAL, BUT WILL NOT PREVENT DRIFT IF APPLICATIONS ARE MADE IMPROPERLY OR UNDER UNFAVORABLE ENVIRONMENTAL CONDITIONS!** See Wind, Temperature and Humidity, and Temperature Inversions sections of this label.

9/10

Controlling Droplet Size - General Techniques

- **Volume** - Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets.
- **Pressure** - Use the lower spray pressures recommended for the nozzle. Higher pressure reduces droplet size and does not improve canopy penetration. **WHEN HIGHER FLOW RATES ARE NEEDED, USE A HIGHER-CAPACITY NOZZLE INSTEAD OF INCREASING PRESSURE.**
- **Nozzle Type** - Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles.

Controlling Droplet Size - Aircraft

- **Number of Nozzles** - Use the minimum number of nozzles with the highest flow rate that provide uniform coverage.
- **Nozzle Orientation** - Orienting nozzles so that the spray is emitted backwards, parallel to the airstream will produce larger droplets than other orientations.
- **Nozzle Type** - Solid stream nozzles (such as disc and core with swirl plate removed) oriented straight back produce larger droplets than other nozzle types.
- **Boom Length** - The boom length should not exceed 3/4 of the wing or rotor length - longer booms increase drift potential.
- **Application Height** - Application more than 10 ft above the canopy increases the potential for spray drift.

BOOM HEIGHT

Setting the boom at the lowest labeled height (if specified) which provides uniform coverage reduces the exposure of droplets to evaporation and wind. For ground equipment, the boom should remain level with the crop and have minimal bounce.

WIND

Drift potential increases at wind speeds of less than 3 mph (due to inversion potential) or more than 10 mph. However, many factors, including droplet size and equipment type determine drift potential at any given wind speed. **AVOID GUSTY AND WINDLESS CONDITIONS.**

Note: Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they effect spray drift.

TEMPERATURE AND HUMIDITY

When making applications in hot and dry conditions, set up equipment to produce larger droplets to reduce effects of evaporation.

TEMPERATURE INVERSIONS

Drift potential is high during a temperature inversion. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain close to the ground and move laterally in a concentrated cloud. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

SHIELDED SPRAYERS

Shielding the boom or individual nozzles can reduce the effects of wind. However, it is the responsibility of the applicator to verify that the shields are preventing drift and not interfering with uniform deposition of the product.

AIR ASSISTED (AIR BLAST) FIELD CROP SPRAYERS

Air assisted field crop sprayers carry droplets to the target via a downward directed air stream. Some may reduce the potential for drift, but if a sprayer is unsuitable for the application and/or set up improperly, high drift potential can result. It is the responsibility of the applicator to determine that a sprayer is suitable for the intended application, is configured properly, and that drift is not occurring.

Note: Air assisted field sprayers can affect product performance by affecting spray coverage and canopy penetration. Consult the application equipment section of this label to determine if use of an air assist sprayer is recommended.

IMPORTANT PRECAUTIONS

SYNCHRONY STS SP Herbicide should be used only in accordance with recommendations on this label or in separately published DuPont recommendations. DuPont will not be responsible for losses or damage resulting from use of this product in any manner not specifically recommended by DuPont.

Do not spray in areas where the chemical may be washed or moved into contact with their roots.

Injury to or loss of desirable trees or vegetation may result from failure to observe the following:

- Do not apply SYNCHRONY STS SP or drain or flush equipment on or near desirable trees or other plants, on areas where their roots may extend, or in locations where the chemical may be washed or moved into contact with their roots.
- Do not use on lawns, walks, driveways, tennis courts or similar areas.
- Prevent spray drift to desirable plants.
- Thoroughly clean all application equipment immediately after use and prior to spraying crops other than soybeans.

SYNCHRONY STS SP Herbicide should be used only in accordance with recommendations on this label or in separately published DuPont recommendations. DuPont will not be responsible for losses or damage resulting from use of this product in any manner not specifically recommended by DuPont.

INFORMATION ON RESISTANT WEEDS

When herbicides with the same mode of action are used repeatedly over several years to control the same weed species in the same field, naturally-occurring resistant weed biotypes may survive a correctly applied herbicide treatment, propagate, and become dominant in that field. These resistant weed biotypes may not be adequately controlled. Cultural practices such as

10/10

tillage, preventing weed escapes from going to seed, and using herbicides with different modes of action within and between crop seasons can aid in delaying the proliferation and possible dominance of herbicide resistant weed biotypes.

STORAGE AND DISPOSAL

Storage: Store product in original container only. Do not contaminate water, other pesticides, fertilizer, food or feed in storage.

Product Disposal: Do not contaminate water, food or feed by disposal. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: Do not reuse the outer box or the resealable plastic bag. When all water-soluble packets are used, the outer packaging should be clean and may be disposed of in a sanitary landfill or by incineration, or if allowed by state and local authorities, by open burning. If burned, stay out of smoke. If the resealable plastic bag contacts the formulated product in any way, the bag must be triple-rinsed with clean water. Add the rinsate to the spray tank and dispose of the outer wrap as described above.

Notice to Buyer: Purchase of these materials does not confer any rights under patents of countries outside of the United States. Use of this quantity of purchased SYNCHRONY STS SP herbicide is permitted under claim 24 of U.S. Patent 5,084,082.

NOTICE OF WARRANTY

Du Pont warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for the purpose stated on such label only when used in accordance with the directions under normal use conditions. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Du Pont. In no case shall Du Pont be liable for consequential, special or indirect damages resulting from the use or handling of this product. All such risks are assumed by the buyer. DU PONT MAKES NO WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS

- 1 Registered trademark of BASF Corporation
- 2 Registered trademark of Valent USA Corporation
- 3 Registered trademark of American Cyanamid Company
- 4 Registered trademark of Zeneca, Inc.

SL - 145 9065 6/15/95