




DuPont™ Staple®

herbicide

DRAFT LABEL



“..... A Growing Partnership With Nature”

(The following Highlights section appears on product subset labeling for the U.S. excluding CA).

DUPONT™ STAPLE® HIGHLIGHTS

- STAPLE® provides control of a wide range of broadleaf weeds in cotton.
- STAPLE® can be used preemergence, postemergence (over the top), or as a post-directed spray.
- Preemergence applications must be applied by ground equipment.
- Post emergence applications may be applied by ground or aerial equipment (except Arizona).
- Post emergence applications may be made any time from emergence up to 60 days before harvest.
- Always include a surfactant for control of emerged weeds.
- STAPLE® may be tank mixed with most herbicides (except "Dual"), insecticides (except malathion), and PGR's.
- Consult label text for complete instructions. Always read and follow label instructions.

(The following Highlights section appears on product subset labeling for CA only)

DUPONT™ STAPLE® HIGHLIGHTS

- STAPLE® provides control of a wide range of broadleaf weeds in cotton.
- STAPLE® can be used postemergence (over the top), or as a post-directed spray.
- Post emergence applications may be applied by ground equipment.
- Post emergence applications may be made any time from emergence up to 60 days before harvest.
- Always include a surfactant for control of emerged weeds.
- STAPLE® may be tank mixed with most herbicides (except "Dual"), insecticides (except malathion), and PGR's.
- Consult label text for complete instructions. Always read and follow label instructions.

TABLE OF CONTENTS

Active Ingredient 1
 Precautionary Statements 1
U.S. except California 2
 General Information 2
 Directions for Use 2
 Agricultural Use Requirements 2
APPLICATION INFORMATION 3
 Spray - Volumes 3
 Sequential Applications 3
 Preemergence Use 4
 Pre/Post Programs 4
 Specific Weed Problems 4
 Postemergence Use 5
 Weeds Controlled 5
 Weeds Suppressed 6
 STAPLE® Plus DuPont™ ASSURE® II 6
 STAPLE® Plus MSMA or DSMA 6
 STAPLE® Plus "Buctril" 6
 STAPLE® Plus "Ignite" 6
 STAPLE® Plus Glyphosate 7
 Arizona Only 7
 Salvage Treatments 7
 Reduced Rate Sequential Applications 7
 Rotational Crop Restrictions 8
 Replanting to Cotton 8
 Cotton Crop Failure 8
 Additional Use Precautions 9
 Sprayer Preparation 9
 Sprayer Clean Up 9
 Spray Drift Management 9
 Storage and Disposal 10
 Notice to Buyer 10
CALIFORNIA 11
 General Information 11
 Directions for Use 11
 Agricultural Use Requirements 11
 Application Information 12
 Postemergence Band Use 12
 Spray Volumes 12
 Sequential Applications 12
 Soluble Packet Handling 12
 Weeds Controlled 13
 Weeds Suppressed 13
 Tank Mixtures 13
 STAPLE® Plus MSMA 13
 STAPLE® Plus Post Grass Herbicides 13
 Rotational Crop Restrictions 13
ADDITIONAL USE INFORMATION 14
 Sprayer Preparation 14
 Sprayer Clean Up 14
 Spray Drift Management 14
 Storage and Disposal 15
 Notice to Buyer 15
 Limitation of Warranty and Liability 16



DuPont™

Staple®


herbicide

For Use on Cotton in the States of AL, AR, AZ, CA, FL, GA, KS, LA, MO, MS, NC, NM, OK, PR, SC, TN, TX, & VA.

Soluble Powder

| <i>Active Ingredient</i> | <i>By Weight</i> |
|--|------------------|
| Pyrithiobac sodium | |
| Sodium 2-chloro-6-[(4,6-dimethoxy pyrimidin-2-yl)thio]benzoate | 85% |
| <i>Inert Ingredients</i> | 15% |
| TOTAL | 100% |

EPA Reg. No. 352-576



KEEP OUT OF REACH OF CHILDREN

WARNING AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand this label, find someone to explain it to you in detail.)

FIRST AID

IF IN EYES: Hold open eye and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

IF SWALLOWED: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

FIRST AID (Cont'd)

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-441-3637 for medical emergencies involving this product.

PRECAUTIONARY STATEMENTS HAZARD TO HUMANS AND DOMESTIC ANIMALS

WARNING! Causes substantial but temporary eye injury. Do not get in eyes or on clothing. Harmful if swallowed. Harmful if absorbed through skin. Avoid contact with skin, eyes or clothing.

PERSONAL PROTECTIVE EQUIPMENT

Some materials that are chemical resistant to this product are listed below. If you want more options follow the instructions for Category A on an EPA chemical resistance category selection chart.

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants.
- Chemical resistant gloves made of any waterproof material such as polyethylene or polyvinylchloride.
- Shoes plus socks.
- Protective eyewear.

Discard clothing or other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

USERS SHOULD: Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

ENVIRONMENTAL HAZARDS

This product is highly toxic to nontarget plants adjacent to area of application. Do not apply this product or allow it to drift to areas where endangered or desired plant species exist.

Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning of equipment or disposing of equipment washwaters.

(U.S. except California)

GENERAL INFORMATION

DuPont™ STAPLE® herbicide is a water soluble powder formulation packaged in premeasured soluble packets that rapidly dissolve in water.

STAPLE® may be applied preemergence, postemergence or post-directed to cotton and weeds by ground application equipment. STAPLE® may also be applied postemergence to cotton and weeds by aerial equipment (except in Arizona).

If STAPLE® is used in a tank mixture with other herbicides, read and follow all use instructions, warnings and precautions on companion herbicide labels.

BIOLOGICAL INFORMATION

STAPLE® is absorbed by weed foliage following postemergence application. Thorough coverage of target weed species, including the weed terminals or growing points, is required to obtain best results. When using a banded spray application, the band spray area should be of sufficient width to ensure thorough coverage of target weeds.

Growth of susceptible weeds is rapidly inhibited. Growing points and leaves of susceptible weeds appear yellow in 5-10 days. Death of leaf tissue and growing points will follow in some species, while others remain green but stunted and non-competitive. Susceptible weeds are controlled in 14-28 days.

Do not apply STAPLE® on any crops other than cotton. Most crops other than cotton are sensitive to STAPLE®. All direct and indirect contact (such as drift) to crops other than cotton or land not scheduled to be planted to cotton in the current growing season should be avoided.

RESISTANCE

When herbicides that affect the same biological site of action are used repeatedly over several years to control the same weed species in the same field, naturally-occurring resistant biotypes may survive a correctly applied herbicide treatment, propagate, and become dominant in that field. Adequate control of these resistant weed biotypes cannot be expected. If weed control is unsatisfactory, it may be necessary to retreat the problem area using a product affecting a different site of action.

To better manage herbicide resistance through delaying the proliferation and possible dominance of herbicide resistant weed biotypes, it may be necessary to change cultural practices within and between crop seasons such as using a combination of tillage, retreatment, tank-mix partners and/or sequential herbicide applications that have a different site of action. Weed escapes that are allowed to go to seed will promote the spread of resistant biotypes.

It is advisable to keep accurate records of pesticides applied to individual fields to help obtain information on the spread and dispersal of resistant biotypes. Consult your agricultural dealer, consultant, applicator, and/or appropriate state agricultural extension service representative for specific alternative cultural practices or herbicide recommendations available in your area.

INTEGRATED PEST MANAGEMENT

This product may be used as part of an Integrated Pest Management (IPM) program that can include biological, cultural, and genetic practices aimed at preventing economic pest damage. IPM principles and practices include field scouting or other detection methods, correct target pest identification, population monitoring, and treating when target pest populations reach locally determined action thresholds. Consult your state cooperative extension service, professional consultants or other qualified authorities to determine appropriate action treatment threshold levels for treating specific pest/crop systems in your area.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency in your State responsible for pesticide regulation.

DuPont will not be responsible for losses or damages resulting from the use of this product in any manner not specifically recommended by DuPont.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls.
- Chemical Resistant Gloves made of any Waterproof material.
- Shoes plus socks.
- Protective eyewear.

APPLICATION INFORMATION

ENVIRONMENTAL CONDITIONS FOR OPTIMUM PERFORMANCE

Weather: Conditions which are conducive to healthy, actively growing weeds optimize DuPont™ STAPLE® postemergence weed control performance. Ideal conditions include warm soil temperatures (70 Deg. F or more) and adequate soil moisture before, during and immediately after application.

Rainfastness: Rainfall immediately after treatment may wash STAPLE® off the weed foliage and result in reduced weed control. A minimum of 4 hours is needed to allow STAPLE® to be absorbed by weed foliage.

SPRAY VOLUMES

Ground Application - Apply uniformly by ground with a properly calibrated low pressure (20-40 psi) boom or cultivator mounted sprayer equipped with flat fan nozzles. Use a minimum of 10 gal. water per acre. Under heavy weed pressure or dense crop foliage, increase minimum spray volume to 20-40 gal. per acre.

Aerial Application (except Arizona) - Use orifice discs, cores and nozzle types and arrangements that will provide for optimum spray distribution and maximum coverage at a minimum of 3 GPA. Do not apply during inversion conditions, when winds are gusty, or when other conditions will favor poor coverage and/or drift.

SEQUENTIAL APPLICATIONS

Annual broadleaf weeds may have more than one flush of emerging seedlings. Also, regrowth of treated annual weeds may occur due to application being made to weeds under stress from adverse growing conditions. To control weeds under these conditions, a sequential application of STAPLE® may be necessary.

If a respray of treated annual weeds is necessary, allow the weeds to begin to regrow prior to making a second application of STAPLE®.

When using STAPLE® in sequential treatment program, allow a minimum of 7 days between applications.

GENERAL USE PRECAUTIONS

- Do not exceed 1.0 oz/A preemergence.
- Do not exceed 1.8 oz/A in a single postemergence application.
- Do not exceed 2.4 oz/A per acre per year.
- In West Texas (broadly defined as West of Highway 83), do not apply more than 1.5 ounce total per acre per year. Where continuous cotton is grown, do not apply more than 2.4 ounce total per acre per year.
- Do not apply this product through any type of irrigation system.
- Do not apply to irrigated land where tail water will be used to irrigate crops other than cotton.
- Do not apply within 60 days of harvest.

SOLUBLE PACKET HANDLING PRECAUTIONS

Exposure to moisture or excessive handling of the soluble packets can cause them to break open.

Do not handle the soluble packets with wet hands or place them on wet surfaces.

Protect unused soluble packets by immediately resealing them in the original barrier bag.

The outer, barrier bag is NOT soluble in water. Do not place it in the spray tank.

TANK MIX PRECAUTIONS

This product can be mixed with pesticide products labeled for use on cotton in accordance with the most restrictive of label limitations. Read all label precautions for tank mix partners prior to use. Follow all manufacturers label recommendations for the companion product. If these recommendations conflict with this label, do not tank mix with STAPLE®.

Since formulations may be changed and new ones introduced, it is recommended that users premix a small quantity of a desired tank mix and observe for possible adverse changes (settling out, flocculation, etc.). Avoid mixtures of several materials and very concentrated spray mixtures.

Do not tank mix STAPLE® with "Dual" herbicide as a postemergence treatment as crop injury may result.

Do not tank mix STAPLE® with malathion-containing insecticides (such as "Cythion" RTU or "Cythion" ULV) as crop injury may result. To avoid crop injury, apply malathion-containing insecticides at least 24 hours before or after application of STAPLE®.

PIMA COTTON PRECAUTION

Foliar injury to Pima cotton varieties from postemergence applications of STAPLE® can be more severe than that occasionally observed on upland cotton varieties (see NOTE: under POSTEMERGENCE USE section of label). Any of the plant stress conditions mentioned in the POSTEMERGENCE USE Note paragraph may further increase the severity of the injury to Pima varieties. Consequently, DuPont is not responsible for any crop injury arising from the use of STAPLE® on Pima cotton varieties.

PREEMERGENCE USE

DuPont™ STAPLE® may be applied preemergence in cotton to aid in the control of many problem weeds.

STAPLE® is absorbed by weed roots following a preemergence application. Susceptible weeds may germinate and emerge, but growth is rapidly inhibited. Death of leaf tissue and growing points will follow in some species while others remain green but stunted and non-competitive.

Preemergence applications of STAPLE® require rainfall or sprinkler irrigation to activate the herbicide. Degree and duration of weed control depend on: rate used, weed spectrum, growing conditions at and following time of treatment, soil texture, organic matter, soil moisture at the time of treatment, and precipitation following treatment.

The amount of rainfall required to activate STAPLE® preemergence treatments depends on the amount of soil moisture available when rainfall is received. Several rainfalls of 0.25 inch or less are not as effective as one rainfall or supplemental irrigation of 0.5-1 inch for activation.

Note: Temporary leaf yellowing and/or stunting may occur following a preemergence treatment. Plant stresses from seedling diseases, cool soil temperatures (60° F or less), thrips injury or excessive soil moisture may increase the sensitivity of cotton to injury from preemergence treatments of STAPLE®.

PREEMERGENCE USE PRECAUTIONS

- Do not use on cotton planted in furrows.
- Do not use on soils with less than 0.5% organic matter (OM).
- Do not use on course soils such as sands or loamy sands.
- Do not apply more than one preemergence application per year.
- Do not apply STAPLE® preemergence by aerial application.

WEEDS CONTROLLED OR SUPPRESSED

STAPLE® may be applied preemergence in cotton (including glyphosate (“Roundup” Ready), glufosinate (“LibertyLink”), and bromoxynil (BXN) tolerant varieties) for the control of pigweed (redroot, smooth), prickly sida, spotted spurge, spurred anoda & velvetleaf, and suppression of annual morningglory (cypressvine, entireleaf, ivyleaf, pitted, purple, red/scarlet, sharppod/cotton, small flower, threelobe, wooly) lambsquarters, smartweed (ladysthumb, pennsylvania), jimsonweed, coffee senna, palmer pigweed, lanceleaf sage, barnyardgrass, broadleaf signalgrass, goosegrass, fall panicum, giant foxtail, seedling johnsongrass & large crabgrass.

APPLICATION RATES

STAPLE® may be applied preemergence at the rates of 0.6 - 1.0 oz product/A. Use the higher rate of STAPLE® for harder to control weeds and/or in fields where high weed infestation levels are known to occur (see Specific Weed Problems section).

All rates are broadcast. Use proportionately less for band application.

PREEMERGENCE COMBINATIONS

For improved control of weeds, such as, prairie sunflower, lanceleaf sage and annual morningglory (entireleaf, ivyleaf, pitted, red morningglory, sharppod), STAPLE® may be applied preemergence in combination with diuron (“Karmex” DF). See “TANK MIX PRECAUTIONS” on page 3 for additional information.

STAPLE® + “Karmex” DF, “Direx” 4L

Medium Soils (sandy loam, loam, silt loam, silt): Apply STAPLE® at 0.6 - 0.8 ounce per acre plus diuron at 1.0 pound a.i. per acre.

Fine Soils (sandy clay loam, clay loam, silty clay loam, sandy clay): Apply STAPLE® at 0.6 - 1.0 ounce per acre plus diuron at 1.25 a.i. pound per acre.

- Do not use on soils with less than 1% organic matter when tank mixing with diuron.
- Do not use soil applied organophosphate insecticides where diuron will be applied preemergence. Refer to the specific diuron labels for further application information and use restrictions.

Note: A second application of STAPLE® may be applied postemergence if needed for extended weed control. See the POSTEMERGENCE USE section of this label for further application information and use restrictions.

PREEMERGENCE/POSTEMERGENCE PROGRAMS

A program of STAPLE® at 0.6 - 1.0 oz /A plus fluometuron (“Cotoran”) applied preemergence followed by STAPLE® early postemergence is recommended for improved control of bristly starbur, coffee senna, common ragweed, Florida beggarweed, hemp sesbania, jimsonweed, ladysthumb smartweed, lambsquarter, annual morningglory (cypressvine, entireleaf, ivyleaf, pitted, purple, red/scarlet, sharppod/cotton, smallflower) Pennsylvania smartweed, pigweed (redroot, smooth, spiny), prickly sida, spotted spurge, spurred anoda, velvetleaf and for suppression of palmer pigweed.

Specific Weed Problems (Sicklepod, Wild Poinsettia, Yellow Nutsedge): For improved control of the above weeds and other labeled weeds that often occur in high populations and/or have multiple seasonal flushes, a program of STAPLE® at 0.8 - 1.0 oz /A plus fluometuron applied preemergence followed by a postemergence application of STAPLE® alone or in combination with “MSMA” or “DSMA” is recommended.

Refer to the **POSTEMERGENCE USE** section of this label for use rates, application timings and restrictions.

POSTEMERGENCE USE

Application should be made postemergence (over-the-top) or as a post-directed spray to cotton (begin at cotyledon stage) and actively growing weeds. The degree of control and duration of effect are dependent on sensitivity and size of target weed and environmental conditions at time of and following application.

Application should be made by ground or aerial equipment (except Arizona). In Arizona, apply DuPont™ STAPLE® by ground equipment only.

Postemergence applications should be made to young, actively growing weeds. Control may be reduced if application is made to weeds under stress due to severe environmental conditions such as drought, excessive soil moisture or cool soil or air temperatures (60° F or less).

Foliar absorption is the primary means of uptake from postemergence applications of STAPLE®. Therefore, select a spray volume, delivery system and uniform spray pattern that will insure thorough coverage of the target weed species (including the growing point) to obtain best results. Increase spray volume as weed density and size increases. Avoid overlapping, and shut off spray booms while starting, turning, slowing or stopping, or injury to the crop may result.

Note: STAPLE® may cause temporary leaf yellowing, bronzing and/or leaf crinkling when applied as a postemergence application. Plant stresses from seedling diseases, insects (thrips injury), blowing sand (sand blasting), hail injury, cool soil or air temperatures (60° F or less), extreme temperature variations and lack of or excessive soil moisture just prior to or soon after treatment may increase the sensitivity of cotton to injury from STAPLE® treatments. To reduce the potential for increased cotton sensitivity, allow cotton plants to recover from stress conditions (approximately 2 days) prior to postemergence applications of STAPLE®.

APPLICATION RATES

Apply STAPLE® at 1.2 to 1.8 oz product / A for control of the weeds listed in "Weeds Controlled" section. Use the higher rate for arid growing conditions or where weed infestations are severe.

All rates are broadcast. Use proportionately less for band applications.

Note: In New Mexico and W. Texas (broadly defined as West of Highway 83) on sand or loamy sand soil types with less than 1% OM, confine in-season applications of STAPLE® to a band of no more than one-third the row width. If replanting back to cotton is necessary, replant outside the original treated band.

SPRAY ADDITIVES

Add a nonionic surfactant cleared for application to growing crops, at the rate of 0.25% V/V with all postemergence applications.

For the states of AZ, KS, NM, OK and TX, add a nonionic surfactant cleared for application to growing crops, at the rate of 0.25-0.5% V/V or a crop oil concentrate cleared for application to growing crops, at the rate of 1-2% V/V with all postemergence applications. Under arid conditions, a crop oil concentrate is recommended.

TIMING

STAPLE® may be used as a postemergence (over-the-top) or post-directed application to young, actively growing weeds.

WEEDS CONTROLLED

| Common Name | Scientific Name | Height or Diameter (inches) |
|----------------------------|---------------------------------|-----------------------------|
| Citronmelon | <i>Citrullus lanatus</i> | 1-4 |
| Cocklebur, common† | <i>Xanthium strumarium</i> | 1-4 |
| Cocklebur, common | (AZ, KS, NM, OK, TX only) | 1-3 |
| Coffee senna | <i>Cassia occidentalis</i> | 1-4 |
| Cowpea | <i>Vigna sinensis</i> | 1-4 |
| Dayflower, common/asiatic | <i>Commelina communis</i> | 1-3 |
| Devils claw | <i>Proboscidea louisianica</i> | 1-2 |
| Dock, curly | <i>Rumex crispus</i> | 1-4 |
| Florida beggarweed | <i>Desmodium tortuosum</i> | 1-4 |
| Goosefoot, nettleleaf | <i>Chenopodium murale</i> | 1-2 |
| Groundcherry, wright | <i>Physalis wrightii</i> | 1-2 |
| Jimsonweed | <i>Datura stramonium</i> | 1-4 |
| Knotweed, silversheath | <i>Polygonum argyrocoleon</i> | 1-2 |
| Ladysthumb | <i>Polygonum persicaria</i> | 1-4 |
| Morningglory, | | |
| cypressvine | <i>Ipomoea quamoclit</i> | 1-4 |
| entireleaf | <i>Ipomoea hederacea</i> | 1-4 |
| ivyleaf | <i>Ipomoea hederacea</i> | 1-4 |
| pitted | <i>Ipomoea lacunosa</i> | 1-3 |
| purple | <i>Ipomoea turbinata</i> | 1-4 |
| red/scarlet | <i>Ipomoea coccinea</i> | 1-3 |
| sharppod/cotton (seedling) | <i>Ipomoea trichocarpa</i> | 1-3 |
| smallflower | <i>Jacquemontia tamnifolia</i> | 1-4 |
| three-lobe | <i>Ipomoea triloba</i> | 1-3 |
| woolly | <i>Ipomoea hirsutula</i> | 1-3 |
| Mustard, black | <i>Brassica nigrum</i> | 1-2 |
| Nightshade | | |
| black | <i>Solanum nigrum</i> | 1-2 |
| hairy | <i>Solanum sarrachoides</i> | 1-2 |
| Pigweed | | |
| redroot | <i>Amaranthus retroflexus</i> | 1-2 |
| smooth | <i>Amaranthus hybridus</i> | 1-2 |
| spiny | <i>Amaranthus spinosus</i> | 1-2 |
| tumble | <i>Amaranthus albus</i> | 1-2 |
| Redweed | <i>Melochia corchorifolia</i> | 1-4 |
| Rocket, London | <i>Sisymbrium irio</i> | 1-2 |
| Sage, lanceleaf | <i>Salvia reflexa</i> | 0.25-0.5 |
| Sesbania, hemp*** | <i>Sesbania exaltata</i> | 1-4 |
| Shepherd's-purse | <i>Capsella bursa-pastoris</i> | 1-2 |
| Sida, prickly | <i>Sida spinosa</i> | 0.25-1 |
| Smartweed, Pennsylvania | <i>Polygonum pennsylvanicum</i> | 1-4 |
| Smellmelon | <i>Cucumis melo</i> | 1-3 |
| Spiderflower, spiny | <i>Cleome spinosa</i> | 1-4 |
| Spurred anoda | <i>Anoda cristata</i> | 1-4 |
| Starbur, bristly | <i>Acanthospermum hispidum</i> | 1-2 |
| Sunflower | | |
| common | <i>Helianthus annuus</i> | 1-4 |
| prairie | <i>Helianthus petiolaris</i> | 1-3 |
| Thisle, Russian | <i>Salsola iberica</i> | 1-2 |
| Velvetleaf | <i>Abutilon theophrasti</i> | 1-4 |
| Waterhemp, common | <i>Amaranthus tamariscinus</i> | 1-4 |
| Watermelon (volunteer) | <i>Citrullus vulgaris</i> | 1-2 |
| Wild poinsettia | <i>Euphorbia heterophylla</i> | 1-2 |
| Wild radish | <i>Raphanus raphanistrum</i> | 1-2 |

WEEDS SUPPRESSED††

| Common Name | Scientific Name | Height or Diameter (inches) |
|------------------|----------------------------|-----------------------------|
| Pigweed, palmer† | <i>Amaranthus palmeri</i> | 1-2 |
| Puncturevine | <i>Tribulus terrestris</i> | 1-2 |
| Purple nutsedge | <i>Cyperus rotundus</i> | 2-4 |
| Purslane, common | <i>Portulaca oleracea</i> | 1-2 |
| Sicklepod | <i>Cassia obtusifolia</i> | 0.5-2 |
| Yellow nutsedge | <i>Cyperus esculentus</i> | 2-4 |

- † Naturally occurring biotypes of this weed resistant to DuPont™ STAPLE® are known to exist. STAPLE® will not control these biotypes. See Information in Resistance section.
- †† Weed suppression is a visual reduction in weed competition (reduced population and/or vigor) as compared to an untreated check. The degree of control will vary with the rate used, size of weeds, crop competition, and environmental conditions.
- * In AL, AZ, FL and GA only, Palmer Pigweed is controlled at the height of 1-2 inches.
- *** Effective control may require sequential applications of STAPLE® as cotyledon to one-leaf stage plants are more difficult to control.

POSTEMERGENCE COMBINATIONS

STAPLE® may be tank mixed with other suitable registered herbicides to control weeds in addition to those listed.

STAPLE® can also be mixed with other suitable registered PGR's and insecticides labeled for use on cotton.

See "TANK MIX PRECAUTIONS" on page 3 for additional information.

STAPLE® plus DuPont™ ASSURE® II: Johnsongrass

STAPLE® may be tank mixed with ASSURE® II for additional early postemergence control of johnsongrass in cotton. This tank mix will also control many other grass species. Refer to ASSURE® II label for rates and timing of application.

Tank mixes of STAPLE® with other post grass herbicides can result in antagonism and partial control of of rhizome johnsongrass or annual grasses. To avoid poor control of rhizome johnsongrass or annual grasses apply other post grass herbicides at least 3 days prior to the application of STAPLE®.

STAPLE® plus MSMA or DSMA

STAPLE® may be tank mixed with MSMA or DSMA and applied post-directed for improved control of certain broadleaf weeds and suppression of sedges. Refer to MSMA or DSMA label for information on weeds, weed sizes, application conditions and use restrictions(follow label guidelines that are most restrictive).

- Treatments of STAPLE® + MSMA should be made only as a post-directed application using two nozzles per row set to provide complete coverage of the weeds while avoiding application over the top or to the growing point of the cotton plant.
- The use of gauge wheels or shielded sprayer equipment is recommended to prevent application of STAPLE® + MSMA over the top of cotton.
- Always add MSMA or DSMA to spray tank after the water soluble bag of STAPLE® has dissolved. Adding MSMA or DSMA to the spray tank first may prevent or delay the

dissolution of the water soluble bag causing filters and or nozzles to plug.

- Certain weeds such as black and hairy nightshade, palmer amaranth, and wright groundcherry have shown antagonism (reduced weed control) from tank mixtures of STAPLE® plus MSMA.

Sicklepod and Yellow nutsedge: STAPLE® will provide partial control (growth suppression) of sicklepod and yellow nutsedge when applied alone at the sizes indicated. For best results, STAPLE® should be applied as a post-directed application in combination with MSMA at 2 2/3 pints / A (2 lbs ai/A at 6 lbs ai /gal). Applications of STAPLE® + MSMA to sicklepod larger than 2 inches or yellow nutsedge larger than 4 inches will only provide partial control (growth suppression).

STAPLE® plus "Buctril" 4 EC (BXN Cotton)

A tank mixture of STAPLE® at 0.8 ounces per acre plus "Buctril" at 1 pint per acre may be applied as a postemergence treatment in Bromoxynil Tolerant (BXN) Cotton for improved control of pigweed (redroot, smooth, spiny and tumble) and prickly sida. (redroot, smooth, spiny and tumble).

The addition of STAPLE® to a BXN program will provide residual control of prickly sida, pigweed (redroot, smooth), spotted spurge, velvetleaf and spurred anoda and suppression of annual morningglory (cypressvine, entireleaf, ivyleaf, pitted, purple, red/scarlett, sharppod/cotton, smallflower, threelobe, wooly), lambsquarters, smartweed (ladysthumb, Pennsylvania), jimsonweed, coffee senna, palmer pigweed, lanceleaf sage, barnyardgrass, broadleaf signalgrass, goosegrass, fall panicum, giant foxtail, seedling johnsongrass and large crabgrass. Rainfall (0.5-1 inch) following the postemergence application is required for residual control.

Refer to the "Buctril" 4EC label for further information and use restrictions. Follow the label guidelines that are the most restrictive.

STAPLE® plus "Ignite" ("LibertyLink" Cotton)

A tank mixture of STAPLE® at 0.6 - 0.9 ounces per acre plus "Ignite" (glufosinate) at 32-40 ounces per acre may be applied as a postemergence treatment in "LibertyLink" cotton. See "Ignite" label for specific postemergence recommendations for weeds, weed sizes and Ignite rates.

The addition of STAPLE® to recommended postemergence rates of "Ignite" in "LibertyLink" cotton will provide residual control of prickly sida, pigweed (redroot, smooth), spotted spurge, velvetleaf and spurred anoda as well as suppression of annual morningglory (cypressvine, entireleaf, ivyleaf, purple, red/scarlet, sharppod/cotton, smallflower, threelobe, wooly), lambsquarters, smartweed (ladysthumb, Pennsylvania), jimsonweed, coffee senna, palmer pigweed, lanceleaf sage, barnyardgrass, broadleaf signalgrass, goosegrass, fall panicum, giant foxtail, seedling johnsongrass, and large crabgrass. Rainfall (0.5 - 1 inch) following the postemergence application is required for residual control.

Tank mixtures of STAPLE® plus "Ignite" must be applied broadcast in a minimum of 15 gallons of water per acre by ground and 10 gallons of water per acre by air.

**DuPont™ STAPLE® plus Glyphosate
(Glyphosate Tolerant Cotton)**

A tank mixture of STAPLE® at 0.6 - 0.8 ounces plus glyphosate at 24 - 32 ounces per acre may be applied as an early postemergence treatment in glyphosate tolerant cotton for improved control of hemp sesbania, morningglory (entireleaf, ivyleaf, pitted, scarlet/red), cutleaf evening primrose, prickly sida and palmer pigweed.

The addition of STAPLE® to a glyphosate only program will provide residual control of prickly sida, pigweed (redroot, smooth), spotted spurge, velvetleaf and spurred anoda and suppression of annual morningglory (cypressvine, entireleaf, ivyleaf, pitted, purple, red/scarlett, sharppod/cotton, smallflower, threelobe, wooly), lambsquarters, smartweed (ladysthumb, Pennsylvania), jimsonweed, coffee senna, palmer pigweed, lanceleaf sage, barnyardgrass, broadleaf signalgrass, goosegrass, fall panicum, giant foxtail, seedling johnsongrass and large crabgrass. Rainfall (0.5-1 inch) following the postemergence application is required for residual control.

Glyphosate rates are based on 4 pound active ingredient per gallon formulation. For other glyphosate formulations, rates should be adjusted proportionally to the active ingredient content of the formulation.

STAPLE® plus glyphosate may be applied postemergence (over-the-top) to glyphosate tolerant cotton through the 4th true leaf stage of growth (when 5th true leaf is the size of a quarter or less).

STAPLE® plus glyphosate may be applied using precision post-directed or hooded sprayers to glyphosate tolerant cotton through layby. When making post-directed applications, be especially careful to minimize contact of the spray with cotton leaves. Any single application should not exceed 0.8 ounces of STAPLE® or 32 ounces of glyphosate. No more than two post-directed applications may be made from the fifth leaf stage through layby. All applications must be 10 days apart and cotton must have at least two nodes of incremental growth between applications.

In New Mexico, Texas and Oklahoma, the STAPLE® rate range is 0.6 - 1.5 ounces in a single application with glyphosate.

Under hard water conditions, always add an appropriate rate of either a spray grade ammonium sulfate (AMS) or a water conditioner (such as Helena's "Quest" or Loveland's "Choice") to the spray water prior to adding the glyphosate.

Refer to the glyphosate formulation label for further application information and use restrictions. Follow the label guidelines that are the most restrictive.

Note: No antagonism has been observed to annual grass species from this tank mixture.

**Reduced Rate Sequential Applications
(Glyphosate Tolerant Cotton)**

Two applications of STAPLE® herbicide at 0.4 - 0.6 ounces per acre plus glyphosate at 16 - 32 ounces per acre (4 pound active ingredient per gallon formulation) are required for the control of the weeds listed in the Weeds Controlled section of this label. Use the higher rates for adverse

growing conditions, heavy weed infestations, or if additional residual control is desired.

Applications should be made postemergence (over-the-top) to glyphosate tolerant cotton from the cotyledon stage until the four leaf (node) stage of cotton development (until the fifth leaf stage reaches the size of a quarter). After the four leaf stage of growth through layby, apply the STAPLE® plus glyphosate tank mix as a post-directed application. For best results, the initial application should be made while weeds are small and actively growing. The applications must be made at least 10 days apart.

Refer to the label of the glyphosate formulation being used for any adjuvant recommendations.

Do not exceed a total of 4 quarts of glyphosate (4 pound per gallon formulation) per acre per season of in-crop use.

ARIZONA ONLY

A tank mix of STAPLE® at 0.8 - 1.8 ounces per acre plus "Roundup" UltraMax 5L at 26 ounces per acre may be applied as a postemergence (over-the-top) treatment in glyphosate tolerant cotton for improved weed control of groundcherry, morningglory, pigweed, puncturevine, purslane and nutsedge. Apply uniformly by ground application in a minimum of 5 - 20 gallons of water per acre.

Salvage Treatments (Glyphosate Tolerant Cotton)

This treatment may be used after the four leaf stage of development, where weeds threaten to cause the loss of the crop. Apply STAPLE® at 0.8 to 1.8 ounces per acre plus "Roundup" Ultra Max 5L at 26 ounces per acre (5 lb active ingredient per gallon) ounces per acre may be applied. Apply this treatment either as an over-the-top application or as a post-directed application sprayed higher on the cotton plants and over the weeds. If at the timing of the salvage treatment the weeds are larger than specified in this label, only partial control may be achieved.

Note: Salvage treatments will result in significant boll loss, delayed maturity and/or yield loss. No more than one salvage treatment should be used per growing season.

The combined total of in-crop over-the-top plus selective equipment applications must not exceed 4 pounds active ingredient of glyphosate per acre per crop.

10/18

ROTATIONAL CROP RESTRICTIONS

These crops may be planted after treatment with DuPont™ STAPLE®:

| CROP† | INTERVAL (MONTHS) |
|---|-------------------|
| Cotton* | Anytime |
| Winter/spring, wheat | 4 |
| Peanuts | 10 |
| Rice | 9 |
| Soybeans | 10 |
| Corn, field # | 9 |
| Corn, field ## | 10 |
| Corn, field IR (imidazilnone resistant) | 9 |
| Sorghum, grain | ‡ |
| Tobacco (transplant) | 10 |
| All other crops** | Field Bioassay |
| Arizona only - (all crops listed in the main table above plus) | |
| Field corn, grain sorghum | 10 |
| Watermelon, cantaloupe | 10 |

Note: When rotating to either cantaloupe or watermelon in the spring season following cotton, use only a single application of STAPLE® at no more than 1.8 ounces per acre.

Note: Where "drip irrigated" cotton is grown, rotate only to cotton.

Southeast US Only - (GA, NC, N.FL, SC, S.AL) (all crops listed in the main table above plus)

| | |
|------------------|----|
| Cabbage | 12 |
| Cantaloupe | 12 |
| Carrots | 12 |
| Collards | 12 |
| English Pea | 12 |
| Mustard (greens) | 12 |
| Onions | ++ |
| Peppers | 12 |
| Snap bean | 12 |
| Squash | 12 |
| Sweet Corn | 12 |
| Sweet Potato | 12 |
| Tomato | 12 |
| Turnips | 12 |
| Watermelon | 12 |

++ Do not rotate to Onions in the fall or spring crop season following a STAPLE® application.

* If initial seeding fails to produce a stand, cotton may be replanted into the treated area. Wherever possible, avoid disturbing original bed. If necessary to rework soil before replanting, use shallow cultivation. Do not rebed nor move soil into the original drill area.

Note: New Mexico and W. Texas (broadly defined as west of highway 83) - On sand or loamy sand soils with less than 1% OM replant cotton outside the original treated band.

** A minimum rotational interval of 10 months is required for all crops not listed above. Field bioassay results may require that this interval be extended. A successful field bioassay means growing to maturity a test strip of the crop(s) intended for production the following year. The test strip should cross the entire field including knolls and low areas.

† In AZ, KS, NM, OK, and TX the rotational crops listed may be planted at the indicated intervals provided the fields are deep plowed prior to planting the rotational crop.

Field corn, corn grown for grain or silage, may be planted at the indicated interval provided STAPLE® is applied on a band (not to exceed 50% of the row width) and the fields have had a thorough soil mixing, for example, two diskings or a deep plowing, prior to planting. Otherwise, do not rotate to field corn in the season following a STAPLE® application.

Note: New Mexico and W. Texas (broadly defined as West of Highway 83) do not rotate to field corn the season following a STAPLE® application.

Limited Geography--Field corn grown for grain or silage, only in the States of AL, AR, FL, GA, LA, MO, MS, NC, SC, TN, and VA, may be planted at the indicated interval provided all the STAPLE® applications made in cotton do not exceed a total of 1.8 ounces broadcast per acre per season. No additional soil mixing (disking or plowing) will be required beyond that which is normally done with the various production systems, e.g. conventional tillage, minimum till, no-till, ridge till, etc.

‡ Do not rotate to grain sorghum in the season following a STAPLE® application.

For Southeast Texas, in an area broadly defined as east of route I-35 and south of route US 90, to include Uvalde, Medina and Bexar counties, grain sorghum may be planted after a 10 month interval provided that in the above outlined area has received a minimum of 25 inches of rainfall following a STAPLE® application and the fields have had a thorough soil mixing, for example two diskings or a deep plowing prior to planting.

For the Rio Grande Valley of Texas, do not rotate to corn or grain sorghum in the fall crop season following a STAPLE® application.

REPLANTING TO COTTON

If initial seeding fails to produce a stand, cotton may be replanted in soil treated preemergence with STAPLE®.

Whenever possible avoid disturbing the original seedbed. If it proves necessary to rework the soil before replanting, use shallow cultivation. Do not relist nor move soil into the original drill area. Plant cotton seed at least 1 inch deep. Do not retreat field with a second preemergence application of STAPLE® during the same year as injury may result. For tank mix applications, see the respective combination product label for further replanting information. Follow the label guidelines that are the most restrictive. **Note:** In New Mexico and W. Texas (broadly defined as West Highway 83) on sand or loamy sand soil types with less than 1% OM, replant outside the original treated band.

COTTON CROP FAILURE

In the event of a cotton crop failure where seasonal constraints do not allow replanting to cotton, pyriithiobac sodium tolerant soybeans, such as DuPont™ STS® soybeans may be used as a replant crop. Pyriithiobac sodium tolerant soybeans may be planted 30 days following the last STAPLE® application to the failed cotton crop.

Where other herbicides have been used with or in conjunction with STAPLE®, refer to the other herbicide label(s) for any information or restrictions prior to replanting with STS® soybeans.

ADDITIONAL USE PRECAUTIONS

Injury to or loss of desirable trees or vegetation may result from failure to observe the following:

- Do not apply, drain or flush equipment on or near desirable trees or other plants or on areas where their roots may extend, or in locations where the chemical may be washed or moved into contact with their roots.
- Do not use on lawns, walks, driveways, tennis courts, or similar areas. Prevent drift of spray to desirable plants.

Injury to or loss of adjacent sensitive crops and vegetation may result from failure to observe the following:

- Avoid all direct or indirect (such as spray drift) contact with crops other than cotton.
- Carefully observe all sprayer cleanup instructions both prior to and after using this product, as spray tank residue may damage crops other than cotton.

SPRAYER PREPARATION

It is important that spray equipment is clean and free of existing pesticide deposits before using DuPont™ STAPLE®. Follow the clean up procedures specified on the label of the product(s) previously used. If no clean up procedure is provided, follow this clean up procedure for all application equipment before using STAPLE®:

1. Thoroughly rinse sprayer, tanks, boom and hoses with clean water.
2. Partially fill tank with water and add ammonia (1 gal. of ammonia per 100 gal. of tank volume) or a tank cleaner. Complete filling the tank and flush the cleaning solution through the boom hoses. Let solution stand for 15 minutes while agitating/recirculating, and then drain the tank by flushing the hoses, booms and nozzles.
3. Thoroughly rinse the sprayer, tanks, boom and hoses with clean water.
4. Follow label directions on product(s) previously sprayed for disposal.

Mix the proper amount of STAPLE® into the necessary volume of water in the spray tank with the agitator running. Continuous agitation is required for a uniform suspension and application. STAPLE® must be added first to the spray tank followed by tank mix partner, if used, then the adjuvant.

Use spray preparation of STAPLE® and approved adjuvant within 7 days or product degradation may occur. If spray preparation is left standing without agitation, thoroughly agitate before using.

PRECAUTION: Do not use chlorine bleach with ammonia. See Sprayer Clean Up Section for more information.

SPRAYER CLEAN UP

Spray equipment must be clean and free of previous pesticide deposits before applying STAPLE® and properly cleaned out after applying STAPLE®. Using the clean up procedures specified on the label of the previously used product, clean all application equipment before applying STAPLE®. If no clean up procedure is provided, use the procedure that

follows. Immediately following applications of STAPLE® thoroughly clean all mixing and spray equipment according to the following instructions:

1. Drain Tank: Thoroughly hose down the interior surfaces of the tank; then flush tank, boom and hoses with clean water for a minimum of 5 minutes. Loosen and physically remove any visible deposits.
2. Fill the tank with clean water and add one gal. of household ammonia* (3% active) for every 100 gal. of water. Flush the cleaning solution through the boom, hoses and nozzles. Add more water to completely fill the tank and allow to agitate/recirculate for at least 15 minutes. Again, flush the boom, hoses and nozzles with the cleaning solution, then drain the tank.
3. Remove the nozzles and screens and clean separately in a bucket containing the cleaning agent and water.
4. Repeat step 2.
5. Thoroughly rinse the tank with clean water for a minimum of 5 minutes, flushing the water through the hoses and boom.
6. Dispose of the rinsate on site or at an approved waste disposal facility.

* Equivalent amounts of an alternate-strength ammonia solution or DuPont approved cleaner can be used in the cleanout procedure. Carefully read and follow the individual cleaner instruction.

PRECAUTION

Do not use chlorine bleach with ammonia when cleaning out spray tanks. All traces of liquid fertilizer containing ammonia, ammonia nitrate or ammonium sulphate must be rinsed with water from the mixing and application equipment before adding any chlorine bleach solution. Failure to do so will release a gas with a musty chlorine odor which can cause eye, nose, throat and lung irritation.

Do not clean equipment in an enclosed area.

SPRAY DRIFT MANAGEMENT

The interaction of many equipment and weather-related factors determines the potential for spray drift. The applicator is responsible for considering all these factors when making application decisions.

AVOIDING SPRAY DRIFT IS THE RESPONSIBILITY OF THE APPLICATOR.

IMPORTANCE OF DROPLET SIZE

The most effective way to reduce drift potential is to apply large droplets (>150 - 200 microns). The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. The presence of sensitive species nearby, the environmental conditions, and pest pressure may affect how an applicator balances drift control and coverage. **APPLYING LARGER DROPLETS REDUCES DRIFT POTENTIAL, BUT WILL NOT PREVENT DRIFT IF APPLICATIONS ARE MADE IMPROPERLY OR UNDER UNFAVORABLE ENVIRONMENTAL CONDITIONS!** See **Wind, Temperature and**

12/18

Humidity, and Temperature Inversions sections of this label.

Controlling Droplet Size - General Techniques

- **Volume** - Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets.
- **Pressure** - Use the lower spray pressures recommended for the nozzle. Higher pressure reduces droplet size and does not improve canopy penetration. **WHEN HIGHER FLOW RATES ARE NEEDED, USE A HIGHER-CAPACITY NOZZLE INSTEAD OF INCREASING PRESSURE.**
- **Nozzle Type** - Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles.

Controlling Droplet Size - Aircraft

- **Number of Nozzles** - Use the minimum number of nozzles with the highest flow rate that provide uniform coverage.
- **Nozzle Orientation** - Orienting nozzles so that the spray is emitted backwards, parallel to the airstream will produce larger droplets than other orientations.
- **Nozzle Type** - Solid stream nozzles (such as disc and core with swirl plate removed) oriented straight back produce larger droplets than other nozzle types.
- **Boom Length** - The boom length should not exceed 3/4 of the wing or rotor length - longer booms increase drift potential.
- **Application Height** - Application more than 10 ft above the canopy increases the potential for spray drift.

BOOM HEIGHT

Setting the boom at the lowest labeled height (if specified) which provides uniform coverage reduces the exposure of droplets to evaporation and wind. For ground equipment, the boom should remain level with the crop and have minimal bounce.

WIND

Drift potential increases at wind speeds of less than 3 mph (due to inversion potential) or more than 10 mph. However, many factors, including droplet size and equipment type determine drift potential at any given wind speed. **AVOID GUSTY OR WINDLESS CONDITIONS.**

Note: Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they affect spray drift.

TEMPERATURE AND HUMIDITY

When making applications in hot and dry conditions, set up equipment to produce larger droplets to reduce effects of evaporation.

TEMPERATURE INVERSIONS

Drift potential is high during a temperature inversion. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain close to the ground and move laterally in a concentrated cloud. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be

indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

SHIELDED SPRAYERS

Shielding the boom or individual nozzles can reduce the effects of wind. However, it is the responsibility of the applicator to verify that the shields are preventing drift and not interfering with uniform deposition of the product.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

Pesticide Storage: Store product in original container only.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: Do not reuse the outer box or the resealable plastic bag. When all water-soluble packets are used, the outer packaging should be clean and may be disposed of in a sanitary landfill or by incineration, or if allowed by state and local authorities, by open burning. If burned, stay out of smoke. If the resealable plastic bag contacts the formulated product in any way, the bag must be triple-rinsed with clean water. Add the rinsate to the spray tank and dispose of the outer wrap as described above.

NOTICE TO BUYER: Purchase of this material does not confer any rights under patents of countries outside of the United States.

The DuPont Oval Logo, DuPont™ and STAPLE®. ASSURE® and STS® are trademarks or registered trademarks of E. I. duPont de Nemours and Company

“Cotoran”, “Cotton-Pro”, “Direx” 4L, are registered trademarks of Griffin LLC

“Karmex” is a registered trademark of E. I. du Pont de Nemours and Company and is exclusively licensed to Griffin LLC

“Buctril” 4EC is a registered trademark of Aventis CropScience

“Cythion” is a registered trademark of Helena Chemical Co.

“Dual” is a registered trademark of Syngenta Crop Protection Inc.

“Choice” is a registered trademark of UAP - Loveland Industries

“Quest” is a registered trademark of Helena Chemical Company

“Roundup” is a registered trademark of Monsanto Company

NET WEIGHT 6 oz per soluble bag.

(CALIFORNIA)

GENERAL INFORMATION

DuPont™ STAPLE® herbicide is a water soluble powder formulation packaged in premeasured soluble packets that rapidly dissolve in water.

STAPLE® may be applied postemergence (over-the-top) or post-directed to cotton and weeds by ground application equipment.

If STAPLE® is used in a tank mixture with other herbicides, read and follow all use instructions, warnings and precautions on companion herbicide labels.

BIOLOGICAL INFORMATION

STAPLE® is absorbed by weed foliage following postemergence application. Thorough coverage of target weed species, including the weed terminals or growing points, is required to obtain best results.

Growth of susceptible weeds is rapidly inhibited. Growing points and leaves of susceptible weeds appear yellow in 5-10 days. Death of leaf tissue and growing points will follow in some species, while others remain green but stunted and non-competitive. Susceptible weeds are controlled in 14-28 days.

Do not apply STAPLE® on any crops other than cotton. Most crops other than cotton are sensitive to STAPLE®. All direct and indirect contact (such as drift) to crops other than cotton or land not scheduled to be planted to cotton in the current growing season should be avoided.

RESISTANCE

When herbicides that affect the same biological site of action are used repeatedly over several years to control the same weed species in the same field, naturally-occurring resistant biotypes may survive a correctly applied herbicide treatment, propagate, and become dominant in that field. Adequate control of these resistant weed biotypes cannot be expected. If weed control is unsatisfactory, it may be necessary to retreat the problem area using a product affecting a different site of action.

To better manage herbicide resistance through delaying the proliferation and possible dominance of herbicide resistant weed biotypes, it may be necessary to change cultural practices within and between crop seasons such as using a combination of tillage, retreatment, tank-mix partners and/or sequential herbicide applications that have a different site of action. Weed escapes that are allowed to go to seed will promote the spread of resistant biotypes.

It is advisable to keep accurate records of pesticides applied to individual fields to help obtain information on the spread and dispersal of resistant biotypes. Consult your agricultural dealer, consultant, applicator, and/or appropriate state agricultural extension service representative for specific alternative cultural practices or herbicide recommendations available in your area.

INTEGRATED PEST MANAGEMENT

This product may be used as part of an Integrated Pest Management (IPM) program that can include biological, cultural, and genetic practices aimed at preventing economic pest damage. IPM principles and practices include field scouting or other detection methods, correct target pest identification, population monitoring, and treating when target pest populations reach locally determined action thresholds. Consult your state cooperative extension service, professional consultants or other qualified authorities to determine appropriate action treatment threshold levels for treating specific pest/crop systems in your area.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency in your State responsible for pesticide regulation.

DuPont will not be responsible for losses or damages resulting from the use of this product in any manner not specifically recommended by DuPont.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls.
- Chemical Resistant Gloves made of any Waterproof material.
- Shoes plus socks.
- Protective eyewear.

14/18

APPLICATION INFORMATION

ENVIRONMENTAL CONDITIONS FOR OPTIMUM PERFORMANCE

Weather: Conditions which are conducive to healthy, actively growing weeds optimize DuPont™ STAPLE® postemergence weed control performance. Ideal conditions include warm soil temperatures (70 Deg. F or more) and adequate soil moisture before, during and immediately after application.

Rainfastness: Rainfall immediately after treatment may wash STAPLE® off the weed foliage and result in reduced weed control. A minimum of 4 hours is needed to allow STAPLE® to be absorbed by weed foliage.

SPRAY VOLUMES

Ground Application - Apply uniformly by ground with a properly calibrated low pressure (20-40 psi) stabilized boom equipped with either Even flat fan, "Twinjet", or under leaf banding nozzles. Use 10-40 GPA with ground spray equipment.

SEQUENTIAL APPLICATIONS

Annual broadleaf weeds may have more than one flush of emerging seedlings. Also, regrowth of treated annual weeds may occur due to application being made to weeds under stress from adverse growing conditions. To control weeds under these conditions, a sequential application of STAPLE® may be necessary.

If a respray of treated annual weeds is necessary, allow the weeds to begin to regrow prior to making a second application of STAPLE®.

When using STAPLE® in sequential treatment program, allow a minimum of 7 days between applications.

GENERAL USE PRECAUTIONS

- Do not exceed 1.8 oz/A in a single postemergence application.
- Do not exceed 2.4 oz/A per acre per year.
- Do not apply this product through any type of irrigation system.
- Do not apply to irrigated land where tail water will be used to irrigate crops other than cotton.
- Do not apply within 60 days of harvest.

SOLUBLE PACKET

HANDLING PRECAUTIONS

Exposure to moisture or excessive handling of the soluble packets can cause them to break open.

Do not handle the soluble packets with wet hands or place them on wet surfaces.

Protect unused soluble packets by immediately resealing them in the original barrier bag.

The outer, barrier bag is NOT soluble in water. Do not place it in the spray tank.

TANK MIX PRECAUTIONS

This product can be mixed with pesticide products labeled for use on cotton in accordance with the most restrictive of label limitations. Read all label precautions for tank mix partners

prior to use. Follow all manufacturers label recommendations for the companion product. If these recommendations conflict with this label, do not tank mix with STAPLE®.

Since formulations may be changed and new ones introduced, it is recommended that users premix a small quantity of a desired tank mix and observe for possible adverse changes (settling out, flocculation, etc.). Avoid mixtures of several materials and very concentrated spray mixtures.

Do not tank mix STAPLE® with "Dual" herbicide as a postemergence treatment as crop injury may result.

Do not tank mix STAPLE® with malathion-containing insecticides (such as "Cythion" RTU or "Cythion" ULV) as crop injury may result. To avoid crop injury, apply malathion-containing insecticides at least 24 hours before or after application of STAPLE®.

PIMA COTTON PRECAUTIONS

Foliar injury to Pima cotton varieties from postemergence applications of STAPLE® can be more severe than that occasionally observed on upland cotton varieties (see NOTE: under POSTEMERGENCE BAND USE section of label). Any of the plant stress conditions mentioned in the POSTEMERGENCE BAND USE Note paragraph may further increase the severity of the injury to Pima varieties.

Consequently, DuPont is not responsible for any crop injury arising from the use of STAPLE® on Pima cotton varieties.

POSTEMERGENCE BAND USE

Application should be made by ground equipment only. Apply STAPLE® as a postemergence band (over-the-top) or as a post-directed band spray over the cotton seed row at 8 - 10 inches wide (not to exceed 10 inches in width). Applications may be made over the top when cotton is at the first visible true leaf stage through 6 inches in height, or post-directed when cotton is up to 10 inches in height. Weeds should be young and actively growing. The degree of control and duration of effect are dependent on the sensitivity and size of the target weed, coverage, rate of STAPLE® applied and the environmental conditions at the time of and following application. Regrowth of susceptible weeds may occur if these conditions are not met. Application should be made to the same number of rows as planted to avoid row width variations.

Postemergence applications should be made to young, actively growing weeds. Control may be reduced if application is made following a cultivation, i.e., dirt clods blocking the spray, dust covered weeds, weeds injured by cultivation equipment, or to weeds under stress due to severe environmental conditions such as drought, excessive soil moisture or cool soil or air temperatures (60° F or less).

Foliar absorption is the primary means of uptake from postemergence applications of STAPLE®. Therefore, select a spray volume, delivery system and uniform spray pattern that will insure thorough coverage of the target weed species (including the growing point) to obtain best results. Increase spray volume as weed density and size increases. Avoid overlapping, and shut off spray booms while starting, turning, slowing or stopping, or injury to the crop may result.

- Do not cultivate within 5 days after application.

- After a minimum of 5 days after application, a cultivation that moves soil to the crop and covers small treated weeds can improve efficacy.
- Do not sprinkler irrigate cotton within 48 hours after application.

Note: DuPont™ STAPLE® may cause temporary leaf yellowing, bronzing and/or leaf crinkling when applied as a postemergence application. Plant stresses from seedling diseases, insects, e.g., thrips injury, blowing sand (sand blasting), hail injury, cool soil or air temperatures (60° F or less), extreme temperature variations and lack of or excessive soil moisture just prior to or soon after treatment may increase the sensitivity of cotton to injury from STAPLE® treatments.

To reduce the potential for increased cotton sensitivity, allow cotton plants to recover from stress conditions prior to postemergence applications of STAPLE®.

APPLICATION RATES

Apply STAPLE® at 1.2 - 1.8 oz product per acre as a single application. Use the higher rates of STAPLE® to aid control under arid growing conditions or where weed infestations are severe.

All rates are broadcast. Use proportionately less for band applications.

SPRAY ADDITIVES

Add a nonionic surfactant, cleared for application to growing crops, at the rate of 0.25 - 0.5% V / V with all postemergence applications.

TIMING

STAPLE® may be used as a postemergence application to young, actively growing weeds.

WEEDS CONTROLLED

| Common Name | Scientific Name | Height or Diameter (inches) |
|------------------------|--------------------------------|-----------------------------|
| Cocklebur, common | <i>Xanthium strumarium</i> | 1-4 |
| Goosefoot, nettleleaf | <i>Chenopodium murale</i> | 1-2 |
| Nightshade | | |
| black | <i>Solanum nigrum</i> | 1-2 |
| hairy | <i>Solanum sarrachoides</i> | 1-2 |
| Knotweed, silversheath | <i>Polygonum argyrocoleon</i> | 1-2 |
| Mustard, black | <i>Brassica nigrum</i> | 1-2 |
| Pigweed | | |
| palmer | <i>Amaranthus palmeri</i> | 1-2 |
| redroot | <i>Amaranthus retroflexus</i> | 1-2 |
| smooth | <i>Amaranthus hybridus</i> | 1-2 |
| spiny | <i>Amaranthus spinosus</i> | 1-2 |
| Rocket, London | <i>Sisymbrium irio</i> | 1-2 |
| Shepherd's-purse | <i>Capsella bursa-pastoris</i> | 1-2 |
| Sunflower | | |
| common | <i>Helianthus annuus</i> | 1-4 |
| Velvetleaf | <i>Abutilon theophrasti</i> | 1-4 |
| Watermelon (volunteer) | <i>Citrullus vulgaris</i> | 1-2 |

WEEDS SUPPRESSED

| Common Name | Scientific Name | Height or Diameter (inches) |
|----------------------|----------------------------|-----------------------------|
| Groundcherry, wright | <i>Physalis wrightii</i> | 1-2 |
| Morningglory, * | | |
| entireleaf | <i>Ipomoea hederacea</i> | 1-4 |
| ivyleaf | <i>Ipomoea hederacea</i> | 1-4 |
| Puncturevine | <i>Tribulus terrestris</i> | 1-2 |
| Purslane, common | <i>Portulaca oleracea</i> | 1-3 |

* To aid control under arid growing conditions STAPLE® may be applied in a single application at up to 1.8 oz per acre. Include a nonionic surfactant at 0.5% V/V or crop oil concentrate at 1% V/V. For best activity, treat at the one to two leaf stage of weed growth.

TANK MIXTURES

STAPLE® plus MSMA

STAPLE® may be tank mixed with MSMA and applied post-directed for improved control of certain broadleaf weeds and suppression of sedges. Refer to MSMA label for information on weeds, weed sizes, application conditions and use restrictions (follow label guidelines that are most restrictive).

Only add MSMA to spray tank after the water soluble bag of STAPLE® has dissolved. Adding MSMA to the spray tank first may prevent or delay the dissolution of the water soluble bag causing filters and/or nozzles to plug.

Note: Certain weeds such as black and hairy nightshade, and wright groundcherry have shown antagonism (reduced weed control) from tank mixtures of STAPLE® plus MSMA.

STAPLE® plus Post Grass Herbicides

Tank mixes of STAPLE® with post grass herbicides can result in antagonism and partial control of grasses. To avoid poor grass control apply post grass herbicides at least 3 days prior to the application of STAPLE®.

ROTATIONAL CROP RESTRICTIONS

The rotational crops listed may be planted at the indicated intervals provided the fields are double disked or deep plowed prior to planting. These crops may be planted after treatment with STAPLE®:

| CROP | INTERVAL (MONTHS) |
|-------------------|-------------------|
| Cotton* | Anytime |
| Tomatoes | 8 |
| Wheat | 6 |
| All other crops** | Field Bioassay |

* If initial seeding fails to produce a stand, cotton may be replanted into the treated area. Wherever possible, avoid disturbing original bed. If necessary to rework soil before replanting, use shallow cultivation. Do not rebed nor move soil into the original drill area. **Note:** Where "drip irrigated" cotton is grown, rotate only to cotton.

** A minimum rotational interval of 10 months is required for all crops not listed above. Field bioassay results may require that this interval be extended. A successful field bioassay means growing to maturity a test strip of the crop(s) intended for production the following year. The test strip should cross the entire field including knolls and low areas.

Note: Shortening of the rotational intervals listed under the Rotational Crop Restrictions may result in crop injury.

16/18

ADDITIONAL USE INFORMATION

Injury to or loss of desirable trees or vegetation may result from failure to observe the following:

- Do not apply, drain or flush equipment on or near desirable trees or other plants or on areas where their roots may extend, or in locations where the chemical may be washed or moved into contact with their roots.
- Do not use on lawns, walks, driveways, tennis courts, or similar areas. Prevent drift of spray to desirable plants.

Injury to or loss of adjacent sensitive crops and vegetation may result from failure to observe the following:

- Avoid all direct or indirect (such as spray drift) contact with crops other than cotton.
- Carefully observe all sprayer cleanup instructions both prior to and after using this product, as spray tank residue may damage crops other than cotton.

SPRAYER PREPARATION

It is important that spray equipment is clean and free of existing pesticide deposits before using DuPont™ STAPLE®. Follow the clean up procedures specified on the label of the product(s) previously used. If no clean up procedure is provided, follow this clean up procedure for all application equipment before using STAPLE®:

1. Thoroughly rinse sprayer, tanks, boom and hoses with clean water.
2. Partially fill tank with water and add ammonia (1 gal. of ammonia per 100 gal. of tank volume) or a tank cleaner. Complete filling the tank and flush the cleaning solution through the boom hoses. Let solution stand for 15 minutes while agitating/recirculating, and then drain the tank by flushing the hoses, booms and nozzles.
3. Thoroughly rinse the sprayer, tanks, boom and hoses with clean water.
4. Follow label directions on product(s) previously sprayed for disposal.

Mix the proper amount of STAPLE® into the necessary volume of water in the spray tank with the agitator running. Continuous agitation is required for a uniform suspension and application. STAPLE® must be added first to the spray tank followed by tank mix partner, if used, then the adjuvant.

Use spray preparation of STAPLE® and approved adjuvant within 7 days or product degradation may occur. If spray preparation is left standing without agitation, thoroughly agitate before using.

PRECAUTION: Do not use chlorine bleach with ammonia. See Sprayer Clean Up Section for more information.

SPRAYER CLEAN UP

Spray equipment must be clean and free of previous pesticide deposits before applying STAPLE® and properly cleaned out after applying STAPLE®. Using the clean up procedures specified on the label of the previously used product, clean all application equipment before applying STAPLE®. If no clean up procedure is provided, use the procedure that

follows. Immediately following applications of STAPLE® thoroughly clean all mixing and spray equipment according to the following instructions:

1. Drain Tank: Thoroughly hose down the interior surfaces of the tank; then flush tank, boom and hoses with clean water for a minimum of 5 minutes. Loosen and physically remove any visible deposits.
2. Fill the tank with clean water and add one gal. of household ammonia* (3% active) for every 100 gal. of water. Flush the cleaning solution through the boom, hoses and nozzles. Add more water to completely fill the tank and allow to agitate/recirculate for at least 15 minutes. Again, flush the boom, hoses and nozzles with the cleaning solution, then drain the tank.
3. Remove the nozzles and screens and clean separately in a bucket containing the cleaning agent and water.
4. Repeat step 2.
5. Thoroughly rinse the tank with clean water for a minimum of 5 minutes, flushing the water through the hoses and boom.
6. Dispose of the rinsate on site or at an approved waste disposal facility.

* Equivalent amounts of an alternate-strength ammonia solution or DuPont approved cleaner can be used in the cleanout procedure. Carefully read and follow the individual cleaner instruction.

PRECAUTION

Do not use chlorine bleach with ammonia when cleaning out spray tanks. All traces of liquid fertilizer containing ammonia, ammonia nitrate or ammonium sulphate must be rinsed with water from the mixing and application equipment before adding any chlorine bleach solution. Failure to do so will release a gas with a musty chlorine odor which can cause eye, nose, throat and lung irritation.

Do not clean equipment in an enclosed area.

SPRAY DRIFT MANAGEMENT

The interaction of many equipment and weather-related factors determines the potential for spray drift. The applicator is responsible for considering all these factors when making application decisions.

AVOIDING SPRAY DRIFT IS THE RESPONSIBILITY OF THE APPLICATOR.

IMPORTANCE OF DROPLET SIZE

The most effective way to reduce drift potential is to apply large droplets (>150 - 200 microns). The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. The presence of sensitive species nearby, the environmental conditions, and pest pressure may affect how an applicator balances drift control and coverage. **APPLYING LARGER DROPLETS REDUCES DRIFT POTENTIAL, BUT WILL NOT PREVENT DRIFT IF APPLICATIONS ARE MADE IMPROPERLY OR UNDER UNFAVORABLE ENVIRONMENTAL CONDITIONS!** See **Wind, Temperature and**

Humidity, and Temperature Inversions sections of this label.

Controlling Droplet Size - General Techniques

- **Volume** - Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets.
- **Pressure** - Use the lower spray pressures recommended for the nozzle. Higher pressure reduces droplet size and does not improve canopy penetration. **WHEN HIGHER FLOW RATES ARE NEEDED, USE A HIGHER-CAPACITY NOZZLE INSTEAD OF INCREASING PRESSURE.**
- **Nozzle Type** - Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles.

BOOM HEIGHT

Setting the boom at the lowest labeled height (if specified) which provides uniform coverage reduces the exposure of droplets to evaporation and wind. For ground equipment, the boom should remain level with the crop and have minimal bounce.

WIND

Drift potential increases at wind speeds of less than 3 mph (due to inversion potential) or more than 10 mph. However, many factors, including droplet size and equipment type determine drift potential at any given wind speed. **AVOID GUSTY OR WINDLESS CONDITIONS.**

Note: Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they affect spray drift.

TEMPERATURE AND HUMIDITY

When making applications in hot and dry conditions, set up equipment to produce larger droplets to reduce effects of evaporation.

TEMPERATURE INVERSIONS

Drift potential is high during a temperature inversion. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain close to the ground and move laterally in a concentrated cloud. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

SHIELDED SPRAYERS

Shielding the boom or individual nozzles can reduce the effects of wind. However, it is the responsibility of the applicator to verify that the shields are preventing drift and not interfering with uniform deposition of the product.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

Pesticide Storage: Store product in original container only.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: Do not reuse the outer box or the resealable plastic bag. When all water-soluble packets are used, the outer packaging should be clean and may be disposed of in a sanitary landfill or by incineration, or if allowed by state and local authorities, by open burning. If burned, stay out of smoke. If the resealable plastic bag contacts the formulated product in any way, the bag must be triple-rinsed with clean water. Add the rinsate to the spray tank and dispose of the outer wrap as described above.

NOTICE TO BUYER: Purchase of this material does not confer any rights under patents of countries outside of the United States.

The DuPont Oval Logo, DuPont™, STAPLE®. and STS® are trademarks or registered trademarks of E. I. duPont de Nemours and Company

“Twinjet” is a registered trademark of Spraying Systems Co.
“Cythion” is a registered trademark of Helena Chemical Co.
“Dual” is a registered trademark of Syngenta Crop Protection Inc.

NET WEIGHT 6 oz per soluble bag.

D - 966[MASTER] 112904

18/18

LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read This Limitation of Warranty and Liability Before Buying or Using This Product. If the Terms Are Not Acceptable, Return the Product at Once, Unopened, and the Purchase Price Will Be Refunded.

It is impossible to eliminate all risks associated with the use of this product. Such risks arise from weather conditions, soil factors, off target movement, unconventional farming techniques, presence of other materials, the manner of use or application, or other unknown factors, all of which are beyond the control of DuPont. These risks can cause: ineffectiveness of the product; crop injury, or; injury to non-target crops or plants.

DuPont does not agree to be an insurer of these risks. WHEN YOU BUY OR USE THIS PRODUCT, YOU AGREE TO ACCEPT THESE RISKS.

DuPont warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for the purpose stated in the Directions for Use, subject to the inherent risks described above, when used in accordance with the Directions for Use under normal conditions.

DUPONT MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR OF MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

TO THE FULLEST EXTENT PERMITTED BY LAW, IN NO EVENT SHALL DUPONT OR SELLER BE LIABLE FOR ANY INCIDENTAL, CONSEQUENTIAL OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. BUYER'S OR USER'S BARGAINED-FOR EXPECTATION IS CROP PROTECTION. THE EXCLUSIVE REMEDY OF THE USER OR BUYER AND THE EXCLUSIVE LIABILITY OF DUPONT OR SELLER, FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY OR CONTRACT, NEGLIGENCE, TORT OR STRICT LIABILITY), WHETHER FROM FAILURE TO PERFORM OR INJURY TO CROPS OR OTHER PLANTS, AND RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT, OR AT THE ELECTION OF DUPONT OR SELLER, THE REPLACEMENT OF THE PRODUCT.

DuPont or its Ag Retailer must have prompt notice of any claim so that an immediate inspection of buyer's or user's growing crops can be made. Buyer and all users shall promptly notify DuPont or a DuPont Ag Retailer of any claims, whether based on contract, negligence, strict liability, other tort or otherwise or be barred from any remedy.

This Limitation of Warranty and Liability may not be amended by any oral or written agreement.

For product information call: 1-888-6-DUPONT

Internet address: <http://cropprotection.dupont.com/>

© 2000-2004 E. I. du Pont de Nemours and Company, Wilmington, Delaware 19898.

All rights reserved.