

Handwritten numbers: 1112, 250-577, 10/12

RESTRICTED USE PESTICIDE

This product is a restricted use herbicide due to reproductive and ground and surface water concerns. Users must read and follow all precautionary statements and instructions for use in order to minimize potential for Atrazine and Cyanazine to reach ground and surface water.

For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification.

ACCEPTED
NOV - 4 1993
DUPONT

EXTRAZINE® II DF

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 352-577

HERBICIDE DISPERSIBLE GRANULE

EXTRAZINE II DF is a water dispersible granule.

ACTIVE INGREDIENTS	BY WEIGHT
Cyanazine: 2-[[4-chloro-6-(ethylamino)-s-triazin- 2-yl]amino]-2-methylpropionitrile	67.5%
Atrazine: 2-chloro-4-(ethylamino)-6-(isopropylamino)-s-triazine	21.4%
Related compounds	1.1%
INERT INGREDIENTS	10.0%
	TOTAL 100.0%

This product contains 90 percent by weight active ingredients.
EPA Reg. No. 352-577

KEEP OUT OF REACH OF CHILDREN WARNING AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand this label, find someone to explain it to you in detail.)

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Call a physician or poison control center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person.

IF IN EYES: Flush with plenty of water. Get medical attention if irritation persists.

IF ON SKIN: Wash with plenty of soap and water.

IF INHALED: Remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth. Get medical attention.

For medical emergencies involving this product, call 1-800-441-3637.

PRECAUTIONARY STATEMENTS

HAZARD TO HUMANS AND DOMESTIC ANIMALS

WARNING: May be fatal if swallowed. Harmful if inhaled or absorbed through the skin. Causes temporary eye injury.

This product may be hazardous to your health. This product has been classified "Restricted Use" because, at doses which caused serious maternal illness in laboratory animals, birth defects were present. Use of protective clothing and equipment and following the precautions below can reduce risk.

Avoid breathing spray mist. Avoid contact with skin, eyes, or clothing. Do not get in eyes or on clothing.

Keep out of reach of domestic animals, particularly cattle. Consumption of this product, spray solutions, or water contaminated with product can result in serious illness or possible death of bovines.

PERSONAL PROTECTIVE EQUIPMENT

Applicators and other handlers must wear:

Long sleeved shirt and long pants.

Chemical-resistant gloves, such as barrier laminate or butyl rubber or nitrile rubber or polyvinyl chloride or viton or neoprene rubber.

Chemical-resistant footwear plus socks

Protective eyewear

Chemical-resistant apron when cleaning equipment, mixing or loading.

BEST AVAILABLE COPY

PRECAUTIONARY STATEMENTS (continued on next page)

2712

PRECAUTIONARY STATEMENTS (continued)

PERSONAL PROTECTIVE EQUIPMENT (continued)

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENGINEERING CONTROL STATEMENTS

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR part 170.240 (d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should: Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove personal protective equipment immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

Cyanazine and Atrazine, the active ingredients of EXTRAZINE II, are pesticides which can move (seep or travel) through soil and can contaminate groundwater which may be used as drinking water. Cyanazine and Atrazine have been found in groundwater as a result of agricultural use. Users are advised not to apply EXTRAZINE II where the water table (groundwater) is close to the surface and where the soils are very permeable (i.e. well drained soils such as loamy sands). Your local agricultural agencies can provide further information on the type of soil in your area and the location of groundwater.

Cyanazine has been detected in surface waters that receive run-off from treated areas. To minimize cyanazine run-off, follow the Best Management Practices outlined in the Directions For Use section of this label.

Atrazine is toxic to aquatic invertebrates. Do not apply directly to water or wetlands. Do not apply when weather conditions favor drift from treated areas. Runoff and drift from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water by cleaning of equipment or disposal of wastes.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls.
- Chemical-resistant gloves, such as barrier laminate or butyl rubber or nitrile rubber or polyvinyl chloride or viton or neoprene rubber.
- Shoes plus socks.

Best Management Practices for Ground and Surface Water Protection

This product may not be mixed or loaded within 50 feet of perennial or intermittent streams and rivers, natural or impounded lakes, and reservoirs. This product may not be mixed, loaded or used within 50 feet of all wells, including abandoned wells, drainage wells, and sinkholes.

Operations that involve mixing, loading, rinsing, or washing of this product into or from pesticide handling or application equipment or containers within 50 feet of any well are prohibited unless conducted on an impervious pad constructed to withstand the weight of the heaviest load that may be positioned on or moved across the pad. Such a pad shall be designed and maintained to contain any product spills or equipment leaks, container or equipment rinse or wash-water, and rain-water that may fall on the pad.

Surface water shall not be allowed to either flow over or from the pad, which means the pad must be self-contained. The pad shall be sloped to facilitate material removal. An unroofed pad shall be of sufficient capacity to contain, at a minimum, 110% of the capacity of the largest pesticide container or application equipment on the pad. A pad that is covered by a roof or sufficient size to completely exclude precipitation from contact with the pad shall have a minimum containment capacity of 100% of the capacity of the largest pesticide container or application equipment on the pad. Containment capacities as described above shall be maintained at all times. The above-specified minimum containment capacities do not apply to vehicles when delivering pesticide shipments to the mixing/loading site.

States may have in effect additional requirements regarding well-head setbacks and operational area containment.

This product may not be applied aerially or by ground within 66 feet of the points where field surface water run-off enters perennial or intermittent streams and rivers or within 200 feet of natural or impounded lakes and reservoirs. This product may only be applied to highly erodible land if the 66 foot buffer or set-back from run-off points is planted to crop or seeded with grass.

Atrazine Rate Limits: One pound of EXTRAZINE II DF contains 0.225 pounds active ingredient atrazine. For soil applications prior to crop emergence (i.e., early preplant, preplant incorporated, preplant surface, at planting or preemergence) the following applies:

- a. On highly erodible land, as defined by the Soil Conservation Service (SCS), if conservation tillage is utilized (> or = 30 % plant residue), the maximum rate of atrazine from all sources is 2 pounds a.i./A. If plant residue is <30%, the maximum rate of atrazine is 1.6 pounds a.i./A.
- b. On land that is not highly erodible, the maximum rate of atrazine is 2 pounds a.i./A.

For postemergence applications, if there has been no previous soil application to that crop, the maximum rate of atrazine from all sources is 2 pounds a.i./A. If there has been a previous soil application to that crop, do not exceed a total of 2.5 pounds atrazine a.i./A per calendar year.

Cyanazine Rate Limits: One pound of EXTRAZINE II DF contains 0.675 lb cyanazine active ingredient (a.i.). Adhere to the use rate recommendations in this or other label. In addition:

- a. Do not apply more than 6.5 lbs. total cyanazine a.i. (all sources) per acre per year to any land.
- b. On highly erodible land, as defined by the Soil Conservation Service, if plant residue cover is less than 30%, do not apply more than 3.0 pounds total cyanazine a.i. (all sources) per acre per year.

Where there are state/local requirements regarding atrazine or cyanazine use (including lower maximum rates and/or higher set-backs) which are different from the label, the more restrictive/protective requirements apply.

GENERAL INFORMATION

EXTRAZINE II DF Herbicide is a selective herbicide for the control of annual grasses and broadleaf weeds in field corn, popcorn, and sweet corn.

Consult your local Agricultural Extension Agent for help in determining soil type, organic matter content, and the most appropriate herbicide rate for local conditions.

EXTRAZINE II DF is not effective when used preemergence on peat or muck soils. Do not use EXTRAZINE II DF on sands or loamy sands (soils consisting of more than 70% sand) containing less than 1% organic matter.

Where surfactants or emulsible vegetable oils are added to EXTRAZINE II DF for over the top of corn postemergence applications, use on field corn only.

Do not apply this product through any type of irrigation system.

Do not apply this product with aerial application equipment.

Postemergence application to corn must be made before corn reaches 12 inches in height.

Application for quackgrass suppression in corn are restricted to a spring application only. No fall applications are permitted.

Weather Effects: As a preemergence herbicide, EXTRAZINE II DF is active mainly through the roots, and, therefore, its effect on weeds is dependent on adequate rainfall to move the herbicide into the root zone. Moisture should be sufficient to thoroughly wet the soil throughout the zone where weed seeds may germinate and enough to make the soil too wet to cultivate. Rotary hoeing or shallow cultivation is recommended for those applications which are not incorporated at the time of treatment, if adequate rainfall or sprinkler irrigation has not occurred within about ten days after application of EXTRAZINE II DF.

Heavy rainfall between planting and crop emergence may cause crop injury or stand loss. Rainfall tends to cause excessive concentrations of herbicide in the seed furrow, resulting in possible crop injury. Level deep planter marks or seed furrows before application.

Under conditions which delay weed germination, such as low temperatures, lack of soil surface moisture, or when germination extends over a long period, the effectiveness of the herbicide may be impaired. Rotary hoeing, shallow cultivation or a post-emergence herbicide treatment may be of benefit under these circumstances.

If the crop is cultivated, tillage should be shallow to minimize herbicide dilution in the soil. Should the crop stand be lost due to adverse weather conditions, insects, etc., the field can be replanted the same season to corn or sorghum.

BEST AVAILABLE COPY

10/12

To enhance weed control in areas of less than 25 inches of rainfall or where long dry periods are common, these treatments may require shallow incorporation with a tool such as a field cultivator operated from 5-7 mph. Incorporation should not be more than three inches deep to keep from burying the herbicide. A spike-toothed harrow, deep tillage disk or rolling basket device is not recommended for incorporating EXTRAZINE II.

When applied as a post-emergence herbicide, EXTRAZINE II DF is also active through foliage as well as through the roots. Yellowing and/or stunting of the crop may result from this treatment, particularly if cold, adverse growing conditions occur after application. Extended or extreme cold and wet conditions may reduce stands. Do not apply EXTRAZINE II postemergence to a crop that is damaged or growing under stress.

OBSERVE ALL CAUTIONS AND LIMITATIONS ON LABELING OF ALL PRODUCTS USED IN MIXTURES.

Rotational crops: (1) Plant only corn, sorghum, or soybeans the year following the use of this mixture. (2) If soybeans are to be planted, injury may occur. (3) If applied after June 10, do not rotate with crops other than corn or sorghum the next year or injury may occur. (4) In the high plains and inter-mountain areas of the West where rainfall is sparse and erratic or where irrigation is required, use only when corn or sorghum is to be planted the following year, or a crop of corn or sorghum not treated with this mixture or atrazine is to precede other rotational crops. (5) Small grains may be planted 15 months following treatment. (6) All other crops may be planted 18 months after application.

Triazine Resistant Weeds: In fields where triazine resistant biotypes of weeds have been identified, EXTRAZINE II should be used in combination with or in sequence with other registered non-triazine herbicides. (Triazine resistant biotypes of Kochia and Pigweed have been identified in some fields in the Western Great Plains and triazine resistant biotypes of Pigweed and Lambsquarters have been identified in some fields in various states.) Consult with appropriate state agricultural extension service representatives for specific recommendations.

APPLICATION DIRECTIONS

GENERAL MIXING AND SPRAYING DIRECTIONS

This product may not be mixed/loaded, or used within 50 feet of all wells including abandoned wells, drainage wells and sink holes.

This product may not be applied by means of chemigation or aerial application.

Use sufficient agitation to ensure that the EXTRAZINE II DF is completely dispersed and in uniform suspension prior to application or tank mixing with other formulations.

The following general mixing instructions are recommended:

I. General

1. Unless otherwise specified, use at least 7.5 gallons of water per acre for soil applications and at least 15 gallons of water per acre for foliar applications for all applications with ground equipment.

NOTE: Sufficient carrier must be used to assure uniform application. Follow label requirements of all products used in tank mix combinations.

2. A nitrogen solution or complete liquid fertilizer may replace all or part of the water as a carrier for preemergence or preplant application on corn. Do not apply fertilizer mixtures after crop emerges, as injury may occur.
 3. Always check the tank mix compatibility (TMC) of this or any other formulation before mixing with liquid fertilizer carriers or other formulations. A simple but generally reliable TMC evaluation procedure has been provided for your use in Step II of these mixing instructions.
 4. Start with thoroughly clean equipment. (See the labels of the previous compounds used for cleaning instructions.)
 5. Fill the tank at least 1/2 full with carrier. Start and maintain consistent agitation through all mixing and spraying procedures. Make sure the agitation system is working properly and creates a rippling or rolling action on the liquid surface.
 6. Slowly add the recommended amount of EXTRAZINE II DF to the tank or inductor.
 7. Fill the tank to 75 percent capacity with carrier. Filling and bypass lines should be kept below liquid surface. Increase tank agitation if necessary to maintain surface action.
 8. When desired, appropriate emulsible crop oil, crop oil concentrate, or other tank mix formulations should be added at this time. Pre-slurry these added ingredients before addition, if the compatibility test shows it to be necessary.
 9. Complete filling tank, maintaining sufficient agitation at all times to ensure surface action. This applies to both spray and nurse tanks.
 10. Tank mixtures should always be applied immediately after preparation. If, for any reason, this is not possible, assure that sufficient agitation has been provided to re-mix all products and check for complete re-suspension prior to application.
 11. Empty tank as completely as possible before refilling to prevent buildup of oil or emulsible concentrate residues when tank mixing with these formulations. Always maintain agitation to avoid separation.
 12. If an oil or emulsible concentrate film starts to build up after using these formulations, drain and clean the tank with strong detergent solution or appropriate solvent.
 13. It is recommended that the sprayer be thoroughly cleaned by flushing with a detergent solution at the end of each work day when any emulsible crop oil, crop oil concentrate, or other emulsible formulation has been used either alone or in tank mix combinations with other pesticide formulations. This precaution will ensure a clean sprayer and continued trouble-free operation.
- II. Tank Mix Compatibility Evaluation Procedure**
1. Add 1 pint of carrier liquid to each of 2 one-quart jars. Mark 1 quart jar "with" and the other "without."

BEST AVAILABLE COPY

25412

- Add 1/4 teaspoon of a suitable compatibility agent (1/4 teaspoon/pint = 2 pints/100 gallons of carrier) to the jar marked "with", cap the jar and shake gently for 5 to 10 seconds to mix.
- Add the appropriate amount of herbicide to both jars, cap each jar and shake gently for 5 to 10 seconds to mix. If problems are encountered in mixing wettable powder or dry flowable formulations into a liquid fertilizer, pre-slurry these formulations in water prior to their addition to the liquid fertilizer and proceed with the test. The following chart has been provided to assist in selecting the approximate EXTRAZINE II DF use rate for this evaluation.

JAR TEST FOR EXTRAZINE II DF COMPATIBILITY						
Gallons of Liquid Carrier per acre	4	7.5	15	20	25	30
Teaspoons of EXTRAZINE II DF per pint of Liquid Carrier	6.0	3.2	1.6	1.2	1.0	0.8

This chart is based on 1 pound of EXTRAZINE II DF (0.9 pounds active ingredient) per acre in the indicated carrier volumes. Intended field use rates are achieved by varying the amount of EXTRAZINE II DF. (e.g., for a field use rate of 3 pounds of EXTRAZINE II DF in 15 gallons of carrier per acre, add 4.8 teaspoons of EXTRAZINE II DF to the quart jars containing 1 pint of carrier. Calculation: 3 pounds of EXTRAZINE II DF/15 gallons of carrier per acre = 3 X 1.6 = 4.8 teaspoons of EXTRAZINE II DF per pint of carrier.)

- Let each jar stand one-half hour and make observations. If any separation, agglomeration, or precipitation has occurred, shake the jar again for 10 to 15 seconds, and note whether any of the following occur:
 - Separated phases do not re-mix uniformly.
 - Screen/nozzle plugging lumps do not disperse.
 - Precipitate does not re-suspend readily.
 - Precipitate sticks tenaciously to the glass.
- If none of the above problems occur in either jar, the herbicides can, in most cases, be safely used in that carrier without a compatibility agent.
- If problem 4.a or 4.b occurs in the jar marked "without" but does not occur in the jar marked "with", the compatibility agent should be used.
- If problem 4.a or 4.b is seen in both jars, then the herbicides and carrier are incompatible and should not be used in the same spray tank. Alternatively, a different tank mix compatibility agent can be evaluated.
- If problem 4.c or 4.d occurs in the jar marked "without" but does not occur in the jar marked "with", the compatibility agent should be used unless constant, thorough agitation can be maintained and immediate clean-out of the spray system is performed.
- If problem 4.c or 4.d is seen in the jar marked "with," the user proceeds with mixing and application at his own risk should the agitation in the system be insufficient or curtailed.

- Those mixtures defined as compatible in this test should then be mixed for use as indicated in Steps 1 through 12 of the general mixing instructions listed above.

If a test such as outlined indicates that components of a proposed mix are compatible, the applicator still has the responsibility of combining materials in sequence to the spray tank in accordance with directions prescribed on the label of the herbicides or pesticides involved.

Tank Mix Combination	Compatibility Agents
EXTRAZINE II DF /ASSO (Liquid Fertilizer Grade)	Probably not needed in 28-0-0, 10-34-0. Compex may help in others.
EXTRAZINE II DF /SUTAN+. 6.7E	Probably not needed in 28-0-0. Incompatible in 10-34-0. Unite, Spray-Mate, Kem-Link, may help in others.
EXTRAZINE II DF /DUAL 8E	Probably not needed in 28-0-0. Unite, Spray-Mate, Ivory Liquid may help in others.

Tests have indicated that compatibility agents, noted below by the various tank mix combinations, may give improved compatibility in liquid fertilizers.

III. Application Equipment

- Use application equipment fitted with nozzles that provide accurate and uniform coverage. Be certain that nozzles are uniformly spaced and the same size. Calibrate sprayer before use and re-check frequently during use whenever possible.
- Use a pump with capacity to:
 - Maintain 35 to 40 psi at the nozzles.
 - Provide sufficient agitation in tank to keep mixture in suspension.
 - Provide a minimum of 20 percent bypass at all times.
- Use centrifugal pumps which provide sufficient shear action for dispersing and mixing this product. The pump should provide a minimum of 10 gallons/minute/100-gallon tank size circulated through the jets of a correctly-positioned sparger tube.
- Use screens to protect the pump and to prevent nozzles from clogging. Screens placed on suction side of pump should be 10 to 16 mesh. Do not place a screen in the recirculation line. Use a 40- to 50-mesh screen between the pump and boom and, where required, 50-mesh screens at the nozzles. Check your equipment manufacturer's literature for specific recommendations.

FERTILIZER IMPREGNATION

EXTRAZINE II DF Herbicide may be applied when coated on or impregnated in dry granular fertilizer for early preplant, preemergence or preplant incorporated weed control in field corn. All recommendations, cautions and special precautions on this label must be followed along with state regulations relating to dry bulk fertilizer blending, impregnating and labeling.

GENERAL BLENDING DIRECTIONS

EXTRAZINE II DF may be coated on or impregnated in dry bulk fertilizers using tower blenders, rotary drum blenders or blending augurs or conveyors. DO NOT impregnate EXTRAZINE II or tank mixes containing EXTRAZINE II on or in fertilizers containing Ammonium Nitrate, Potassium Nitrate,

BEST AVAILABLE COPY

100712

or Sodium Nitrate. Do not use on straight limestone since absorption will not be achieved. Fertilizer blends containing limestone can be impregnated When using EXTRAZINE II alone. Use a minimum of 200 lbs. and a maximum of 450 lbs. per acre of dry fertilizer.

Use equipment that will give a uniform distribution of the herbicide throughout each batch of impregnated fertilizer. Non-uniform impregnation can cause crop injury or unsatisfactory performance.

A. EXTRAZINE II DF may be used as the only herbicide for impregnation.

1. Add EXTRAZINE II DF to 1/2 the total fertilizer volume required.
2. Spray one gallon of water (to break down DF) and one gallon of diesel fuel (to prevent evaporation and crusting) per ton of fertilizer and allow to mix thoroughly. NOTE: If the fertilizer is dusty, add the diesel fuel before adding the herbicide.
3. Add remaining fertilizer and mix thoroughly. (3 minutes or more for rotary drum blenders.)
4. Add 2-3% Ag-Sorb or 1-2% MP-79 drying agent (or a suitable amount of another effective drying agent) to insure a spreadable herbicide/fertilizer mixture. The need for a drying agent is determined by the wetness of the fertilizer batch. Wetness can change with humidity, nitrogen content, fertilizer types, fertilizer rates and herbicide rates.

B. EXTRAZINE II DF may be used in tank mixes with other dry herbicides including PRINCEP CALIBER 90. Follow the procedure as above in "A".

C. EXTRAZINE II DF may be used in tank mixes where an EC or other liquid herbicide acts as the sticking agent. This may eliminate the need for water and/or diesel fuel.

1. While fertilizer is blending, add the EXTRAZINE II DF. Experience has shown that this will provide the most consistent performance due to the grinding action of the fertilizer on the 90DF.
2. Spray in the EC herbicide and mix thoroughly. (3 minutes or more for rotary drum blenders).
3. Add drying agent to insure a spreadable herbicide/fertilizer mixture. Usually less drying agent is required when using EXTRAZINE II DF.

D. Pre-slurried EXTRAZINE II DF can be used alone or in a tank mix for impregnation. For rotary-drum mixers, the liquids can be moved into the drum using an air system or liquid pump. Do not add extra water. Add drying agent to insure a spreadable herbicide/fertilizer mixture.

(For more information on Drying Agents, Application Equipment, Calibration Guide and variations of these methods, see Du Pont Bulletin on "Fertilizer Impregnation").

CLEAN OUT: Equipment used to impregnate or apply fertilizer impregnated with EXTRAZINE II or combinations including EXTRAZINE II must be cleaned out by running at least 1,000 lbs. of fertilizer not impregnated with EXTRAZINE II through the impregnation equipment and application equipment, if the next batch of material is to be applied to a crop for which EXTRAZINE II or a combination herbicide is not registered.

APPLICATION: Uniform application of EXTRAZINE II DF which has been impregnated in or coated on dry fertilizer is essential for satisfactory weed control and crop safety. Accurate

calibration of the fertilizer applicator is necessary. Applying while turning at the ends of the fields may result in excessive application rates causing crop injury. Do not double apply across the ends or sides of the field.

Crop injury and/or poor weed control may occur where the impregnated fertilizer is not uniformly applied. Air flow or augur metered application equipment is preferred (one pass application). If other equipment is used, the recommended method of application is to apply 1/2 the recommended rate and overlap 50 percent to double apply by splitting the middles to obtain the best distribution pattern.

Apply immediately after impregnation. Impregnated fertilizer may become lumpy and difficult to spread if stored.

RATES AND TIMING: Use the application rates and timing shown in the appropriate sections of this label. Follow the precautions on the labels of all products used.

CORN

Weeds Controlled by EXTRAZINE II DF Alone and in Combination with Other Herbicides on Corn.

Grasses


Annual bluegrass	Giant foxtail
Annual fescues	Goosegrass
Annual (Italian) ryegrass	Green foxtail
Annual sedge	Junglerice
Barnyardgrass(1)	Stinkgrass (Indian
Bullgrass	lovegrass)
Crabgrass	Witchgrass
Fall panicum	Yellow foxtail

Broadleaves

Annual groundcherry	Pigweed(1)
Annual morningglory	Pineappleweed
Black mustard	Plantain
Buffalobur	Poorjoe
Buttercup (annual)	Prickly sida (teaweed)
Carpetweed	Prostrate knotweed
Common chickweed	Prostrate spurge
Cocklebur(2)	Ragweed (Common)
Common groundsel	Russian Thistle
Common mallow	Shepherdspurse
Common purslane	Smallflower galinsoga
Corn spurry	Smartweed
Curly dock (seedling)	(Pennsylvania)
Fiddleneck	Spiny sida
Florida pusley	Sunflower(2)
(Florida purslane)	(wild, annual, common)
Hedge mustard	Tarweed cuphea
Jimsonweed(1)	(Gumweed)
Kochia	Velvetleaf(1)
Ladysthumb	Wild buckwheat
Lambsquarters	Wild mustard
Mayweed	Wild radish
Nightshade (annual)	Wild turnip

(1) Under conditions such as low temperatures, lack of soil surface moisture, or other factors that may cause delay in germination of the seeds, the effectiveness of EXTRAZINE II may be impaired against these weeds.

(2) The degree of control will be reduced if soil moisture and temperature conditions cause deep germination of the seed.



7/10/12

**PREEMERGENCE
-PREPLANT INCORPORATED**

Apply EXTRAZINE II treatments just before, at, or after planting but before crop has emerged. Avoid removal of treated soil from seedrow prior to or during the planting operation.

EXTRAZINE II may also be applied early prior to planting or in a split application if pre-season weed control is desired. For split applications, do not exceed the total amount of EXTRAZINE II for the soil texture and organic matter shown in Table 1. If EXTRAZINE II is applied early, more than 15 days before planting, a split application of EXTRAZINE II or some other herbicide treatment may be necessary at or after planting to provide additional length of weed control. For further information, see recommendations in the Conservation Tillage section of this label.

Rotary hoeing is recommended for preemergence applications which do not receive adequate rainfall or sprinkler irrigation to wet the top 2 inches of soil or depth of germinating weeds within about 10 days after application.

EXTRAZINE II alone or in tank mix combinations should not be incorporated more than three inches deep to keep from burying the herbicide. Single or two pass incorporation with a tool such as a field cultivator operated at 5-7 mph is acceptable. A spike-toothed harrow, deep tillage disk or rolling basket device is not recommended for incorporating EXTRAZINE II.

See the Rotational Crops statement in the General Information section of the label.

EXTRAZINE II APPLIED ALONE

Use at the proper rate for soil texture and organic matter indicated in Table 1.

EXTRAZINE II COMBINATIONS

EXTRAZINE II plus LASSO 4EC

Use EXTRAZINE II DF at the proper rate for the soil texture and organic matter shown in Table 2 plus 2 quarts per acre of LASSO (Use 2.5 quarts LASSO on clay soils containing 5 percent organic matter and over).

EXTRAZINE II plus SUTAN+ or ERADICANE 6.7E

Use EXTRAZINE II DF at the proper rate for the soil texture and organic matter shown in Table 2 plus 1.8 quarts per acre of SUTAN+ or ERADICANE for control of many annual grasses and broadleaf weeds. (Use 2.4 quarts of SUTAN+ or ERADICANE on loam soils containing 5 percent or more organic matter.) Do not use on sands and loamy sands of less than 1 percent organic matter in the light sandy soils of eastern coastal states. Do not use on corn seed stock.

Apply before planting. Incorporate the mixture immediately upon application using power-driven cultivation equipment set for 2 to 3 inch depth, or a tandem disc set to cut to a depth of about 4 inches while operating at 4 to 6 mph. For thorough mixing, disc in two directions (cross disc), and follow with a harrow, drag, or other leveling device. Prior to the second discing, readjust the disc to prevent cutting deeper than 4 inches. EXTRAZINE II DF may be applied preemergence as an overlay over previous incorporated SUTAN+ or ERADICANE if desired.

Existing stands of quackgrass, purple and yellow nutsedge must be turned under and thoroughly chopped up prior to chemical treatments.

**TABLE 1
PREEMERGENCE BROADCAST APPLICATION RATES PER ACRE FOR EXTRAZINE II DF
APPLIED ALONE ON CORN**

Soil Texture Description	Pounds of EXTRAZINE II DF**					
	Percent Organic Matter in Soil*					
	Less than 1%	1%	2%	3%	4%	5% & Over
Sand, Loamy sand	DO NOT USE	1.4	1.7	2.2	2.8	3.6
Sandy loam	1.4	1.7	2.2	2.8	3.6	4.4
Loam, Silt loam, Silt	1.7	2.2	3.1	3.6	4.4	5.0
Sandy clay loam, Clay loam, Silty clay loam	2.2	3.1	3.6	4.4	5.0	5.3
Sandy clay, Silty clay, Clay	3.1	3.6	4.4	5.0	5.3	5.8
Peat or Muck	NOT RECOMMENDED					

* For organic matter content between those listed, adjust the rate proportionately.

** Maximum rate limits per acre per year for all applications is 6.5 lbs. cyanazine (9.6 lbs. EXTRAZINE II DF) except on highly erodible land with less than 30% plant residue cover, the rate limit is 3.0 lbs. cyanazine (4.4 lbs. EXTRAZINE II DF).

20712

TABLE 2

PREEMERGENCE BROADCAST APPLICATION RATES PER ACRE FOR EXTRAZINE II DF USED IN TANK-MIX COMBINATIONS WITH LASSO, SUTAN+, ERADICANE, OR DUAL ON CORN

Soil Texture Description	Pounds of EXTRAZINE II DF***						5% & Over
	Percent Organic Matter in Soil*						
	Less than 1%	1%	1%	2%	3%	4%	
Sand, Loamy sand	0.7**	0.8	1.4	1.7	1.9	2.2	
Sandy loam	0.8	1.4	1.7	1.9	2.2	2.5	
Loam, Silt loam, Silt	1.4	1.7	1.9	2.2	2.5	2.8	
Sandy clay loam, Clay loam, Silty clay loam	1.7	1.9	2.2	2.5	2.8	3.1	
Sandy clay, Silty clay, Clay		1.9	2.2	2.5	2.8	3.1	3.3
Peat or Muck	NOT RECOMMENDED						

* For organic matter content between those listed, adjust the rate proportionately.

**Do not use on the light sandy soils of the Atlantic Coastal Plain.

*** Maximum rate limits per acre per year for all applications is 6.5 lbs. cyanazine (9.6 lbs. EXTRAZINE II DF) except on highly erodible land with less than 30% plant residue cover, the rate limit is 3.0 lbs. cyanazine (4.4 lbs. EXTRAZINE II DF).

Additional weeds controlled by SUTAN+ or ERADICANE combinations:

Grasses

Sandbur	Shattercane (Wild Cane)*
Texas Panicum	Quackgrass (ERADICANE only)
	Wild Proso Millet*(ERADICANE only)

Perennial Weeds

- Yellow Nutsedge (nutgrass)
- Purple Nutsedge (nutgrass)

* Suppression only -- refer to SUTAN+ or ERADICANE label for appropriate supplement cultural and tillage practices.

For fields with moderate to heavy infestations of these weeds, refer to the SUTAN+ or ERADICANE labels for appropriate higher rates.

EXTRAZINE II PLUS DUAL 8E

Use EXTRAZINE II DF at the proper rate for soil texture and organic matter shown in Table 2. Use DUAL as follows:

SOIL TEXTURE	BROADCAST RATE PER ACRE FOR DUAL 8E
Coarse	
Sand, Loamy sand, Sandy loam	1.25 - 1.5 pints
Medium	
Loam, Silt loam, Silt	1.5 - 2.0 pints
Fine	
Sandy clay loam, Silty clay loam, Clay loam, Sandy clay, Silty clay, Clay	1.5 - 2.5 pints

The low end of the rate range should be used for lowest organic matter soils and the rate increased as organic matter increases to a point that soils containing 4 percent organic matter or more require the highest rate shown for that soil texture. Refer to the DUAL label for additional precautions on rotational crops.

CONSERVATION TILLAGE WEED CONTROL

(30 days prior to planting up to corn emergence)

EXTRAZINE II DF may be used for Early Preplant or Preemergence weed control for land going into the production of corn under conservation tillage (including no-till) programs. Complete any planned early spring tillage prior to application. Tillage after application may reduce the effectiveness of the herbicide treatment. EXTRAZINE II DF when used according to the following directions will kill most existing small weeds and suppress many emerged perennial weeds when corn is planted into no-till stalk ground (corn, sorghum), stubble ground (soybean, small grains), and any minimum-till situation. This treatment then provides residual control of annual weeds as in conventional tillage. A nitrogen solution or complete fertilizer solution may replace all or part of the water as a carrier. The spray gallonage and spray boom design must be adequate to give thorough uniform coverage of the weed foliage. Follow label requirements of all products used in tank mix combinations. Use the proper rate of EXTRAZINE II DF shown in Table 1. Where heavy crop residues exist, the rates shown in Table 1 should be increased by 25 percent.

Grass and Broadleaf Weeds Up to 3 Inches: Use EXTRAZINE II DF alone and add 1-2 quarts/acre of crop oil concentrate(COC) if weeds are emerged at the time of application. For best burndown results use a minimum of 20 gal/acre of liquid fertilizer as the carrier.

Broadleaf Weeds Exceed 3 Inches: If broadleaf weeds are taller than 3 inches at application, add 2,4-D LV Ester and/or BANVEL and non-ionic surfactant at recommended rates. Additional weeds controlled with 2,4-D are buckwheat, dandelion, dock, giant ragweed, mare's tail, pennycress, prickly lettuce and tansy mustard. To control existing alfalfa, add 0.3 to 0.5 pint/acre of BANVEL to the spray mixture. Apply before the alfalfa exceeds 6 inches in height.



Grass Weeds Exceed 3 Inches: If grass weeds are taller than 3 inches at application, add either GRAMOXONE EXTRA or ROUNDUP to the tank at the recommended rates for these products. Add 1 to 2 pints of a non-ionic surfactant per 100 gallons of spray. With GRAMOXONE EXTRA, well established weeds over 6 inches tall will not be well controlled. Do not apply GRAMOXONE EXTRA in a suspension type liquid fertilizer containing clay.

Burndown of Sod Grasses or Under Dry Conditions: For burndown of existing sod grasses such as orchardgrass, bromegrass, rye or timothy, or when very dry conditions exist, add GRAMOXONE EXTRA to the tank-mix at the recommended rates.

Perennial Grass Weeds: For improved control of perennial grasses such as johnsongrass or quackgrass, add ROUNDUP at the recommended rates or follow with a postemergence application of ACCENT.

Other Labeled Tankmixes: Other labeled products may be tankmixed with EXTRAZINE II DF according to the directions for those treatments in the Preemergence section of the label.

Early preplant applications of EXTRAZINE II DF may be tankmixed with 1 quart/acre of PRINCEP 4L or 1.1 pounds of PRINCEP CALIBER 90. Apply 30 days or more prior to planting.

Sequential Treatments: Depending upon the application timing and the weather conditions following an early preplant application, a sequential herbicide treatment may be necessary to provide additional length of weed control.

This may be a postemergence treatment with BLADEX or EXTRAZINE II or some other herbicide treatment applied at or after planting.

Rotational Crops: Refer to the General Information section of this label and to the labels of all products used in combination or in sequence.

POSTEMERGENCE

Under dry, arid conditions of low humidity and the absence of dew formation at night, add a nonionic surfactant or an emulsible vegetable (EV) oil suitable for use on growing corn at its recommended rate. Do not use petroleum-based crop oils. Addition of a surfactant or EV oil is not recommended under moist, rainy conditions and when dew forms at night as injury may occur.

Postemergence application must be made before corn reaches 12 inches in height, or before fifth leaf is visible.

Do not apply this treatment under cold, wet, weather conditions or to corn growing under stress caused by weather, insects, disease, etc. Yellowing of the corn may result from this treatment, particularly if cold, adverse growing conditions occur after application. Extended or extreme cold and wet conditions may reduce stands.

Do not apply postemergence on popcorn, sweet corn or corn grown for seed.

EXTRAZINE II APPLIED ALONE

Use EXTRAZINE II DF at the proper rate for the soil texture and organic matter shown in Table 3 or 4. Use rates shown in Table 3 if BLADEX or EXTRAZINE II has not been applied to the soil this season. Use rates shown in Table 4 if BLADEX or EXTRAZINE II has been applied to the soil this season. This treatment may be used on peat or muck soils for burndown and suppression of existing weeds but will not provide residual control. Apply from crop emergence through the four-leaf stage of corn growth before weeds exceed about 1-1/2 inches in height. Do not apply over the top of corn if the fifth leaf is visible or if corn exceeds 12 inches in height. Apply in water only. Do not spray emerged corn plants in a liquid fertilizer carrier or in tank mix combinations with EC formulation herbicides.

Rotational Crops: Refer to the General Information section of the label.

TABLE 3

**POSTEMERGENCE BROADCAST APPLICATION RATES PER ACRE FOR EXTRAZINE II DF ON CORN
NO PRIOR APPLICATION OF BLADEX OR EXTRAZINE II**

Soil Texture	Pounds of EXTRAZINE II DF			
	Percent Organic Matter in Soil*			
	Less than 1%	1%	2%	Over 2%
Sand, Loamy sand	DO NOT USE	1.3	1.8	2.2
Sandy loam	1.3	1.8	2.2	2.2
Loam, Silt loam, Silt	1.8	2.2	2.2	2.2
All other textures	2.2	2.2	2.2	2.2

* For organic matter content between those listed, adjust the rate proportionately.

10/12

TABLE 4

POSTEMERGENCE BROADCAST APPLICATION RATES PER ACRE FOR EXTRAZINE II DF ON CORN BLADEX OR EXTRAZINE II USED IN PRIOR APPLICATION

Soil Texture	Pounds of EXTRAZINE II DF**			
	Percent Organic Matter in Soil*			
	Less than 1%	1%	2%	Over 2%
Sand, Loamy sand	DO NOT USE	DO NOT USE	1.5	1.5
Sandy loam	DO NOT USE	DO NOT USE	1.75	2.2
Loam, Silt loam, Silt	DO NOT USE	1.5	2.2	2.2
All other textures	DO NOT USE	2.0	2.2	2.2

* For organic matter content between those listed, adjust the rate proportionately.

** Maximum rate limits per acre per year for all applications is 6.5 lbs. cyanazine (9.6 lbs. EXTRAZINE II DF) except on highly erodible land with less than 30% plant residue cover, the rate limit is 3.0 lbs. cyanazine (4.4 lbs. EXTRAZINE II DF).

SWEET CORN

EXTRAZINE II DF may be applied preemergence or preplant incorporated for the control of annual grasses and broadleaf weeds in sweet corn.

NOTE: EXTRAZINE II may cause injury or stand loss on new or "super sweet" varieties of sweet corn. Consult with Agricultural Extension Agencies and sweetcorn seed suppliers about the sensitivity of new varieties to potential injury.

Apply EXTRAZINE II treatments just before, at or after planting but before crop has emerged. Avoid removal of treated soil from seedrow prior to or during the planting operation. Do not apply postemergence to sweet corn.

Rotary hoeing is recommended for preemergence applications which do not receive adequate rainfall or sprinkler irrigation to wet the top 1 1/2 to 2 inches of soil within about 10 days after application.

If an EXTRAZINE II mixture is to be incorporated, except as noted, single or two pass incorporation is acceptable. Care should be taken to incorporate the EXTRAZINE II mixture no deeper than the top two inches of soil.

Rotational Crops: (1) Plant only corn, sorghum or soybeans the year following the use of this mixture. (2) If soybeans are to be planted, injury may occur due to the carryover of Atrazine. (3) If applied after June 10, do not rotate with crops other than corn or sorghum the next year or injury may occur. (4) Small grains may be planted 15 months following treatments. (5) All other crops may be planted 18 months after application.

EXTRAZINE II ALONE

Apply EXTRAZINE II DF at the proper rate for soil texture and organic matter indicated in Table 5.

EXTRAZINE II PLUS LASSO 4EC

Use EXTRAZINE II DF at the proper rate for the soil texture and organic matter shown in Table 6 plus 2 quarts per acre of LASSO (Use 2.5 quarts LASSO on clay soils containing 5 percent organic matter and over.)

EXTRAZINE II PLUS SUTAN+ OR ERADICANE 6.7E

Do not use combinations with SUTAN+ or ERADICANE in New Jersey.

Use EXTRAZINE II DF at the proper rate for the soil texture and organic matter shown in Table 6 plus 1.8 quarts per acre of SUTAN+ or ERADICANE for control of many annual grasses and broadleaf weeds. (Use 2.4 quarts of SUTAN+ or ERADICANE on loam soils containing 5 percent or more organic matter, and clay loams and clays containing 4 percent or more organic matter.) Do not use on sands and loamy sands having less than 1 percent organic matter nor on the light sandy soils of eastern coastal states. Do not use on corn grown for seed.

Apply before planting. Incorporate the mixture immediately upon application using power-driven cultivation equipment set for 2-3 inch depth, or tandem disc set to cut about 4 inches deep while operating at 4-6 mph. For thorough mixing, disc in two directions (cross disc), and follow with a harrow, drag, or other leveling device. Prior to the second discing, readjust the disc to prevent cutting deeper than 4 inches. EXTRAZINE II DF may be applied preemergence as an overlay over previously incorporated SUTAN+ or ERADICANE, if desired.

Existing stands of quackgrass, purple and yellow nutsedge must be turned under and thoroughly chopped up prior to chemical treatments.

Additional weeds controlled by SUTAN+ or ERADICANE combinations:

Grasses

Sandbur	Shattercane (Wild Cane)*
Texas Panicum	Quackgrass (ERADICANE only)
	Wild Proso Millet*(ERADICANE only)

Perennial Weeds

- Yellow Nutsedge (nutgrass)
- Purple Nutsedge (nutgrass)

* Suppression only -- refer to SUTAN+ or ERADICANE label for appropriate supplement cultural and tillage practices.

For fields with moderate to heavy infestations of these weeds refer to the SUTAN+ or ERADICANE labels for appropriate higher rates.

BEST AVAILABLE COPY

11-2-12

TABLE 5
PREEMERGENCE BROADCAST APPLICATION RATES PER ACRE FOR EXTRAZINE II DF
APPLIED ALONE ON SWEET CORN

Soil Texture Description	Pounds of EXTRAZINE II DF**					5% & Over
	Percent Organic Matter in Soil*					
	Less than 1%	1%	2%	3%	4%	
Sand, Loamy sand	DO NOT USE	1.3	1.6	2.1	2.6	3.5
Sandy loam	DO NOT USE	1.6	2.1	2.6	3.3	4.4
Loam, Silt loam, Silt	DO NOT USE	2.1	2.6	3.3	4.1	4.9
Sandy clay loam, Clay loam, Silty clay loam	DO NOT USE	2.6	3.1	4.1	4.9	5.4
Sandy clay, Silty clay, Clay	DO NOT USE	3.1	4.4	4.9	5.4	5.8
Peat or Muck	NOT RECOMMENDED					

* For organic matter content between those listed, adjust the rate proportionately.

** Maximum rate limits per acre per year for all applications is 6.5 lbs. cyanazine (9.6 lbs. EXTRAZINE II DF) except on highly erodible land with less than 30% plant residue cover, the rate limit is 3.0 lbs. cyanazine (4.4 lbs. EXTRAZINE II DF).

TABLE 6
PREEMERGENCE BROADCAST APPLICATION RATES PER ACRE FOR EXTRAZINE II DF USED IN
TANK MIX COMBINATIONS WITH LASSO, SUTAN+, ERADICANE OR DUAL ON SWEET CORN

Soil Texture Description	Pounds of EXTRAZINE II DF					5% & Over
	Percent Organic Matter in Soil*					
	Less than 1%	1%	2%	3%	4%	
Sand, Loamy sand	DO NOT USE	0.9	1.3	1.6	1.8	2.2
Sandy loam	DO NOT USE	1.3	1.6	1.8	2.2	2.4
Loam, Silt loam, Silt	DO NOT USE	1.6	1.8	2.2	2.4	2.9
Sandy clay loam, Clay loam, Silty clay loam	DO NOT USE	2.0	2.2	2.4	2.9	3.1
Sandy clay, Silty clay loam	DO NOT USE	2.2	2.7	2.9	3.1	3.3
Peat or Muck	NOT RECOMMENDED					

*For organic matter content between those listed, adjust the rate proportionately.

EXTRAZINE II PLUS DUAL 8E

Use EXTRAZINE II DF at the proper rate for soil texture and organic matter shown in Table 6. Use DUAL as follows:

SOIL TEXTURE	BROADCAST RATE PER ACRE FOR DUAL 8E
Coarse	
Sand, Loamy sand, Sandy loam	1.25 - 1.5 pints
Medium	
Loam, Silt loam, Silt	1.5 - 2.0 pints
Fine	
Sandy clay loam, Silty clay loam, Clay loam, Sandy clay, Silty clay, Clay	1.5 - 2.5 pints

The low end of the rate range should be used for lowest organic matter soils and the rate increased proportionately as the organic matter increases. Soils containing 4 percent organic matter or more require the highest rate shown for that soil texture. Refer to the DUAL label for precautions or additional information on rotational crops.

1204/2

STORAGE AND DISPOSAL

Storage: Store product in original container only. Do not contaminate water, other pesticides, fertilizer, food or feed in storage. Do not use or store around the home environment. Avoid contact with water. In case of spill or leak, soak up with sand, earth, or synthetic absorbent. Do not use alkaline absorbents and dispose of wastes in compliance with local, state and Federal regulations. Ground water contamination may be reduced by diking and flooring of permanent liquid bulk storage sites with an impermeable material.

Product Disposal: Pesticide, spray mixture, or rinsate that cannot be used according to label instructions must be disposed of according to applicable Federal, state, or local procedures.

Container Disposal: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

BLADEX., EXTRAZINE II, ACCENT

- Trademarks of E.I. Du Pont de Nemours and Co. (Inc.)

BANVEL Trademark of Sandoz Crop Protection Corp.

DUAL Trademark of CIBA-Geigy Corp.

ERADICANE Trademark of ICI Americas, Inc.

LASSO, ROUNDUP Trademarks of Monsanto Co.

PRINCEP, CALIBER Trademarks of CIBA-Geigy Corp.

SUTAN+. Trademark of ICI Americas, Inc.

D-091593/092893/101993

NOTICE OF WARRANTY

Du Pont warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for purposes stated on such label only when used in accordance with directions under normal use conditions. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Du Pont. In no case shall Du Pont be liable for consequential, special or indirect damages resulting from the use or handling of this product. All such risks shall be assumed by the buyer. **DU PONT MAKES NO WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.**

