	Ō	PM-	25		4-10	-98	F	10f 3
Please read instructions on	Form Approve	proved, OMB No. 2070-0060, Approval expires 05-31-98						
United States Environmental Protection Agency Weshington, DC 20460					Registrati Amendme Other			
		Application	n for Pestic	ide - Sectio	n 1			
1. Company/Product Number 352-580				2. EPA Product Menager J. A. Tompkins		3. Proposed Classifi		fication
4. Company/Product (Name DuPont Reliance® ST		РМ# 25						
5. Name and Address of Ap E. I. du Pont de Nemours Agricultural Products, P. C Wilmington, DE 19880-00 Attn: Jackie M. Marks, Wi	odei	6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No						
Check if this	s is a new address			uct Name				
			Section -	11				
Amendment - Explain Resubmission in resp Notification - Explain	[Final printed labels in response to Agency letter dated "Me Too" Application. APR 1 0 1998 Other - Explain below.						
Ref. letter from J. M. M.		•	Section -	188				
Child-Resistant Packaging	Unit Packaging	Water Soluble	Peckeging	2. Type of Container				
Yes* No	Yes No		Yes No			Metal Plastic Glass		
Certification must	lf "Yes" Unit Packaging wgt	No. per . container	lf "Yes" Package wgt	No.per container	Other (Specify)			
3. Location of Net Contents	4. Size(s) Reta	il Container	5.	5. Location of Label Directions On Label On Label On Labeling accompanying product				
6. Manner in Which Label is	Affixed to Product	Lithogra Paper g Stencie	lued	Other _	······································			
		•	Section - I	V.				
1. Contact Point (Complete	items directly below	for identification	of individual to l	e contected, if n	ecessery, to proce	ess this a	pplication.)	
Name Jacqueline M. Marks			Fite Product Regi		an (302) 992-6486			
l certify that the state I acknowledge that ar both under applicable	y knowingly false or i		attachments ti		-	ete.	5. Dete Appl Riceived (Stan	
2. Signature			3. Title Product Registration Technician					· · · · · · · · · · · · · · · · · · ·
4. Typed Neme Jacqueline M. Marks			5. Dete March 30, 1998					• • • • •
							•••	و

-

EPA Form 8570-1 (Rev. 8-94) Previous editions are obsciete.

÷

DUPOND DuPont Agricultural Products

"..... A Growing Partnership With Nature"



RELIANCE[®] STS⁻ SP TANK MIX WITH "PURSUIT" HERBICIDE FOR CONTROL OF NIGHTSHADE IN SOYBEANS NOT DESIGNATED AS STS

SUPPLEMENTAL LABELING

Z-0f3

RELIANCE® STS® SP EPA Reg. No. 352-580

TANK MIX OR SEQUENTIAL APPLICATION WITH A REDUCED RATE OF "PURSUIT" or "PURSUIT ECO PAK" HERBICIDE FOR CONTROL OF NIGHTSHADE IN SOYBEANS NOT DESIGNATED AS STS

DuPont "Reliance" STS SP Herbicide may be applied in a tank mix or the first application of a sequential program with a reduced rate of "Pursuit" or "Pursuit Eco Pak" for the control of nightshade in addition

those weeds listed on the "Reliance" STS SP label.

JIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

HOW TO USE

Tank Mix

"Reliance" STS SP, at a rate of 0.5 oz per acre or at the rate of 1 pack per 4 acres plus either "Pursuit" at a rate of 2.0 ounces per acre or "Pursuit Eco Pak" at a rate of 1 soluble bag per 10 acres can be tank mixed for postemerge control of the weeds listed on the "Reliance" STS SP labels and eastern black nightshade less than 2" in height. Refer to the "Reliance" STS SP, labels for other weeds controlled and maximum heights.

Best results are obtained when either "Reliance" STS SP plus either "Pursuit" or "Pursuit Eco Pak" is applied to weeds that are young (after the first true leaves have expanded, but before they exceed the size indirated on this label) and actively growing. This is generally 21-30 days iter planting of soybeans. Applications made to weeds that are in the cotyledon stage, to weeds larger than the sizes indicated, or to weeds under stress (weather, herbicide, or other) may result in unsatisfactory control.

Sequential Applications:

If eastern black nightshade becomes a problem following an application of "Reliance" STS SP, an application of a reduced rate of "Pursuit" or "Pursuit" Eco Pak can be made. "Pursuit" or "Pursuit" Eco Pak may be applied at a rate of 2.0 ounces per acre or 1 soluble bag per 10 acres, respectively, for control of small, actively growing eastern black nightshade that does not exceed 2" in height. A minimum interval of 7 days between "Reliance" STS SP application and follow-up "Pursuit" or "Pursuit" Eco Pak treatment is advised to reduce the potential for unsatisfactory weed control. Weeds should be free from stress and actively growing when the follow-up application is made. The soybean plants should have at least one newly emerged fully expanded trifoliate prior to the follow-up application of "Pursuit" or "Pursuit" Eco Pak. This program is recommended for the control of broadleaf weeds only. Other measures should be used to control grassy weeds.

ADJUVANTS: Postemerge applications of "Reliance" STS SP tank mixed or applied as the first application of a sequential program with either "Pursuit" or "Pursuit Eco Pak" must include the addition of a nonionic surfactant and ammonium nitrogen fertilizer.

- Use a nonionic surfactant at the rate of 1 pint per 100 gal of solution (0.125% v/v). Under dry, cool (generally 70° F or less) conditions the rate of nonionic surfactant may be increased to 2 pints per 100 gal. of solution (0.25% v/v).
- Use a high quality nitrogen fertilizer product such as 28-0-0 at a rate of 4 - 8 pints per acre, or 10-34-0 at a rate of 2 - 4 pints per acre. Alternately, a high-quality, sprayable grade of ammonium sulfate (21-0-0) may be used at a rate of 2 - 4 pounds per acre. Use the lower rate for spray volumes less than 15 gal. per acre.
- Do not use "Dash"², "Dash" HC², crop oil concentrates or methylated seed oil products such as "Sun-It" II⁹ with "Reliance" STS SP when applied alone as a first application of a planned program or tank mixed with either "Pursuit" or "Pursuit" Eco Pak as excessive crop injury may occur.

For additional information refer to the DuPont bulletin "Approved Adjuvants for Use With DuPont Row Crop and Cereal Herbicides" for a list of approved adjuvants and suggested use rates.

APPLICATION INFORMATION

Broadcast Application: Use flat fan nozzles at 25 - 40 psi. Do not use flood, hollow cone, rain drop, whirl chamber or controlled droplet applicator (CDA) type nozzles as unacceptable crop injury, excessive spray drift, or poor weed control may result. Use 10 - 25 gallons of water per acre. For proper spray coverage, adjust the boom and nozzle height according to the specifications listed by the nozzle manufacturer.

Band Application: For band application, use proportionately less spray mixture. To avoid crop injury, carefully calibrate the band applicator so as not to exceed the desired use rate. Carefully follow the manufacturer's instructions for nozzle type, (flat fan preferred), nozzle orientation, distance of the nozzles from the crop and weeds, spray volumes, calibration, and spray pressure. For additional information on row banders, see DuPont bulletin "Application Accuracy - Row Banders."

For product information call 1-888-6-DUPONT

© 1998 E. I. du Pont de Nemours and Company, Agricultural Products, Wilmington, Delavare: 19898 NOTIFICATION

H - 63388

Page 1 of 2

APR 1 0 1998

Aerial Application: Use nozzle types and arrangements that will provide for optimum spray distribution and maximum coverage at 5 to 10 gallons per acre. Do not apply during a temperature inversion, when winds are gusty, or when other conditions will favor poor coverage and/or off target spray movement. Use a minimum of 5 gallons of water per acre. Consult the respective product labels for special directions for aerial application.

IMPORTANT PRECAUTIONS

- Soybeans should be free from stress and actively growing at the time
 of application. Stress may be caused by abnormally hot or cold
 weather, growing conditions such as drought or water-saturated soil,
 disease, soil nutrient deficiencies such as iron chlorosis, or injury
 from nematodes, insects, or prior herbicide applications.
- Applications of "Reliance" STS SP when tank mixed with either "Pursuit" or "Pursuit Eco Pak" may shorten stem internodal length and cause temporary crop injury. Crop response may be increased when applications are made to soybeans that are under stress. Soybeans will recover quickly under normal growing conditions.
- Cultivation may put weeds under stress by pruning roots, thus reducing weed control. Avoid cultivation 7-10 days prior to or following application of any of the recommended herbicide treatments. For maximum weed control, cultivate 7-10 days after application.
- Apply the tank mix treatment or the first treatment in a planned sequential program after the first trifoliate of the soybean has fully expanded and the plants are actively growing, but before soybeans begin to flower.
- Refer to the "Reliance" STS SP and "Pursuit" or "Pursuit Eco Pak" Herbicide labels for additional use directions, use restrictions, rotational crop intervals and precautions. The most restrictive provision on either label will apply.
- Applications within 1 hour of rain may reduce weed control.

H - 63388 . . .

IMPORTANT

BEFORE USING "RELIANCE" STS SP, "PURSUIT" or "PURSUIT ECO PAK" HERBICIDES, READ AND FOLLOW ALL APPLICABLE DIRECTIONS, RESTRICTIONS, AND PRE-CAUTIONS ON THE EPA-REGISTERED LABELS.

This bulletin contains new or supplemental instructions for use of these products which do not appear on the EPA-registered package labels. Follow the instructions carefully.

This labeling must be in the possession of the user at the time of pesticide application.

- ¹ Registered trademark of American Cyanamid Company
- ² Registered trademark of BASF AG.
- ³ Registered trademark of Agsco, Inc.

110397, 033098 (Replaces H-63297)

12/11/97

For product information call 1-888-6-DUPONT © 1998 E. I. du Pont de Nemours and Company, Agricultural Products, Wilmington, Delaware 19898