



352-581

02/06/2003

1/24

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY

WASHINGTON, D.C. 20460

OFFICE OF
PREVENTION, PESTICIDES
AND TOXIC SUBSTANCES

Mr. Jack H. Cain
Product Registration Manager
DuPont Crop Protection
Stine-Haskell Research Center
P.O. Box 30
Newark, DE 19714-0030

FEB - 6 2003

Dear Mr. Cain:

Subject: DuPont® Velpar® DF Herbicide

EPA Registration No. 352-581

Application and Letter Dated December 19, 2002:

Request To Amend Registration by Revising Label To:

- o Updated resistance Management language to DuPont-Adopted and standardized, EPA approved language.
- o Addition of DuPont-adopted and standardized, EPA Approved Integrated Pest Management language.
- o Minor formatting and organizational changes.
- o Addition of Velpar® DF and diuron tank mix usage in LA and TX sugarcane.
- o Addition of Western Hemlock to Forestry -Release -Hardwood Suppression in Western U.S.
- o Addition of Supplemental Labels to Master Section 3 labeling:
 - o Highbush blueberries
 - o Alfalfa -Chemigation (With minor changes. Changes are highlighted.)
 - o Alfalfa - Velpar DF + Gramoxone
 - o Lowbush blueberries

The proposed subject labeling amendments for the pesticide product "DuPont Velpar® DF Herbicide" have been reviewed and found to be acceptable under the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA), as amended, provided that you:

1. At the next printing of the container label incorporate all use-patterns that are on the supplemental labeling listed above into a "primary" label and submit 5 copies of the revised labeling. Such labeling will be used by this Agency for administrative purposes.
2. Submit one (1) copy of the your final printed labeling before you release this product for shipment under the subject labeling.


2/26

-2-

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA, section 6(e). Your release for shipment of this product under this labeling constitutes acceptance of these conditions.

A stamped copy of the proposed labeling is enclosed for your records.

Sincerely yours,


for Joanne I. Miller
Product Manager (23)
Herbicide Branch
Registration Division (7505C)

Enclosure



DuPont™ Velpar® DF

herbicide

Dispersible Granules

<i>Active Ingredient</i>	<i>By Weight</i>
Hexazinone [3-cyclohexyl-6-(dimethylamino) -1-methyl-1,3,5-triazine-2,4(1H,3H)-dione]	75%
Inert Ingredients	25%
TOTAL	100%

EPA Reg. No. 352-581

**ACCEPTED
with COMMENTS
In EPA Letter Dated**

FEB - 6 2003

**Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.
352-581**

KEEP OUT OF REACH OF CHILDREN
DANGER PELIGRO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand this label, find someone to explain it to you in detail.)

FIRST AID

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

IF SWALLOWED: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-441-3637 for medical emergencies involving this product.

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS
AND DOMESTIC ANIMALS
DANGER! CAUSES EYE DAMAGE.**

Corrosive, causes irreversible eye damage. Harmful if swallowed. Do not get in eyes or on clothing. Avoid contact with skin. Wash thoroughly with soap and water after handling.

PERSONAL PROTECTIVE EQUIPMENT

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants.
- Shoes plus socks.
- Protective eyewear.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

USERS SHOULD: Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

The active ingredient, hexazinone, in this product is known to leach through soil into ground water under certain conditions as a result of agricultural use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in ground-water contamination.

GENERAL INFORMATION

DuPont™ VELPAR® DF herbicide is a water-dispersible granule that is mixed in water and applied as a spray for weed control in certain crops, Christmas trees, forestry site preparation and release areas, and industrial areas. It may also be applied as a basal soil treatment for brush control in reforestation areas, rangeland, pastures and noncrop areas.

VELPAR® DF is an effective general herbicide providing both contact and residual control of many annual and biennial weeds and woody plants. It is also effective for control of most perennial weeds.

VELPAR® DF is noncorrosive to equipment.

Caution should be exercised when applying VELPAR® DF near desirable trees or shrubs as they can absorb VELPAR® DF through roots extending in to treated areas.

Refer to supplemental labeling entitled "VELPAR® DF Herbicide Chemigation Use on Alfalfa" for use directions for chemigation. Do not apply this product through any irrigation system unless the supplemental labeling on chemigation is followed.

ENVIRONMENTAL CONDITIONS AND BIOLOGICAL ACTIVITY

VELPAR® DF is absorbed through the roots and foliage. Moisture is required to activate VELPAR® DF in the soil. Best results are obtained when the soil is moist at the time of application and 1/4–1/2 inches of rainfall occurs within 2 weeks after application.

For best results, apply VELPAR® DF preemergence or postemergence when weeds are less than 2 inches in height or diameter. Herbicidal activity is most effective under conditions of high temperature (above 80 °F), high humidity, and good soil moisture. Herbicidal activity may be reduced when vegetation is dormant, semi-dormant, or under stress (e.g. temperature or moisture).

Herbicidal activity will usually appear within 2 weeks after application to susceptible plants under warm, humid conditions; while 4–6 weeks may be required when weather is cool or dry, or when susceptible plants are under stress. If rainfall after application is inadequate to activate VELPAR® DF in the soil, plants may recover from contact effects and continue to grow.

On woody plants, symptoms usually appear within 3–6 weeks after sufficient rainfall has carried the herbicide into the root zone during periods of active growth. Defoliation and subsequent refoliation may occur, but susceptible plants are killed.

The degree and duration of control will depend on the following:

- Use rate
- Weed spectrum and size at time of application
- Environmental conditions at and following treatment

Where a rate range is shown, use the higher levels of the dosage range on hard-to-control species, fine-textured soils, or soils containing greater than 5% organic matter or carbon. Use the lower levels of the dosage range on coarse-textured soils and/or on soils low in organic matter. Refer to specific uses for rate ranges.

APPLICATION INFORMATION

VELPAR® DF may be applied by ground equipment and, where permitted, aerial equipment. Use rates, minimum spray gallonage, and other application information are described for various uses.

Dispose of the equipment washwater by applying it to a use-site listed on this label or in accordance with directions given in the "Storage and Disposal" section of this label.

MIXING

Before spraying, calibrate equipment to determine the quantity of water necessary to uniformly and thoroughly cover the vegetation and soil in a measured area to be treated. Make sure the volume of water is sufficient to completely suspend the VELPAR® DF.

RESISTANCE

When herbicides that affect the same biological site of action are used repeatedly over several years to control the same weed species in the same field, naturally-occurring resistant biotypes may survive a correctly applied herbicide treatment, propagate, and become dominant in that field. Adequate control of these resistant weed biotypes cannot be expected. If weed control is unsatisfactory, it may be necessary to retreat the problem area using a product affecting a different site of action.

To better manage herbicide resistance through delaying the proliferation and possible dominance of herbicide resistant weed biotypes, it may be necessary to change cultural practices within and between crop seasons such as using a combination of tillage, retreatment, tank-mix partners and/or sequential herbicide applications that have a different site of action. Do not let weed escapes go to seed. If applicable, see Weeds Controlled section of label for additional information on managing herbicide resistant weed biotypes.

It is advisable to keep accurate records of pesticides applied to individual fields to help obtain information on the spread and dispersal of resistant biotypes. Consult your agricultural dealer, consultant, applicator, and/or appropriate state agricultural extension service representative for specific alternative cultural practices or herbicide recommendations available in your area.

INTEGRATED PEST MANAGEMENT

DuPont recommends the use of Integrated Pest Management (IPM) programs to control pests. This product may be used as part of an Integrated Pest Management (IPM) program which can include biological, cultural, and genetic practices aimed at preventing economic pest damage. Application of this product should be based on IPM principles and practices including field scouting or other detection methods, correct target pest identification, population monitoring, and treating when target pest populations reach locally determined action thresholds. Consult your state cooperative extension service, professional consultants or other qualified authorities to determine appropriate action

moved
this
section

treatment threshold levels for treating specific pest/crop systems in your area.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

DuPont™ VELPAR® DF should be used only in accordance with recommendations on this label, or in supplemental DuPont labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

The correct use rates by crop and geographical area, specified on the label, and proper mixing/loading site considerations and application procedures must be followed to minimize potential for hexazinone movement into ground water. Users are encouraged to consult with their state Department of Agriculture, Extension Service, or other pesticide lead agency for information regarding soil permeability, aquifer vulnerability, and best management practices for their area.

AGRICULTURAL USES

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical resistant gloves in Category A (such as butyl rubber, natural rubber, neoprene rubber, or nitrile rubber) all ≥ 14 mls.
- Shoes plus socks
- Protective eyewear

ALFALFA

VELPAR® DF is recommended for control of certain weeds in established alfalfa grown for hay. Do not use on alfalfa grown for seed in any state except California.

APPLICATION TIMING*

NON-DORMANT AND SEMI-DORMANT VARIETIES

In the following states, make a single application during winter months when alfalfa plants are in the least active stage of growth:

Arizona	Montana	Oklahoma	Washington
California	Nebraska	Oregon	Wyoming
Colorado	Nevada	South Dakota	
Idaho	New Mexico	Texas	
Kansas	North Dakota	Utah	

In the following states, make a single application in the spring before new growth exceeds 2 inches in height or to stubble after cutting following hay removal and before regrowth exceeds 2 inches in height:

Connecticut	Maine	New Hampshire	Vermont
Delaware	Maryland	New Jersey	Virginia
Illinois	Massachusetts	New York	West Virginia
Indiana	Michigan	Ohio	Wisconsin
Iowa	Minnesota	Pennsylvania	
Kentucky	Missouri	Rhode Island	

Severe alfalfa injury may result following application after cutting if either the regrowth is more than 2 inches high, significant stubble is left after alfalfa cutting, or the air temperature is above 90 °F.

DORMANT VARIETIES

Make a single application after alfalfa becomes dormant and before new growth begins in the spring. Where weeds have emerged, use a surfactant.

USE RATES

Use higher rates on hard-to-control species, fine textured soils, soils containing greater than 5% organic matter, or under adverse environmental conditions such as temperature extremes or when weeds are stressed due to low rainfall.

Select the appropriate dose for soil texture and organic matter content as follows:

Soils	VELPAR® DF (Lb/Acre)*		
	Percent Organic Matter in Soil <1%	1-5%	>5%
Coarse Texture			
Loamy sand, sandy loam	2/3 - 1	2/3 - 1	1 1/3 - 2
Medium Texture			
Loam, silt loam, silt, clay loam, sandy clay loam	2/3 - 1	1 - 2	1 1/3 - 2
Fine Texture			
Silty clay loam, sandy clay, silty clay, clay	1 - 2	1 - 2	1 1/3 - 2

* Do not exceed 1 lb per acre on alfalfa less than one year old.

WEEDS CONTROLLED

VELPAR® DF, when applied preemergence or early postemergence at the following rates is recommended for the control or suppression of the following species:

1/3 - 2/3 Lb/Acre

Tansy-mustard

2/3 - 1 1/3 Lb/Acre

Annual bluegrass	Jim Hill mustard
Blue mustard	London rocket
Cheatgrass (downy brome)	Miners lettuce
Common chickweed	Salsify
Common groundsel	Seedling orchardgrass
Dogfennel (mayweed)	Shepherd's-purse
English catchfly	Spurry
Fiddleneck (tarweed)	Tansy-mustard
Field pennycress	Wild radish
Filaree	Yellow rocket
Flixweed	

1 1/3 - 2 Lb/Acre

Common dandelion* (i.e. Canada bluegrass)	Perennial bluegrass* (spring only)
False dandelion	Prickly lettuce*
Foxtail (Setaria spp)*	Ryegrass, annual
Mexican tea*	Quackgrass*
White cockle*	Seedling alfalfa*

* Partial control

DuPont™ VELPAR® DF, when applied in late spring or after cutting at the following rates, will control these species:

2/3 - 2 Lb/Acre

Common lambsquarters	Foxtail (Setaria spp.)
Crabgrass	Jimsonweed
Fleabane	Redroot pigweed

SPRAY EQUIPMENT

Apply VELPAR® DF using a fixed boom power sprayer or aerial equipment.

MIXING INSTRUCTIONS

Add VELPAR® DF to a water-filled tank and mix it thoroughly. Apply it in at least 20 gal of water per acre by ground or 5 to 10 gal of water per acre by air.

REPLANTING (FOLLOWING ALFALFA)

- Do not replant treated areas to any crop except corn, root crops or sugarcane within two years after treatment, as crop injury may result.
- Corn may be planted 12 months after the last treatment, provided the use rate did not exceed 1 lb per acre, except in areas of low rainfall (20 inches or less).
- Root crops such as potatoes, sugarbeets, radish and carrots may be planted 12 months after last treatment, provided the use rate not exceed 2/3 lb per acre. Sites with use rates higher than 2/3 lb per acre should not be replanted to any crop within 2 years of application, or unacceptable crop injury may result.
- Sugarcane may be planted any time following treatment.
- In California, do not replant seed alfalfa areas to any crop within two years after treatment, as crop injury may result.

USE PRECAUTIONS - ALFALFA ONLY

Best results are obtained when 1/2-1 inches of rainfall or sprinkler irrigation occurs within two weeks after application, when soil is moist at time of application, and when weeds have not germinated or are less than 2 inches in height or diameter. Heavy rainfall or excessive irrigation after application may result in crop injury or poor performance of the herbicide.

- Do not apply to snow-covered or frozen ground.
- Since the effect of VELPAR® DF on alfalfa varies with soil conditions, uniformity of application, and environmental conditions, growers should limit their first use to small areas.
- If abnormally dry conditions exist following application, restrict the first irrigation to no more than 1/2 acre inch of water.
- Temporary yellowing of alfalfa may occur following VELPAR® DF applications.
- Treat only stands of alfalfa established for one year or for one growing season (except in California), provided the following conditions are met:
 - The alfalfa root system must be well established.
 - The crop must be healthy, vigorous and not under stress from adverse weather conditions, insect or disease damage.
 - The alfalfa crop must be free of excessive winter injury from freezing and thawing.
- In California, fall planted alfalfa may be treated in the following winter months with VELPAR DF at 1/3 to 2/3 lb per acre to suppress or control Black Mustard, Common groundsel, London rocket, Shepard's-purse, Common chickweed, Fiddleneck or Filaree, provided:
 - alfalfa root growth exceeds 6 inches in length
 - vegetative top growth of alfalfa has lateral development of secondary growth
 - alfalfa is healthy and vigorous, not growing under stress from insect, disease, winter injury or other types of stress.
 Injury may result to alfalfa plants that fail to meet these growth criterion listed above.
- Do not use VELPAR® DF on seedling alfalfa, alfalfa-grass mixtures, or other mixed stands as injury may result to the seedling alfalfa or companion crop.
- At elevations above 4,000 feet, do not use VELPAR® DF on alfalfa that has been established with a cover or nurse crop until the alfalfa has gone through two summer growing seasons.
- Do not use VELPAR® DF in low desert valleys in California or Arizona.
- Do not add a surfactant to VELPAR® DF when treating nondormant alfalfa.
- Do not use VELPAR® DF on gravelly or rocky soils, exposed subsoils, hardpan, sand, poorly drained soil, or alkali soils.
- Crop injury, including mortality, may result in fields with restricted root growth due to nonuniform soil profiles such as gravel bases and clay lenses.
- Do not graze or feed forage or hay to livestock within 30 days following application.

SEED ALFALFA (CALIFORNIA ONLY)

- ADDITIONAL USE PRECAUTIONS

- Do not use DuPont™ VELPAR® DF on fields with sandy loam or loamy sand soils having less than 1% organic matter.
- Do not exceed 2/3 lb per acre on fields with sandy loam or loamy sand soils having 1-2% organic matter.
- Do not exceed 2/3 lb per acre on seed alfalfa that has been established for only one growing season.

ALFALFA - IMPREGNATION ON DRY BULK FERTILIZER (EXCEPT CALIFORNIA AND ARIZONA)

Dry bulk fertilizer may be impregnated or coated with VELPAR® DF for application to established alfalfa. All recommendations, cautions and special precautions on this label must be followed along with state regulations relating to dry bulk fertilizer blending, impregnating and labeling.

If fertilizer materials are excessively dusty, use a suitable additive to reduce dust prior to impregnation, as dusty fertilizer will result in poor distribution during application. The dry fertilizer must be properly impregnated and uniformly applied to the alfalfa to avoid crop injury and/or poor weed control.

To impregnate the fertilizer, use a system consisting of a conveyor or closed drum used to blend dry bulk fertilizer. Any commonly used fertilizer can be impregnated with VELPAR® DF, except potassium nitrate or sodium nitrate. Do not use VELPAR® DF on limestone.

Use a minimum of 250 lb dry bulk fertilizer per acre and up to a maximum of 450 lb per acre. To impregnate or coat the dry bulk fertilizer with VELPAR® DF, mix 2 2/3 pounds of VELPAR® DF with sufficient water to make one gallon of suspension and thoroughly agitate. Direct the nozzles to deliver a fine spray of this suspension toward the fertilizer for thorough coverage while avoiding spray contact with mixing equipment. Uniform impregnation of VELPAR® DF to dry bulk fertilizer will vary, and if the absorptivity is not adequate, the use of an absorptive powder may be required to produce a dry, free-flowing mixture. "Microcel E" is the recommended absorbent powder. When another herbicide is used with VELPAR® DF, mix and impregnate the fertilizer immediately.

Apply impregnated fertilizer as soon as possible after impregnation for optimum performance.

Select the rate of VELPAR® DF to apply per acre from the appropriate section of this label. Then refer to the rate chart below to determine the amount of VELPAR® DF that should be impregnated on a ton of dry bulk fertilizer, based on the amount of fertilizer to be distributed in one acre.

Rate Chart for Impregnating Fertilizer with VELPAR® DF

Fertilizer Rate/Acre	VELPAR® DF Rate Per Acre			
	2/3 Lbs	1 Lbs	1 1/3 Lbs	2 Lbs
250 lbs	5.3 lbs/ton	8.0 lbs/ton	10.6 lbs/ton	16.0 lbs/ton
300 lbs	4.4 lbs/ton	6.6 lbs/ton	8.8 lbs/ton	13.3 lbs/ton
350 lbs	3.7 lbs/ton	5.7 lbs/ton	7.5 lbs/ton	11.4 lbs/ton
400 lbs	3.3 lbs/ton	5.0 lbs/ton	6.7 lbs/ton	10.0 lbs/ton
450 lbs	2.9 lbs/ton	4.4 lbs/ton	5.9 lbs/ton	8.9 lbs/ton

For rates other than those listed, use the following formula to calculate the amounts of VELPAR® DF to be impregnated per ton of dry fertilizer.

$$\text{VELPAR® DF} \times \frac{\text{2000}}{\text{Lbs Fertilizer Per Acre}} = \frac{\text{Lbs}}{\text{Ton of Fertilizer}} \text{ VELPAR® DF Per}$$

APPLICATION

Uniform application of VELPAR® DF-impregnated dry fertilizer is essential for satisfactory weed control. Accurate calibration of the application equipment is essential for uniform distribution to the surface. The recommended method of application is to apply 1/2 the recommended rate and overlap 50%. This results in the best distribution pattern.

CHRISTMAS TREES

VELPAR® DF is recommended for control of certain weeds where the following species are grown:

- | | |
|-------------------|----------------|
| Austrian pine | Noble fir |
| Douglas fir | Ponderosa pine |
| (western US only) | Scotch pine |
| Grand fir | Sitka spruce |
| Loblolly pine | |

Do not use VELPAR® DF on Christmas trees in the following states:

- | | | | |
|-------------|---------------|----------------|---------------|
| Alabama | Louisiana | New Jersey | Texas |
| Arkansas | Maine | New York | Vermont |
| Connecticut | Maryland | North Carolina | Virginia |
| Delaware | Massachusetts | Pennsylvania | West Virginia |
| Georgia | Mississippi | Rhode Island | |
| Florida | New Hampshire | South Carolina | |

APPLICATION TIMING

EASTERN US

Apply VELPAR® DF as a broadcast spray in the spring prior to bud break. If application is made after bud break, use directional spray equipment to prevent contact with foliage.

WESTERN US

Areas of greater than 20 inches annual rainfall - VELPAR® DF may be applied as a broadcast spray in the spring prior to conifer bud break. If application is made after bud break, use directional spray equipment to prevent contact with foliage.

Areas of less than 20 inches annual rainfall - VELPAR® DF may be applied in the fall before the soil freezes or in the spring after snow cover melts, but before conifer bud break occurs.

USE RATES

The rates listed below are for broadcast application. For band application, use proportionately less; for example, use 1/2 of the broadcast rates when treating a 3-foot band where row spacing is 6 feet.

Do not use more than one application of VELPAR® DF per year.

DuPont™ VELPAR® DF (Lb/Acre)		
Soils	First Year Plantings	Established Trees
Coarse Texture		
Loamy sand, sandy loam (50-85% sand)	1 1/3	1 1/3 - 1 2/3
Medium Texture		
Loam, silt loam silt, clay loam, sandy clay loam	1 1/3 - 1 2/3	1 2/3 - 2 1/3
Fine Texture		
Silty clay loam, clay loam, sandy clay, silty clay, clay	1 2/3 - 2	2 1/3 - 2 2/3

First year plantings - Transplant stock that is 2 years old or more (1 year old for loblolly pine). Apply VELPAR® DF only if rainfall has settled the soil around the base and root systems of the transplants.

Established trees - Trees that have been planted in the plantation for 1 year or more.

WEEDS CONTROLLED

VELPAR® DF is recommended for the control or suppression of the following species:

- | | |
|----------------------------|------------------------|
| Annual bluegrass | Fleabane |
| Barnyardgrass | Foxtail |
| Bentgrass | Goldenrod* |
| Bromegrass | Heath aster* |
| Catscar* (false dandelion) | Horseweed* |
| Common groundsel | Orchardgrass* |
| Common ragweed | Oxeye daisy |
| Crabgrass* | Pennsylvania smartweed |
| Curly dock* | Ryegrass* |
| Dandelion* | Velvetgrass |
| Fescue* | Wild carrot* |
| Fireweed (willowweed)* | |

* Partial control

SPRAY EQUIPMENT

VELPAR® DF may be applied by ground equipment or by air.

MIXING INSTRUCTIONS

Select a spray volume that will ensure a thorough and uniform application.

USE PRECAUTIONS

- CHRISTMAS TREES

- Do not use VELPAR® DF in nurseries, seed beds, or ornamental plantings.
- Do not add a surfactant in applications over the top of conifers.
- Weed control results from spring applications depend on sufficient moisture to activate VELPAR® DF.
- Do not cut treated vegetation for forage or hay nor graze domestic animals on treated areas for 60 days following application.
- Poor weed and brush control may result from the following:
 - Heavy duff or slash present at the time of application.

-Use on poorly drained sites.

-Applications made when soil is saturated with water and rain is imminent within 24 hours.

-Applications to soils high in organic matter (greater than 5%).

- Injury may occur when VELPAR® DF is used on the following:

-Trees that show poor vigor, insect damage, disease, winter injury, or other stress conditions.

-Any soil containing less than 1% organic matter.

-Loamy sand or sandy loam with less than 2% organic matter (except Jeffrey Pine and Ponderosa Pine).

-Foliage after bud break.

-Gravelly or rocky soils, exposed subsoils, clay knobs, sand, or sandy soil with 85% or more sand.

PINEAPPLE

VELPAR® DF is recommended for control of certain weeds in pineapple.

APPLICATION TIMING

- USE RATES - MIXING INSTRUCTIONS

Use a sprayer properly calibrated to a constant speed and rate of delivery.

Mix the proper amount of VELPAR® DF in water. Add a surfactant at 0.25% by volume of water.

- Intercrop period - Apply VELPAR® DF as a broadcast spray in 100-400 gal of water per acre at the rate of 1/3 - 2 1/3 lb per acre. For aerial application, use at least 10 gal water per acre.
- Post mulch, preplant - Apply VELPAR® DF as a broadcast spray in 100-400 gal of water per acre at the rate of 1/3 - 2 1/3 lb per acre.
- Post plant, before planted cuttings start active growth - Apply VELPAR® DF as a broadcast spray in 100-400 gal of water per acre at the rate of 1/3 - 2 1/3 lb per acre.

A post-plant application should be made after planted cuttings start to grow only when weed growth has escaped control by other herbicide applications.
- Prior to forcing first ratoon - Apply VELPAR® DF as a broadcast spray in 100-400 gal of water per acre at the rate of 1/3 - 2 1/3 lb per acre.
- Directed postemergence (pineapple and weeds) inter-space application - Apply VELPAR® DF as a directed spray 3-10 months after planting in 50-200 gal of water per acre (broadcast basis) at the rate of 1/3 - 2 1/3 lb per acre (broadcast basis) using a stroller boom or knapsack.
- Directed spot treatments for perennial grasses before floral induction - Spray perennial grasses postemergence to wet (50-200 gals per acre depending on size) with 1 1/3 - 2 1/3 lb per 100 gal of water as a spot treatment.
- Treatments to field edges and roadsides - Apply VELPAR® DF at 2 1/3 - 4 8/10 lb per acre in 100-400 gal of water.

WEEDS CONTROLLED

DuPont™ VELPAR® DF is recommended for the control or suppression of the following species:

Ageratum	Kao haole*
Balsam apple	Mauna loa*
Castor bean	Morningglory
Crabgrass	Oxalis
Crotalaria	Popolo
Dallisgrass	Richardsonium
Guinea grass	Vaseygrass
Jungle rice	

* Suppression

USE PRECAUTIONS - PINEAPPLE

- Use the lower rates on coarse-textured soils or in areas where rainfall exceeds 65 inches per year.
- Use the higher rates on fine-textured soils or in areas where rainfall is less than 65 inches per year.
- Do not exceed 4.8 lb VELPAR® DF per acre per crop.
- Do not apply VELPAR® DF within 181 days of harvest.

SUGARCANE

VELPAR® DF is recommended for selective weed control in sugarcane except in the State of Florida.

APPLICATION TIMING

- USE RATES - MIXING INSTRUCTIONS

Apply a single treatment of VELPAR® DF per year using a fixed-boom sprayer and a minimum of 25 gal per acre unless otherwise directed.

HAWAII

Apply VELPAR® DF pre- or postemergence at the following rates for the indicated soil texture:

Soils	VELPAR® DF (Lb/Acre) (Plus surfactant 0.25% by volume)
Coarse Texture	
Sand, loamy sand, sandy loam	2/3 - 1 2/10
Medium Texture	
Loam, silt loam, silty clay loam	2/3 - 2 1/3
Fine Texture	
Clay, gray hydromorphic clay	2 1/3 - 4 8/10

Use the higher levels of the recommended dosage ranges on soils high in organic matter. Do not apply more than twice the highest recommended rate for the indicated soil texture per crop (18–24 months).

A surfactant is recommended for all uses.

For preemergence use only, VELPAR® DF may be applied with aerial equipment using at least 10 gal of spray per acre.

For spot treatments of emerged weeds, VELPAR® DF may be applied with a knapsack sprayer in concentrations of 0.6 - 4.8 lb per 100 gal of water. Apply a sufficient volume to thoroughly wet weed foliage, but do not exceed 40 gal of spray per treated acre. Use the lower concentrations on coarse-textured soils that are low in organic matter, and use the higher concentrations on fine-textured soils that are high in organic matter.

LOUISIANA

Apply 2/3 - 1 2/10 lb of VELPAR® DF per acre broadcast in the fall before sugarcane emerges or in the spring before active cane tillering begins. Fall treatments of 2/3 - 1 2/10 lb per acre may be followed by a spring treatment of 2/3 - 1 2/10 lb per acre. Do not apply more than 2 lb per year. Use the higher levels of the recommended dosage range on fine-textured soils.

PUERTO RICO

For preemergence treatments, apply 1/3 - 2/3 lb of VELPAR® DF per acre.

For postemergence treatments, apply 1/3 - 2/3 lb of VELPAR® DF per acre to weeds after they have emerged. Use the lower rates on coarse-textured soils and the higher rates on fine-textured soils (high in clay or organic matter). Each ratoon may receive up to 2/3 lb of VELPAR® DF per acre.

For spot treatment of emerged weeds, VELPAR® DF may be applied with a knapsack sprayer in concentrations of 1/3 - 2/3 lb per 100 gal of water. Apply a sufficient volume to wet the weed foliage. Do not exceed 100 gal of spray per treated acre. Use the lower concentration on coarse-textured soils and the higher concentration on fine-textured soils.

Note: Since it is difficult to calibrate "spot" knapsack applications, extra care must be taken not to exceed the rate equivalent of the maximum of 2/3 lb VELPAR® DF per acre.

Do not apply more than 1 1/3 lb of VELPAR® DF per acre per crop season.

TEXAS

Apply 2/3 - 2 1/3 lb of VELPAR® DF per acre. On plant cane, apply the herbicide before the cane emerges or as a directed layby treatment. On stubble cane, apply VELPAR® DF preemergence (up to the 3-leaf stage) or as a directed layby treatment. A pre- or early postemergence treatment may be followed by a layby treatment, provided at least 60 days have elapsed and 3 inches of rainfall or sprinkler irrigation have occurred since the first treatment.

Do not apply more than 2 1/3 lb of VELPAR® DF per acre per season.

Use the following rates according to the different soil textures:

Soils	VELPAR® DF (Lb/Acre)	
	Preemergence	Layby
Coarse Texture*		
Sandy loam	1/3	1/3
Medium Texture		
Loam, silt loam	9/10	9/10
Fine Texture		
Clay loam	1 1/3	1 1/3

* With at least 2% organic matter

On dormant cane, a surfactant may be added to the spray mixture to increase control of emerged weeds.

WEEDS CONTROLLED

DuPont™ VELPAR® DF is recommended for the control or suppression of the following species:

Ageratum*	Johnsongrass (from seed)
Alexandergrass	Jungle rice
Amaranth (slender, smooth)	Lambsquarter
American burnweed (fireweed)	Morningglory (hairy, three-lobed)
Balsam apple	Mustard (wild)
Barnyardgrass	Oxalis
Bermudagrass*	Panicum (brownleaf, browntop, Texas millet)
Carolina geranium	Paspalum (ricegrass, sour)
Chickweed	Pigweed (common, smooth)
Crabgrass (hairy, large, smooth)	Popolo
Crotalaria (fuzzy, showy)	Purslane
Cuphea (tarweed)	Sandbur
Dallisgrass	Sensitive plant (hila hila)
Fingergrass (radiate, swollen)	Signalgrass (broadleaf)
Flora's paintbrush	Sowthistle
Foxtail (bristly, yellow)	Spanish needle
Goosegrass	Sprangletop
Guineagrass	Spurge (prostrate, graceful)
Henbit	Sunflower
Itchgrass*	Vaseygrass
Jobs tears	Waltheria (hialoa)

* Partial control

TANK MIXTURES (LOUISIANA AND TEXAS)

A tank mixture of VELPAR® DF at 0.67 pounds per acre plus diuron ("Karmex" DF or "Direx" DF) at 2.25 lbs per acre may be applied to newly planted sugarcane prior to crop emergence or to ratoon (stubble) sugarcane following harvest for improved control of annual sedge, cutleaf eveningprimrose, red morningglory, pitted morningglory, Texas panicum, smellmelon, swinecress, toadflax and also provide additional suppression of itchgrass.

An early Spring application of the tank mix combination may be used postemergence (over the top) of sugarcane until it reaches a height of 18 inches. If weeds are present at the time of application, a non-ionic surfactant at the use rate of 0.25% V/V or a crop oil concentrate at the use rate of 1% V/V is recommended. If a tank mix partner is being used, follow the most restrictive adjuvant recommendation. Temporary crop chlorosis may result from postemergence applications to emerged sugarcane.

The tank mix combination may also be used as a post-directed/layby application. For best weed control, post-directed/layby applications should be made prior to weed germination. Post-directed/layby applications are required when the sugarcane plants reach or exceed a height of 18 inches. Post-directed/layby applications should be directed at the base of the sugarcane plants so the spray does not contact green leaves or the plant whorl.

If weeds are present at application, a tank mixture partner, labeled for use on sugarcane, is recommended. When adding other tank mix partners, read and follow all use instructions, warnings and precautions on the companion label(s).

Do not apply as a postemergence (over-the-top) application to sugarcane beyond a height of 18 inches.

The tank mix combination of VELPAR® plus diuron may be applied to fallow sugarcane fields up to 60 days prior to planting. For best results, applications should be made to a newly prepared seedbed that is free of clods and existing vegetation. If weeds are present, either a separate application of a contact herbicide or a tank mix partner is recommended for improved postemergence control.

USE PRECAUTIONS - SUGARCANE

Do not plant any crop other than sugarcane following an application of VELPAR® DF.

Do not feed sugarcane forage to livestock.

Do not apply VELPAR® DF:

- Within 180 days of harvest in Hawaii.
- Within 234 days of harvest in Louisiana.
- Within 288 days of harvest in Puerto Rico.
- Within 234 days of harvest in Texas.

To avoid injury to sugarcane, observe the following precautions:

- Do not use VELPAR® DF on cane that shows poor vigor because of insect damage, disease, or winter injury, or shows symptoms of other stress conditions such as drought stress.
- Do not add a surfactant in applications unless otherwise specified.
- Do not use VELPAR® DF on gravelly or rocky soils, thinly covered subsoils, or coarse-textured soils (sands to sandy loams) with less than 1% organic matter.
- Temporary chlorosis of the crop may result from application over emerged cane. Applications during active cane growth should be directed to cover the weeds and soil while minimizing crop contact.
- Do not use VELPAR® DF on varieties known to be susceptible to weed killers.

Extremely heavy rainfall after application may result in poor weed control and/or crop injury, especially if the application is made to dry soil.

FORESTRY

SITE PREPARATION

VELPAR® DF is recommended for weed and brush control in areas where the following species are grown:

EASTERN US AND LAKE STATES

Austrian pine	Red spruce
Balsam fir	Scotch pine
Black spruce	Shortleaf pine
Loblolly pine	Slash pine
Longleaf pine	Spruce
Ponderosa pine	Virginia pine
Red pine	White spruce

WESTERN US

- Blue spruce
- Douglas fir
- Engleman spruce
- Grand fir
- Jeffrey pine
- Lodgepole pine
- Noble fir
- Ponderosa pine
- Sitka spruce
- White fir

APPLICATION TIMING

EASTERN US

Apply DuPont™ VELPAR® DF from early spring to early summer after hardwoods have broken bud and before the foliage has hardened off.

WESTERN US

Rainbelt (areas of high spring rainfall): For best results, apply in late winter or spring when weeds and brush are actively growing.

Snowbelt (areas of low spring rainfall): For best results, apply in the fall before soil freezes, or in the spring after snow cover melts in anticipation of rainfall. Weed and brush control results from spring applications will be dependent on sufficient rainfall following application to activate VELPAR® DF.

USE RATES

Use the lower rates on coarse-textured soils and soils low in organic matter; use the higher rates on fine-textured soils and on soils high in organic matter. Use the higher rates where hard-to-kill species predominate.

The rates listed below are for broadcast application.

Soils	VELPAR® DF (Lb/Acre)	
	Eastern US	Western US
Coarse Texture		
Sand, loamy sand, sandy loam	2 2/3 - 4	1 1/3 - 1 1/2
Medium Texture		
Loam, silt loam, sandy clay loam	4 - 5 1/3	1 1/2 - 2 2/3
Fine Texture		
Silty clay loam, clay loam, sandy clay, silt, silty clay, clay	5 1/3 - 6 2/3	2 2/3 - 4

PLANTS CONTROLLED

VELPAR® DF is recommended for the control or suppression of the following species:

Herbaceous Plants

- Annual bluegrass
- Asters
- Barnyardgrass
- Bentgrass
- Bromegrass
- Canada thistle*
- Catsear (false dandelion)*
- Common groundsel
- Common ragweed
- Crabgrass*
- Curly dock*
- Dandelion*
- Elksedge
- Fescue*
- Fireweed (willowweed)*
- Fleabane
- Foxtail
- Goldenrod*
- Heath aster*
- Horseweed*
- Orchardgrass*
- Oxeye daisy
- Pennsylvania smartweed*
- Pinegrass
- Quackgrass*
- Ryegrass*
- Squawcarpet
- Velvetgrass
- Wild carrot

*Partial control; use higher rates for soil textures indicated above.

Woody Plants

- Ash
- Aspen (big tooth, trembling)
- Balsam poplar
- Birch
- Blackgum
- Deerbrush ceanothus
- Elm
- Flowering dogwood*
- Greenleaf manzanita
- Hawthorne
- Hazel
- Hickory
- Honeysuckle*
- Oaks
- Red maple*
- Snowbush ceanothus (varnishleaf ceanothus)
- Sourwood*
- Sweetgum
- Whitehorn
- Wild cherry
- Willows

*Partial control; use the higher rates for the soil textures indicated above.

Within several weeks after VELPAR® DF activation by rainfall, affected vegetation may be burned, if desired. This burn may further enhance control of vegetation. Burn the vegetation only after any residual stand is completely defoliated, at least twice, allowing for sufficient root uptake of VELPAR® DF. In the West, results may take one to two years in areas of low rainfall.

SPRAY EQUIPMENT

When applied as a liquid spray using water as the carrier, VELPAR® DF may be applied by ground equipment or by air (helicopter only).

MIXING INSTRUCTIONS

For ground application, use enough water for thorough coverage, usually a minimum of 25 gal per acre. For aerial applications, use at least 5 gal of water per acre.

SITE PREPARATION - TANK MIXTURES

VELPAR® DF PLUS "Tordon 101" MIXTURE

For improved, broad-spectrum brush control, tank mix VELPAR® DF with "Tordon 101" Mixture. Only the following trees may be planted on sites with this combination:

- Austrian pine
- Loblolly pine
- Longleaf pine
- Ponderosa pine
- Red pine
- Scotch pine
- Shortleaf pine
- Slash pine
- Virginia pine

APPLICATION TIMING

Apply DuPont™ VELPAR® DF plus “Tordon 101” Mixture from late spring to early summer after the trees have reached full leaf, but before leaf tissue hardens. In the states of Maine, Michigan, Minnesota, New Hampshire, New York, Vermont, and Wisconsin, apply this tank mixture after bud break.

WOODY PLANTS CONTROLLED - USE RATES

Predominant Species	DuPont™ VELPAR® DF (Lb/Acre)	“Tordon 101” M (Qt/Acre)
Blackberry, Elm, Oak (Prunus spp.), Sweetgum	4 - 6 2/3	4
Blackgum, Dogwood, Pine, Red maple, Sassafras, Sourwood	2 2/3 - 6 2/3	6-8

Refer to the USE RATES table in the FORESTRY SITE PREPARATION section of this label for recommended rates based on soils.

SPRAY EQUIPMENT

VELPAR® DF plus “Tordon 101” Mixture may be applied by ground equipment or by air (helicopter only).

MIXING INSTRUCTIONS

For ground application, use enough water for thorough coverage, usually a minimum of 25 gal per acre. For aerial applications, use at least 5 gal of water per acre. For best results, full foliar coverage is required.

USE PRECAUTIONS**- VELPAR® DF + “Tordon 101” Mixture**

- To avoid injury, do not plant pine before the following intervals following application with “Tordon 101” Mixture:
 - Southern states - 6 months
 - Lake states - 9 months
 - Northeastern states - 9 months
- Burning treated sites after initial contact (browning) effect on foliage may result in poor control. Burn the vegetation only after brush has completely defoliated at least once, allowing sufficient root uptake of VELPAR® DF (usually 60-90 days after adequate rainfall).
- Before using “Tordon 101” Mixture, read and carefully observe the cautionary statements and all other information appearing on the product label. “Tordon 101” Mixture is a restricted use pesticide, for retail sale to and use only by certified applicators or persons under their direct supervision and only for those uses covered by certified applicator’s certification.

SITE PREPARATION**- GRID & SINGLE STEM APPLICATIONS**

VELPAR® DF is recommended for control of brush in site preparation.

GRID APPLICATION

Mix 2 2/3 pounds of VELPAR® DF with sufficient water to make one gallon of suspension and thoroughly agitate.

Intermittent agitation may be required to maintain the VELPAR® DF in suspension.

Apply the VELPAR® DF suspension directly to the soil surface in a grid pattern using an exact delivery handgun applicator. This equipment delivers a thin stream of predetermined volume. VELPAR® DF should be applied during the period from hardwood bud break to early summer.

Application rate and grid pattern will depend on soil texture and woody plant composition. Use the lower rates on coarse textured soils and when the major component of the hardwoods are susceptible species. Use the high rates on fine-textured soils and hard-to-control species.

Application Patterns and Rates For VELPAR® DF Suspension

	ML/Spot	Grid (Ft)	Qt/Acre
Coarse	0.6	3 X 3	3
	2.0	4 X 4	6
	3.1	4 X 6	6
Medium/Fine	1.6	3 X 3	8
	2.8	4 X 4	8
	3.5	4 X 4	10
	5.2	4 X 6	10

SINGLE STEM (BASAL SOIL)

Mix 2 2/3 pounds of VELPAR® DF with sufficient water to make one gallon of suspension and thoroughly agitate. Apply the VELPAR® DF suspension with an exact-delivery handgun applicator. This equipment delivers a thin stream of predetermined volume when triggered. Apply the VELPAR® DF suspension at the rate of 2 to 4 ml for each inch of stem diameter at breast height. Direct the treatment to the soil within 3 feet of the root collar of woody plants to be controlled.

For multi-stemmed and low-growing brush that have stem diameters that are difficult to determine, apply the VELPAR® DF suspension at the rate of 2 to 4 ml per 3 feet of canopy width. For tall, slender (columnar) brush types, apply 4 to 8 ml per 3 feet of height. Base the rate on whichever canopy dimension is greater (width or height).

When treating brush that requires more than a single delivery of the VELPAR® DF suspension, apply subsequent deliveries equally spaced around the target plant. If treating brush on sloping sites, apply most of the suspension on the uphill side of the stem. If treating resprouts from brush disturbed by cutting or other mechanical methods, the rate of application should be proportional to the original tree size, not just the size of sprout regrowth.

USE PRECAUTIONS**- SITE PREPARATION**

Where burning is desired, burn the vegetation only after any residual brush has completely defoliated, at least twice, allowing for sufficient root uptake of VELPAR® DF.

Following harvest, allow sufficient time for stumps and injured trees to adequately resprout before applying VELPAR® DF.

**RELEASE
- HARDWOOD SUPPRESSION**

DuPont™ VELPAR® DF is recommended for conifer release where the following species are grown:

EASTERN US

Balsam fir	Red spruce
Black spruce	Shortleaf pine
Loblolly pine	Slash pine
Longleaf pine	Virginia pine
Norway spruce	White spruce
Red pine	

WESTERN US

Blue spruce	Noble fir
Douglas fir	Ponderosa pine
Engleman spruce	Sitka spruce
Grand fir	Western hemlock
Jeffrey pine	White fir
Lodgepole pine	

APPLICATION TIMING

EASTERN US

Apply VELPAR® DF from early spring to early summer after hardwoods have broken bud and before full leaf expansion.

Applications made over the top of pines may result in excessive pine injury under conditions of high humidity and temperature (80 degrees F).

WESTERN US

Rainbelt (areas of high spring rainfall): For best results, apply in late winter or spring when brush is actively growing, but prior to conifer budbreak. If application is made after bud break, use directional spray equipment to prevent contact with conifer foliage, as injury may result.

Snowbelt (areas of low spring rainfall): For best results, apply in the fall before soil freezes and after the final resting bud has hardened on the conifers. Or, spring applications may be made after snow cover melts in anticipation of rainfall prior to conifer budbreak. Brush control results from spring treatments will be dependent on sufficient rainfall following application to activate VELPAR® DF.

USE RATES

The rates listed below are for broadcast application. Do not use more than one application of VELPAR® DF per year.

EASTERN US

Crop Species	Soil Description	VELPAR® DF (Lb/Acre) Established Trees
Loblolly pine	Loamy sand, sandy loam	1 1/3 - 2
Longleaf pine		
Shortleaf pine	Loam, silt loam,	
Virginia pine	silt, sandy clay loam	1 1/3 - 2 2/3
Slash pine	Silty clay loam, clay loam, sandy clay, silty clay, clay	3 - 4
Red pine	Loamy sand, sandy loam	1 1/3 - 2 2/3
	Loam, silt loam, silt, sandy clay loam	2 2/3 - 4
	Silty clay loam, clay loam, sandy clay, silty clay, clay	4 - 5 1/3

Established Trees - 4 years of age from transplanting on coarse-textured soils
 - 3 years of age from transplanting on medium-textured soils
 - 2 years of age from transplanting for Red Pine

WESTERN US

Crop Species	Soil Description	VELPAR® DF (Lb/Acre)
Blue spruce	Loamy sand, sandy loam	1 1/3 - 3
Douglas fir		
Engleman spruce	Loam, silt loam, sandy clay loam	2 2/3 - 4
Grand fir		
Jeffrey pine		
Lodgepole pine	Silt, silty clay loam, clay loam,	
Noble fir	sandy clay, silty clay, clay	3 - 4
Ponderosa pine		
Sitka spruce		
Western hemlock		
White fir		

For first year plantings utilizing bare root stock, treat only transplant stock that is 2 years old (2-0, 1-1) or more, except (1-0) for Ponderosa and Jeffrey pines. Apply VELPAR® DF only if rainfall has settled the soil around the base and root systems of the transplants.

BRUSH SUPPRESSION

VELPAR® DF is recommended for the control or suppression of the following species:

Ash	Hawthorne
Aspen	Hazel
Balsam poplar	Honeysuckle
Birch	Oaks
Box elder	Red maple*
Brambles	Snowbush ceanothus
Cherry (black, pin)	Sumac*
Deerbrush ceanothus	Sweetgum*
Dogwood*	Whitehorn
Elm	Willow
Greenleaf manzanita	

* Partial control

In addition to brush controlled, herbaceous species listed in Weeds Controlled section of Release-Herbaceous Weed Control may be controlled with these applications.

SPRAY EQUIPMENT

When applied as a liquid spray using water as the carrier, DuPont™ VELPAR® DF may be applied by ground equipment or by air (helicopter only).

MIXING INSTRUCTIONS

For ground application, use enough water for thorough coverage, usually a minimum of 25 gal per acre. For aerial applications, use at least 5 gal of water per acre.

RELEASE

- GRID & SINGLE STEM APPLICATIONS

VELPAR® DF is recommended for hardwood suppression in conifer release sites.

GRID APPLICATION

Mix 2 2/3 pounds of VELPAR® DF with sufficient water to make one gallon of suspension and thoroughly agitate. Intermittent agitation may be required to maintain the VELPAR® DF in suspension.

Apply the VELPAR® DF suspension directly to the soil surface in a grid pattern using an exact delivery handgun applicator. This equipment delivers a thin stream of predetermined volume. VELPAR® DF should be applied during the period from hardwood bud break to early summer.

Application rate and grid pattern will depend on soil texture and woody plant composition. Use the lower rates on coarse textured soils and when the major component of the hardwoods are susceptible species. Use the high rates on fine-textured soils and hard-to-control species.

Application Patterns and Rates For VELPAR® DF Suspension

	ML/Spot	Grid (Ft)	Qt/Acre
Coarse	0.5	3 X 4	3*
	1.2	3 X 6	6
	2.1	4 X 6	6
Medium/Fine	1.2	3 X 3	8
	2.3	3 X 6	8
	1.6	3 X 3	10
	3.1	3 X 6	10

* Use on deep sands with pines four years or more of age.

SINGLE STEM (BASAL SOIL)

Mix 2 2/3 pounds of VELPAR® DF with sufficient water to make one gallon of suspension and thoroughly agitate. Apply the VELPAR® DF suspension with an exact-delivery handgun applicator. This equipment delivers a thin stream of predetermined volume when triggered. Apply the VELPAR® DF suspension at the rate of 2 to 4 ml for each inch of stem diameter at breast height. Direct the treatment to the soil within 3 feet of the root collar of woody plants to be controlled.

For multi-stemmed and low-growing brush that have stem diameters that are difficult to determine, apply the VELPAR® DF suspension at the rate of 2 to 4 ml per 3 feet of canopy width. For tall, slender (columnar) brush types, apply 4 to 8 ml per 3 feet of height. Base the rate on whichever canopy dimension is greater (width or height).

When treating brush that requires more than a single delivery of the VELPAR® DF suspension, apply subsequent deliveries equally spaced around the target plant. If treating brush on sloping sites, apply most of the suspension on the uphill side of the stem. If treating resprouts from brush disturbed by cutting or other mechanical methods, the rate of application should be proportional to the original tree size, not just the size of sprout regrowth.

USE PRECAUTIONS

- RELEASE GRID & SINGLE STEM

- Application of VELPAR® DF spots closer than 36 inches to conifer seedlings in their first season or directly up slope from these seedlings may result in injury or mortality.
- Use VELPAR® DF on seedlings in their first or fourth year and older. Injury may result from use on two and three year old seedlings where root growth is extensive but hardiness is lacking.

RELEASE

- HERBACEOUS WEED CONTROL

VELPAR® DF is recommended for controlling herbaceous weeds where the following species are grown:

EASTERN US

- | | |
|---------------|------------|
| Loblolly pine | Slash pine |
| Longleaf pine | Red pine |

WESTERN US

- | | |
|-----------------|-----------------|
| Blue spruce | Noble fir |
| Douglas fir | Ponderosa pine |
| Engleman spruce | Sitka spruce |
| Grand fir | Western hemlock |
| Jeffrey pine | White fir |
| Lodgepole pine | |

APPLICATION TIMING

EASTERN US

Apply VELPAR® DF as a broadcast or banded spray in the spring prior to conifer bud break to lessen conifer injury potential.

WESTERN US

Rainbelt (areas of high spring rainfall): For best results, apply as a broadcast or banded spray in the late winter or spring when weeds are actively growing, but prior to conifer budbreak. If application is made after conifer bud break, use directional spray equipment to prevent contact with conifer foliage, as injury may result.

Snowbelt (areas of low spring rainfall): For best results, apply as a broadcast or banded spray in the fall before soil freezes and after the final resting bud has hardened on the conifers. Or, spring applications may be made after snow cover melts in anticipation of rainfall prior to conifer budbreak. Weed control results from spring treatments will be dependent on sufficient rainfall following application to activate VELPAR® DF.

USE RATES

The rates listed below are for broadcast application. For band application, use proportionately less. For example, use 1/2 of the broadcast rates when treating a 3-foot band where row spacing is 6 feet.

EASTERN US

Soil Description	DuPont™ VELPAR® DF (Lb/Acre)	
	First Year Plantings	Established Trees
Loamy sand, sandy loam(50-85% sand)	1 1/3	1 1/3 - 1 2/3
Loam, silt loam, silt, sandy clay loam	1 1/3 - 1 1/2	1 2/3 - 2 1/3
Silty clay loam, clay loam, sandy clay, silty clay, clay	1 1/2 - 1 8/10	2 1/3 - 2 2/3

Red pine only - Refer to recommended rates in the HARDWOOD SUPPRESSION - Eastern US table on page 11.

WESTERN US

Refer to recommended rates in the HARDWOOD SUPPRESSION-Western US table on page 11.

WEEDS CONTROLLED

VELPAR® DF is recommended for the control or suppression of the following species:

Annual bluegrass	Fireweed (willowweed)*
Aster	Fleabane
Barnyardgrass	Foxtail
Bentgrass	Goldenrod*
Bracken Fern	Heath aster*
Bromegrass	Horseweed*
Catsear (false dandelion)	Orchardgrass*
Common groundsel	Oxeye daisy
Common ragweed	Panicums
Crabgrass*	Pennsylvania smartweed
Curly dock*	Ryegrass*
Dandelion*	Squawcarpet
Dogfennel	Velvetgrass
Fescue*	Wild carrot*

* Partial control

FORESTRY—IMPREGNATION ON DRY BULK FERTILIZER

VELPAR® DF is recommended for impregnating or coating dry bulk fertilizer to be applied on forested sites for the establishment or release of conifer plantations (except longleaf pine) as specified on this label.

PLANTS CONTROLLED

Fertilizer impregnated with VELPAR® DF is recommended for the control and suppression of the weeds and brush identified for the specific applications on this label. Consult the appropriate segment of this label to determine the appropriate rate of VELPAR® DF to be applied per acre. Apply this amount of VELPAR® DF to the volume of fertilizer to be applied per acre.

IMPREGNATION EQUIPMENT

To impregnate or coat the fertilizer use a system consisting of conveyor or closed drum used to blend dry bulk fertilizer.

IMPREGNATION INSTRUCTIONS

To impregnate dry bulk fertilizer with VELPAR® DF, mix the amount as prescribed above in a sufficient quantity of water to uniformly coat the desired amount of fertilizer. Suspensions of VELPAR® DF will require thorough agitation.

Direct the spray nozzles of the impregnation equipment to deliver a fine spray of the mixture toward the fertilizer for thorough coverage while avoiding contact with mixing equipment. The use of a colorant or dye may be beneficial to visually determine the uniformity of impregnation.

Uniform impregnation of dry bulk fertilizer may vary. If absorption of the spray is not adequate, the use of an absorptive powder or additive, such as “Microcel E” or “HiSil 233”, may be required to produce a dry, free flowing mixture.

Apply the fertilizer as soon as possible after impregnation for optimum performance. Impregnated fertilizer may become lumpy and difficult to apply following storage.

Diammonium phosphate, potassium chloride, 16-16-16 and 24-4-4 have been successfully impregnated.

APPLICATION EQUIPMENT

Applications of impregnated fertilizer may be made by ground equipment or by air (helicopter or fixed wing). Accurate calibration and patterning of the equipment is essential for uniform distribution of the impregnated fertilizer on the soil surface.

USE PRECAUTIONS—IMPREGNATED FERTILIZER FOR FORESTRY

- If fertilizer materials are excessively dusty, use a suitable additive to reduce dust prior to impregnation. Application of dusty fertilizer which has been impregnated may result in off-target drift and injury to desirable vegetation. Such drift and associated injury may be aggravated by high wind conditions.
- The dry fertilizer must be properly impregnated and uniformly applied to avoid pine injury/mortality and poor weed and brush control.
- Uniform and precise application of the impregnated fertilizer is essential for satisfactory weed and brush control and to minimize pine injury. Overlaps or skips between adjoining swaths or non-uniform distribution of impregnated fertilizer within the swath will deliver poor results and may result in pine injury or mortality.
- Do not impregnate potassium nitrate, sodium nitrate or triple super phosphate fertilizers with VELPAR® DF as herbicidal action will be lost.

USE PRECAUTIONS - FORESTRY

- Do not use VELPAR® DF in nurseries, seedbeds, or ornamental plantings.
- On tracts of land where various soil types are present and VELPAR® DF rate selection is difficult, conifer damage

or less-than-expected vegetation suppression may occur due to the different rates required for various soil types.

- Poor weed and brush control may result from the following:
 - Heavy duff or slash present at time of application
 - Use on poorly drained sites
 - Applications made when the soil is saturated with water and rain is imminent within 24 hours.
 - Applications to soils high in organic matter (greater than 5%).
- Following harvest, allow stumps and injured trees sufficient time to adequately resprout before applying DuPont™ VELPAR® DF.
- Where burning is desired, burn vegetation after any brush has completely defoliated, at least twice, allowing for sufficient root uptake of VELPAR® DF.
- Do not use VELPAR® DF on frozen soils; use in spring after snow melt.
- Do not add a surfactant in applications over the top of conifers.
- Weed control results from spring applications depend on sufficient moisture to activate VELPAR® DF.
- When applying VELPAR® DF after transplanting, wait until rainfall has settled the soil around the base and root systems of the transplants before making the treatment.
- Crop injury may occur when VELPAR® DF is used:
 - On trees that show poor vigor, insect damage, disease, winter injury, or other stress conditions
 - On any soil containing less than 1% organic matter
 - On loamy sand or sandy loam with less than 2% organic matter, except Jeffrey pine and Ponderosa pine
 - On conifer foliage after conifer bud break.
 - On gravelly or rocky soils, exposed subsoils, clay knobs, sand, or sandy soil with 85% or more sand.
- Do not cut treated vegetation for forage or hay nor graze domestic animals on treated areas for 60 days following application.

NON-AGRICULTURAL USES

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Industrial and Pasture/Rangeland weed and brush control applications as described on this label for DuPont VELPAR DF are not within the scope of the Worker Protection Standard.

The area being treated must be vacated by unprotected persons.

Do not enter or allow entry into treated areas until sprays have dried to perform hand tasks.

APPLICATION INFORMATION

VELPAR® DF is recommended for general weed and brush control in noncrop sites such as railroads, highways, utility and pipeline rights-of-way, petroleum tank farms, storage areas, industrial plant sites, and other similar areas.

NONCROP, INDUSTRIAL SITES

VELPAR® DF is recommended for control of many annual, biennial, and perennial weeds in noncrop, industrial sites.

APPLICATION TIMING

Apply VELPAR® DF as a preemergence or postemergence spray when weeds are actively germinating or growing.

WEEDS CONTROLLED - USE RATE

VELPAR® DF effectively controls the following weeds when applied at the use rates shown. When applied at lower rates, VELPAR® DF provides short-term control of the weeds listed; when applied at higher rates, weed control is increased and extended.

Use lower rate on coarse-textured soils (sand to sandy loam). Use the higher rate on fine-textured soils (clay loam to clay) and on soils high in organic matter.

2 2/3 - 6 2/3 Lb/Acre

Barnyardgrass	Lespedeza
Bindweed*	Milkweed*
Bouncingbet*	Mustard
Bromegrass	Nutsedge*
Buffalograss*	Orchardgrass*
Burdock	Oxalis
Cocklebur	Paragrass
Crabgrass	Pigweed
Crown vetch	Purslane
Curly dock*	Quackgrass
Dandelion*	Ryegrass, annual
Dogbane*	Smartweed
Fiddleneck	Spurge
Filaree	Star thistle
Fleabane	Trumpet creeper*
Goatsbeard vine	Wild oats*
Goldenrod	Wild parsnip

8 - 10 2/3 Lb/Acre

Bahiagrass*	Guineagrass
Bermudagrass*	Heath aster
Blackberry	Honeysuckle
Bluegrass	Lantana
Broomsedge	Marestail
Camphorweed	Natalgrass
Canada thistle*	Plantain
Chickweed	Prickly lettuce
Clovers	Ragweed
Dewberry	Smutgrass†
Dogfennel*	Spanish needle
Fescue*	Vaseygrass
Fingergrass	Wild carrot
Foxtail	

* Partial control

† Partial control may result with some of the giant (larger) smutgrass species.

SPECIFIC WEED PROBLEMS

Control of Canada Thistle in Crown Vetch - VELPAR® DF is recommended for control of Canada thistle in established stands of crown vetch on noncrop sites. Make a single application of 1

- 1 2/3 lb of DuPont™ VELPAR® DF from late spring through mid-summer, when thistle is actively growing prior to flowering. Do not use a surfactant. Some discoloration of the crown vetch foliage may occur after application.

SPRAY EQUIPMENT

Apply VELPAR® DF uniformly over the desired area using ground equipment or helicopter. Do not apply more than 8 lbs per acre by air.

MIXING INSTRUCTIONS

Use enough water for thorough coverage. For ground application this is usually a minimum of 25 gal per acre. Higher application volumes may be needed to obtain uniform application with handgun equipment. For aerial applications (helicopter only) this is usually a minimum of 5 gal per acre. Higher volumes of water may be needed when water temperatures are cold or the higher rates of VELPAR DF are used.

INDUSTRIAL TURF (UNIMPROVED ONLY)

VELPAR® DF is recommended for selective weed control in established stands of bermudagrass and/or bahiagrass in noncrop areas.

APPLICATION TIMING

Make a single application of VELPAR® DF per year when weeds are actively growing.

WEEDS CONTROLLED - USE RATE

VELPAR® DF effectively controls the following weeds at the rates shown. Use a lower rate on coarse-textured soils (sand to sandy loam). Use the higher rate on fine-textured soils (clay loam to clay) and on soils high in organic matter.

9/10 - 1 1/2 Lb/Acre

Barnyardgrass	Maypop (passion flower)
Dogfennel	Oxalis
Fescue	Pepperweed
Lespedeza	Pigweed
Little barley	Smutgrass †

† Partial control may result with some of the giant (larger) smutgrass species.

SPRAY EQUIPMENT

Apply VELPAR® DF uniformly over the desired area using ground equipment only.

MIXING INSTRUCTIONS

For ground application, use enough water for thorough coverage usually a minimum of 25 gal per acre. The use of a surfactant is not recommended.

USE PRECAUTIONS

- INDUSTRIAL UNIMPROVED TURF

- Use VELPAR® DF only in stands of bermudagrass and bahiagrass established for at least one year. Do not treat newly sprigged or sodded areas.
- Some discoloration of the bermudagrass or bahiagrass may occur after application.
- Injury may result when desirable grasses are under stress from drought, insects, disease, cold temperature, or poor fertility.
- Severe turf injury may occur if applications are made on gravelly or rocky soils, thinly covered subsoils, or soils with less than 1% organic matter.

BRUSH CONTROL

VELPAR® DF is recommended for the control of undesirable woody plants in noncrop sites.

APPLICATION TIMING

Apply VELPAR® DF from late winter through summer, prebud break until new growth hardens off.

In areas where the soil remains frozen during the winter and spring rains are usually inadequate for soil activation, a fall or winter treatment may be applied before the soil freezes.

WOODY PLANTS CONTROLLED - USE RATE

VELPAR® DF is recommended for the control or suppression of the following species. Use lower rate on coarse-textured soils (sand to sandy loam). Use the higher rate on fine-textured soils (clay loam to clay) and on soils high in organic matter.

5 1/3 - 10 2/3 Lb/Acre

Alder	Manzanita
American elm	Mesquite
Ash	Mulberry
Aspen	Multiflora rose
Balsam poplar	Myrtle
Birch	Oaks
Black cherry	Osage orange
Blackgum	Persimmon
Catclaw acacia	Privet
Chinaberry*	Red maple
Chinese elm	Sassafras*
Chinese tallow	Small soapweed (yucca)
Deerbrush	Snowbrush
Dogwood	Sourwood
Eastern red cedar*	Sumac
Hackberry	Sweet bay
Hawthorne	Sweet gum
Hazel	Whitebrush
Hickory	Whitehorn
Huisache	Wild plum
Juniper	Willow
Locust	Yellow Poplar
Lotebush	

*Partial control

SPRAY EQUIPMENT AND APPLICATION TECHNIQUES

BROADCAST

Apply 5 1/3 to 10 2/3 lb of VELPAR® DF per acre as a coarse spray by ground equipment or 5 1/3 to 8 lb per acre by air (helicopter only). Use enough water for thorough coverage. For ground equipment, usually a minimum of 25 gal per acre. For aerial equipment, usually a minimum of 10 gal per acre. Higher volumes of water may be needed when water temperatures are cold or the higher rates of VELPAR® DF are used.

BASAL (SOIL)

SINGLE STEM - Mix 2 2/3 pounds of DuPont™ VELPAR® DF with sufficient water to make one gallon of suspension and thoroughly agitate. Apply the VELPAR® DF suspension with an exact-delivery handgun applicator. This equipment delivers a thin stream of predetermined volume when triggered. Apply the VELPAR® DF suspension at the rate of 2 to 4 ml for each inch of stem diameter at breast height. Direct the treatment to the soil within 3 feet of the root collar of woody plants to be controlled.

For multi-stemmed and low-growing brush that have stem diameters that are difficult to determine, apply the VELPAR® DF suspension at the rate of 2 to 4 ml per 3 feet of canopy width. For tall, slender (columnar) brush types, apply 4 to 8 ml per 3 feet of height. Base the rate on whichever canopy dimension is greater (width or height).

When treating brush that requires more than a single delivery of the VELPAR® DF suspension, apply subsequent deliveries equally spaced around the target plant. If treating brush on sloping sites, apply most of the suspension on the uphill side of the stem. If treating resprouts from brush disturbed by cutting or other mechanical methods, the rate of application should be proportional to the original tree size, not just the size of sprout regrowth.

LACING/STREAKING - Mix VELPAR® DF with water to form a concentrated suspension. Apply 5 1/3 to 10 2/3 lbs of VELPAR® DF per acre. Adjust the application equipment to deliver a narrow or straight stream spray pattern such that the swath width on the soil surface is 6 to 12 inches wide. Direct the spray at the base of the brush. Swaths or treated bands should be 2 to 4 feet apart.

USE PRECAUTIONS - NONCROP

- Injury to or loss of desirable trees or other plants may result if VELPAR® DF is applied or if equipment is drained or flushed on or near desirable trees or other plants, on areas where their roots may extend, or in locations where the chemical may be washed or moved into contact with their roots.
- Prevent spray from drifting to desirable plants.
- Poor weed and brush control may result from the following:
 - Use on poorly drained sites
 - Applications made when the soil is saturated with water and rain is imminent within 24 hours.
 - Applications to soils high in organic matter (greater than 5%).
- Following mechanical cutting or clearing, allow stumps and injured trees sufficient time to adequately resprout before applying VELPAR® DF.
- Do not use VELPAR® DF on frozen soils.
- Do not use VELPAR® DF on lawns, driveways, tennis courts, or other residential or recreational areas.
- Weed and brush control results from spring applications depend on sufficient moisture to activate VELPAR® DF.

- Do not cut treated vegetation for forage or hay nor graze domestic animals on treated areas for 60 days following application. For rates above 8 lb per acre, do not cut treated vegetation for forage or hay nor graze domestic animals for 1 year.

PASTURE

VELPAR® DF is recommended for control of brush and weeds in pasture.

BERMUDAGRASS/BAHIAGRASS PASTURES

VELPAR® DF is recommended for control of smutgrass and other weeds in established stands of bermudagrass and bahiagrass.

APPLICATION TIMING

Make a single application of VELPAR® DF per year when weeds are actively growing.

WEEDS CONTROLLED - USE RATES

VELPAR® DF effectively controls the following weeds at the rates shown. Use a lower rate on coarse-textured soils (sand to sandy loam). Use the higher rate on fine-textured soils (clay loam to clay) and on soils high in organic matter.

9/10 - 1 1/2 Lb/Acre

Barnyardgrass	Maypop (passion flower)
Dogfennel	Oxalis
Fescue	Pepperweed
Lespedeza	Pigweed
Little barley	Smutgrass*

- * Partial control may result with some of the giant (larger) smutgrass species.

SPRAY EQUIPMENT

Apply VELPAR® DF uniformly over the desired area using ground equipment only.

MIXING INSTRUCTIONS

For ground application, use enough water for thorough coverage usually a minimum of 25 gal per acre. The use of a surfactant is not recommended.

USE PRECAUTIONS - PASTURE

- Use VELPAR® DF only in stands of bermudagrass and bahiagrass established for at least one year. Do not treat newly sprigged or sodded areas.
- Some temporary discoloration of the bermudagrass or bahiagrass may occur after application.
- Treatment of mixed pastures containing forage species other than bermudagrass or bahiagrass may result in injury or mortality to the other forage species.
- Injury may result when desirable grasses are under stress from drought, insects, disease, cold temperature, or poor fertility.
- Injury to or loss of desirable trees or other plants may result if VELPAR® DF is applied or if equipment is drained or flushed on or near desirable trees or other plants, on areas where their roots may extend, or in

locations where the chemical may be washed or moved into contact with their roots.

- Severe crop injury may occur if applications are made on gravelly or rocky soils, thinly covered subsoils, or soils with less than 1% organic matter.
- For broadcast pasture applications of DuPont™ VELPAR® DF, do not cut treated vegetation for forage or hay nor graze domestic animals on treated areas for 60 days.

BRUSH CONTROL PASTURE/RANGELAND

VELPAR® DF is recommended for the control of undesirable woody plants in pasture or rangeland.

APPLICATION TIMING

Apply VELPAR® DF from late winter through summer, pre-budbreak until new growth hardens off.

In areas where the soil remains frozen during the winter and spring rains are usually inadequate for soil activation, a fall or winter treatment may be applied before the soil freezes.

WOODY PLANTS CONTROLLED

VELPAR® DF is recommended for the control or suppression of the following species:

Alder	Manzanita
American elm	Mesquite
Ash	Mulberry
Aspen	Multiflora rose
Balsam poplar	Myrtle
Birch	Oaks
Black cherry	Osage orange
Blackgum	Persimmon
Catclaw acacia	Privet
Chinaberry*	Red maple
Chinese elm	Sassafras*
Chinese tallow	Small soapweed
Deerbrush	Snowbrush
Dogwood	Sourwood
Eastern red cedar*	Sumac
Hackberry	Sweet bay
Hawthorne	Sweet gum
Hazel	Whitebrush
Hickory	Whitehorn
Huisache	Wild plum
Juniper	Willow
Locust	Yellow poplar
Lotebush	

- Partial control.

SPRAY EQUIPMENT AND APPLICATION TECHNIQUES

Basal (Soil)-Mix 2 2/3 pounds of VELPAR® DF with sufficient water to make one gallon of suspension and thoroughly agitate. Apply the VELPAR® DF suspension with an exact-delivery handgun applicator. This equipment delivers a thin stream of predetermined volume when triggered. Apply the VELPAR® DF suspension at the rate of 2 to 4 ml for each inch of stem diameter at breast height. Direct the treatment to soil within 3 inches of the root collar of woody plants to be controlled. When treating large stems and when more than one delivery of the VELPAR® DF suspension is needed per stem, make applications on opposite sides of the stem. Do not apply more

than 1/3 gallon of the VELPAR® DF suspension per acre per year. Intermittent agitation may be required to maintain the VELPAR® DF in suspension.

USE PRECAUTIONS

- PASTURE /RANGELAND

- Injury to or loss of desirable trees or other plants may result if VELPAR® DF is applied or if equipment is drained or flushed on or near desirable trees or other plants, on areas where their roots may extend, or in locations where the chemical may be washed or moved into contact with their roots.
- Poor weed and brush control may result from the following:
 - Use on poorly drained sites
 - Applications made when the soil is saturated with water and rain is imminent within 24 hours
 - Applications to soils high in organic matter (greater than 5%)
- Following mechanical cutting or clearing, allow stumps and injured trees sufficient time to adequately resprout before applying VELPAR® DF.
- Do not use VELPAR® DF on frozen soils.
- Weed and brush control results depend on sufficient moisture to activate VELPAR® DF.
- When VELPAR® DF is applied as a basal soil treatment, there is no restriction on grazing by domestic animals nor on cutting surrounding vegetation for forage or hay.
- For broadcast pasture applications of VELPAR® DF, do not cut treated vegetation for forage or hay nor graze domestic animals on treated areas for 60 days.

ADDITIONAL USE INFORMATION

SPRAY DRIFT MANAGEMENT

The interaction of many equipment- and weather-related factors determines the potential for spray drift. The applicator is responsible for considering all these factors when making application decisions. Avoiding spray drift is the responsibility of the applicator.

IMPORTANCE OF DROPLET SIZE

The most effective way to reduce drift potential is to apply large droplets (greater than 150-200 microns). The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. The presence of sensitive species nearby, the environmental conditions, and pest pressure may affect how an applicator balances drift control and coverage. Applying larger droplets reduces drift potential, but will not prevent drift if applications are made improperly or under unfavorable environmental conditions! See the Wind, Temperature and Humidity, and Temperature Inversions sections below.

CONTROLLING DROPLET SIZE

- GENERAL TECHNIQUES

- Volume - Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets.
- Pressure - Use the lower spray pressures recommended for the nozzle. Higher pressure reduces droplet size and does not improve canopy penetration. When higher flow rates are

needed, use a higher-capacity nozzle instead of increasing pressure.

- Nozzle Type - Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles.

CONTROLLING DROPLET SIZE - AIRCRAFT

- Number of Nozzles - Use the minimum number of nozzles with the highest flow rate that provide uniform coverage.
- Nozzle Orientation - Orienting nozzles so that the spray is emitted backwards, parallel to the airstream will produce larger droplets than other orientations.
- Nozzle Type - Solid stream nozzles (such as disc and core with swirl plate removed) oriented straight back produce larger droplets than other nozzle types.

BOOM LENGTH AND HEIGHT

- **Boom Length (aircraft)** - The boom length should not exceed 3/4 of the wing length, using shorter booms decreases drift potential. For helicopters use a boom length and position that prevents droplets from entering the rotor vortices.
- **Boom Height (aircraft)** - Application more than 10 feet above the canopy increases the potential for spray drift.
- **Boom Height (ground)** Setting the boom at the lowest labeled height (if specified) which provides uniform coverage reduces the exposure of droplets to evaporation and wind. The boom should remain level with the crop and have minimal bounce.

WIND

Drift potential increases at wind speeds of less than 3 mph (due to variable direction and inversion potential) or more than 10 mph. However, many factors, including droplet size and equipment type determine drift potential at any given wind speed. AVOID GUSTY OR WINDLESS CONDITIONS.

Note: Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they effect spray drift.

TEMPERATURE AND HUMIDITY

When making applications in hot and dry conditions, set up equipment to produce larger droplets to reduce effects of evaporation.

SURFACE TEMPERATURE INVERSIONS

Drift potential is high during a surface temperature inversion. Surface inversions restrict vertical air mixing, which causes small suspended droplets to remain close to the ground and move laterally in a concentrated cloud. Surface inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates a surface inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

SHIELDED SPRAYERS

Shielding the boom or individual nozzles can reduce the effects of wind. However, it is the responsibility of the applicator to verify that the shields are preventing drift, and not interfering with uniform deposition of the product.

SPRAY TANK CLEAN OUT

Thoroughly clean all traces of DuPont™ VELPAR® DF from application equipment immediately after use. Flush the tank, pump, hoses, and boom with several changes of water after removing nozzle tips and screens (clean these parts separately). Dispose of the equipment wash water by applying it to a use-site listed on this label.

STORAGE AND DISPOSAL

Storage: Store product in original container only. Do not contaminate water, other pesticides, fertilizer, food or feed in storage. Store in a cool, dry place.

Product Disposal: Do not contaminate water, food, or feed by disposal. Waste resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: For Plastic Containers: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke. **For Fiber Sacks:** Completely empty fiber sack by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into manufacturing or application equipment. Then dispose of sack in a sanitary landfill or by incineration if allowed by State and local authorities. **For Fiber Drums With Liners:** Completely empty liner by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application equipment. Then dispose of liner in a sanitary landfill or by incineration if allowed by State and local authorities. If drum is contaminated and cannot be reused, dispose of in the same manner. **For Paper and Plastic Bags:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

NOTICE TO BUYER: Purchase of this material does not confer any rights under patents of countries outside of the United States.

The DuPont Oval, DuPont™ and VELPAR® are trademarks or registered trademarks of E. I. duPont de Nemours and Company

“Microcel E” is a trademark of Johns Manville Product Corporation

“HiSil 233” is a trademark of Pittsburg Plate Glass

“Tordon 101” is a registered trademark of Dow AgroSciences

“Direx”, “Karmex” are trademarks of Griffin LLC

D - 856[MASTER] 121302

**LIMITATION OF
WARRANTY AND LIABILITY**

NOTICE: Read This Limitation of Warranty and Liability Before Buying or Using This Product. If the Terms Are Not Acceptable, Return the Product at Once, Unopened, and the Purchase Price Will Be Refunded.

It is impossible to eliminate all risks associated with the use of this product. Such risks arise from weather conditions, soil factors, off target movement, unconventional farming techniques, presence of other materials, the manner of use or application, or other unknown factors, all of which are beyond the control of DuPont. These risks can cause: ineffectiveness of the product; crop injury, or; injury to non-target crops or plants.

DuPont does not agree to be an insurer of these risks. **WHEN YOU BUY OR USE THIS PRODUCT, YOU AGREE TO ACCEPT THESE RISKS.**

DuPont warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for the purpose stated in the Directions for Use, subject to the inherent risks described above, when used in accordance with the Directions for Use under normal conditions.

DUPONT MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR OF MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

IN NO EVENT SHALL DUPONT OR SELLER BE LIABLE FOR ANY INCIDENTAL, CONSEQUENTIAL OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. BUYER'S OR USER'S BARGAINED-FOR EXPECTATION IS CROP PROTECTION. THE EXCLUSIVE REMEDY OF THE USER OR BUYER AND THE EXCLUSIVE LIABILITY OF DUPONT OR SELLER, FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY OR CONTRACT, NEGLIGENCE, TORT OR STRICT LIABILITY), WHETHER FROM FAILURE TO PERFORM OR INJURY TO CROPS OR OTHER PLANTS, AND RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT, OR AT THE ELECTION OF DUPONT OR SELLER, THE REPLACEMENT OF THE PRODUCT.

DuPont or its Ag Retailer must have prompt notice of any claim so that an immediate inspection of buyer's or user's growing crops can be made. Buyer and all users shall promptly notify DuPont or a DuPont Ag Retailer of any claims, whether based on contract, negligence, strict liability, other tort or otherwise or be barred from any remedy.

This Limitation of Warranty and Liability may not be amended by any oral or written agreement.

For product information call: 1-888-6-DUPONT

Internet address: <http://cropprotection.dupont.com/>

© 1998-2002 E. I. du Pont de Nemours and Company, Wilmington, Delaware 19898.

All rights reserved.



DuPont Agricultural Products

"..... A Growing Partnership With Nature"



SUPPLEMENTAL LABELING

DUPONT™ VELPAR® DF HERBICIDE HIGHBUSH BLUEBERRIES

DUPONT™ VELPAR® DF HERBICIDE

(EPA REG. NO. 352-581)

FOR USE ON HIGHBUSH BLUEBERRIES

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

DuPont™ VELPAR® DF Herbicide is recommended for control of certain herbaceous and woody weeds in established highbush blueberry fields.

HOW TO USE

Apply 1.3 to 2.6 pounds VELPAR® DF per acre (broadcast; for band treatment use proportionately less) to pruned blueberry fields in the spring before blueberry leaf emergence. VELPAR® DF spray should be directed to the soil and weeds. When applied properly, VELPAR® DF controls many herbaceous and woody plants including: barnyard-grass, briars* (blackberry, raspberry), wild cherry, goldenrod, hawkweed, quackgrass, sheep sorrel, red sorrel, witch grass, yellow foxtail broomsedge*, dog fennel, fall panicum, and panic grass. Using properly calibrated ground equipment, make application in sufficient water to provide thorough and uniform coverage of the treated area, (usually 20 gallons per acre). Spray booms must be shut off while starting, turning, slowing or stopping, as injury to the crop may result.

*Suppression or partial control only.

Do not apply within 90 days of harvest.

USE PRECAUTIONS

Since the effect of VELPAR® DF on blueberries varies with soils, plant vigor, uniformity of application, and amount of rainfall, it is suggested that growers limit their first use to small areas. Most blueberries are resistant to VELPAR® DF, but some clones are susceptible to injury. Application to blueberry plants after leaf emergence will result in injury. Use lower rates on poorly drained soils, but do not use if there is standing water. Do not use on sands, loamy sands or sandy loam soils. Do not apply to frozen or snow covered ground.

Treat only plantings established for 3 years or more.

Do not apply this product through any type of irrigation system.

IMPORTANT

BEFORE USING VELPAR® DF HERBICIDE, READ AND CAREFULLY OBSERVE THE CAUTIONARY STATEMENTS AND ALL OTHER INFORMATION ON ITS PACKAGE LABEL.

This labeling contains new and supplemental instructions for use of DuPont VELPAR® DF Herbicide.

This labeling must be in the immediate possession of the applicator at the time of application.

R-180 013102 03-11-02



**DuPont Agricultural
Products**

"..... A Growing Partnership With Nature"



SUPPLEMENTAL LABELING

**DUPONT™ VELPAR® DF
HERBICIDE
CHEMIGATION
USE ON ALFALFA**

DUPONT™ VELPAR® DF HERBICIDE

EPA Reg. No. 352-581

FOR USE ON ALFALFA APPLIED VIA CHEMIGATION

DuPont™ VELPAR® DF herbicide can be applied using chemigation to alfalfa. Make a single application during the winter months when alfalfa plants are in the least active stage of growth and/or to stubble after cutting following hay removal and before regrowth exceeds 2" in height.

Severe alfalfa injury may result following application after cutting if either the regrowth is more than 2" high or significant stubble is left after alfalfa cutting.

DIRECTIONS FOR USE

Dormant Applications

Follow the EPA-registered package label for rate instructions and weeds controlled when applying VELPAR® DF Herbicide by chemigation during winter months when alfalfa plants are in the least active stage of growth. Select the appropriate dose for soil texture and organic matter content using 0.25" to 0.75" of sprinkler irrigation as a continuous injection during the application. Best results are obtained when soil is moist at time of application, and when weeds have not germinated or are less than 2" tall or across.

Application After Cutting

Apply VELPAR® DF Herbicide at 1/3 lb per acre to stubble after cutting, following hay removal, and before regrowth exceeds 2" in height. Apply VELPAR® DF Herbicide using 0.25" to 0.75" of sprinkler irrigation as a continuous injection during the application. Best results are obtained when soil is moist at time of application and when weeds have not germinated or are less than 2" tall or across.

Refer to the EPA-registered package label for information regarding use restrictions, rotational cropping recommendations, sprayer cleanup, use precautions and other information. It is a violation of federal law to use this product in a manner inconsistent with its labeling.

CHEMIGATION

Apply this product only through center pivot sprinkler irrigation systems. Do not apply this product through any other type of irrigation system.

Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water.

If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts.

Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label prescribed safety devices for public water systems are in place.

A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

SPRINKLER CHEMIGATION

The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from back-flow.

The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.

The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.

The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.

23/26

Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock. Do not apply when wind speed favors drift beyond the area intended for treatment.

MIXING INSTRUCTIONS

1. Fill the supply tank 1/4 to 1/3 full of water.
2. While agitating, add the required amount of VELPAR® DF and continue agitation until the VELPAR® DF is fully dispersed, at least 5 minutes.
3. Once the VELPAR® DF is fully dispersed, maintain agitation and continue filling tank with water. VELPAR® DF should be thoroughly mixed with water before adding any other material.
4. As the tank is filling, add tank mix partners (if desired).
5. After thorough mixing, the agitation system can be stopped to prevent excessive foaming in the tank. Once thoroughly mixed the solution in the supply tank does not require additional agitation unless specified on the companion products label. Read and follow all manufacturers label recommendations for the companion products. A de-foamer can be added to the supply tank if companion products require continuous agitation in the injection supply tank during application.
6. Apply VELPAR® DF spray mixture within 48 hours of mixing to avoid product degradation.

PRECAUTIONS

- Distributing treated water in an uneven manner can result in crop injury, lack of effectiveness, or over-tolerance pesticide residues in the crop. Therefore, to ensure that the mixture is applied evenly at the recommended rate, use sufficient water, apply the mixture for the proper length of time and ensure sprinkler produces a uniform water pattern.
- On soils high in organic matter (greater than 5%), the effectiveness of VELPAR® DF can be significantly reduced and weed control may be unsatisfactory.
- Since the effect of VELPAR® DF on alfalfa varies with soil conditions, uniformity of application, and environmental conditions, growers should limit their first use to small area.
- Temporary yellowing of alfalfa may occur following VELPAR® DF applications.
- Crop injury, including mortality, may result in fields with restricted root growth due to non-uniform soil profiles such as gravel bases and clay lenses.
- Crop injury may result if extremely hot weather (around 100° F or higher) occurs within a few days after application.

RESTRICTIONS

- Do not permit run-off during chemigation.
- Do not apply when wind speed favors drift beyond the area intended for treatment.
- Do not connect an irrigation system used for DuPont VELPAR® DF application to a public water system.

- Treat only stands of alfalfa established for two seasons or more (greater than 13 months) provided the alfalfa root system is well established and the crop is healthy, vigorous, and not under stress from weather conditions, insects, or disease damage.
- In the PNW region, treat only stands that have a well developed tap root structure that is at least 10 inches in length throughout the field and the crop is healthy, vigorous, and not under stress from weather conditions, low fertility, insects or disease damage.
- Do not use on seedling alfalfa, alfalfa/grass mixtures, or other mixed stands.
- Do not apply to snow-covered or frozen ground.
- Do not add a surfactant to VELPAR® DF when treating non-dormant alfalfa varieties via chemigation.
- Do not use VELPAR® DF on gravelly or rocky soils, exposed subsoils, hardpan, sand, poorly drained soil, or alkali soils.
- Do not graze or feed forage or hay to livestock within 30 days following application.

POSTING OF AREAS TO BE TREATED

Posting of areas to be chemigated is required when 1) any part of a treated area is within 300 feet of sensitive areas such as residential areas, labor camps, businesses, daycare centers, hospitals, in-patient clinics, nursing homes, or any public areas such as schools, parks, playgrounds, or other public facilities not including public roads, or 2) when the chemigated area is open to the public such as golf courses or retail greenhouses.

Posting must conform to all the following requirements. Treated areas shall be posted with signs at all usual points of entry and along likely routes of approach from the listed sensitive areas. When there are no usual points of entry, signs must be posted in the corners of the treated areas and in any other location affording maximum visibility to sensitive areas. The printed side of the sign should face away from the treated area towards the sensitive area. The signs shall be printed in English. Signs must be posted prior to application and must remain posted until foliage has dried and soil surface water has disappeared. Signs may remain in place indefinitely as long as they are composed of materials to prevent deterioration and maintain legibility for the duration of the posting period.

All words shall consist of letters at least 2 1/2 inches tall, and all letters and the symbol shall be a color which sharply contrasts with their immediate background. At the top of the sign shall be the words "KEEP OUT", followed by an octagonal stop sign symbol at least 8 inches in diameter containing the word "STOP". Below the symbol shall be the words "PESTICIDE IN IRRIGATION WATER".

Posting required for chemigation does not replace other posting and reentry requirements for farm worker safety.

**IMPORTANT
BEFORE USING THESE PRODUCTS, READ
AND FOLLOW ALL APPLICABLE DIRECTIONS,
RESTRICTIONS AND PRECAUTIONS
ON THE EPA-REGISTERED LABEL.**


This bulletin contains new or supplemental instructions for use of this product which do not appear on the EPA-registered package label. Follow the instructions carefully.

This labeling must be in the possession of the user at the time of pesticide application.

DR-259 121602



DuPont Crop Protection



SUPPLEMENTAL LABELING

DUPONT™ VELPAR® DF HERBICIDE LOWBUSH BLUEBERRIES

DUPONT™ VELPAR® DF HERBICIDE

(EPA REG. NO. 352-581)

FOR USE ON LOWBUSH BLUEBERRIES

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

DuPont™ VELPAR® DF Herbicide is recommended for control of certain herbaceous and woody weeds in established lowbush blueberry fields.

HOW TO USE

Applied in the spring before blueberry leaf emergence VELPAR® DF controls:

* Considered hard to kill - control may not be adequate.

0.6 to 1.2 lbs/Acre

Certain grasses
Certain flowering herbaceous weeds

Aster
Cinquefoil
Fireweed
Goldenrod
Hawkweeds
Pearly everlasting
Sheep sorrel
Wild strawberry
Yarrow

2.4 to 3.6 lbs/Acre(1)

1.2 to 2.4 lbs/Acre

Dogbane*

Meadowsweet
Trailing blackberry
Sheep-laurel
Wild rose*

(1) Do not exceed 2.4 lbs/Acre if soil has been treated with hexazinone within the past 8 years.

USE PRECAUTIONS

VELPAR® DF may only be applied to pruned blueberry fields in the spring before emergence of blueberry foliage.

Most lowbush blueberries are resistant to VELPAR® DF but some clones are susceptible to injury. The phytotoxic effect of VELPAR® DF to these susceptible blueberry clones may vary with soil type, texture, uniformity of application, and amount of rainfall.

If excessive leaf drop is observed after treatment, reduce rate in future applications.

Use lower rates on poorly drained soils.

Do not use if there is standing water.

Do not apply within 450 days of harvest.

Do not apply through any type of irrigation system.

Maintain a 50 foot buffer from any well head or water reservoir.

IMPORTANT

BEFORE USING VELPAR® DF HERBICIDE, READ AND FOLLOW ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON ITS PACKAGE LABEL.

This labelling contains new and supplemental instructions for use of "VELPAR® DF" Herbicide.

This labeling must be in the immediate possession of the applicator at the time of application.

(Replaces H-63027)

R-239 081402 10-10-96