

Please read instructions on reverse before completing form.

Form Approved. OMB No. 2070-0060. Approval expires 05-31-98

	United States <b>Environmental Protection Agency</b> Washington, DC 20460	<input type="checkbox"/> Registration <input type="checkbox"/> Amendment <input checked="" type="checkbox"/> Other	OPP Identifier Number
--	---	--	-----------------------

**Application for Pesticide - Section I**

1. Company/Product Number 352-597	2. EPA Product Manager Geraldine R. McCann	3. Proposed Classification <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) DuPont™ Avaunt® Insecticide	PM#	
5. Name and Address of Applicant (Include ZIP Code) E. I. du Pont de Nemours & Company Crop Protection, P. O. Box 30 Newark, DE 19714-0030 Attention: Rebecca M. Ashley  <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____	

**Section - II**

<input type="checkbox"/> Amendment - Explain below.	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____	<b>NOTIFICATION</b>  <b>MAR 12 2002</b>
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application.	
<input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Other - Explain below.	

**Explanation:** Use additional page(s) if necessary. (For section I and Section II.)

- \* One copy of letter to Geri McCann
- \* Three copies of labeling changes submitted via notification for DuPont™ Avaunt® Insecticide (SL-820-1 022602 01-11-02) with changes highlighted on one copy (addition of "Not for use or sale in Suffolk and Nassau Counties, Long Island, New York")
- \* One copy of currently approved DuPont™ Avaunt® Insecticide label for reference (H-64181) (SL-820-1 102501 01-11-02)

**Section - III**

1. Material This Product Will Be Packaged In:				2. Type of Container	
Child-Resistant Packaging <input type="checkbox"/> Yes* <input type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No		<input type="checkbox"/> Metal	<input type="checkbox"/> Plastic
* Certification must be submitted		If "Yes" Unit Packaging wgt. No. per container	If "Yes" Package wgt. No. per container	<input type="checkbox"/> Glass	<input type="checkbox"/> Paper
				<input type="checkbox"/> Other (Specify) _____	
3. Location of Net Contents Information <input type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) Retail Container		5. Location of Label Directions <input type="checkbox"/> On Label <input type="checkbox"/> On Labeling accompanying product	
6. Manner in Which Label is Affixed to Product		<input type="checkbox"/> Lithograph <input type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled		<input type="checkbox"/> Other _____	

**Section - IV**

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)		
Name Rebecca M. Ashley	Title Registration Coordinator	Telephone No. (include Area Code) (302) 451-0629
<b>Certification</b> I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment as both under applicable law.		6. Date-Application Received (Stamped)
2. Signature <i>Rebecca M Ashley</i>	3. Title Registration Coordinator	
4. Typed Name Rebecca M. Ashley	5. Date 2/27/02	

H-2/10




**NOTIFICATION**

MAR 12 2002

# DuPont™ Avaunt®

**insecticide**

---



*“..... A Growing Partnership With Nature”*



# DuPont™ Avaunt®

## insecticide

### Dispersible Granules

<i>Active Ingredient</i>	<i>By Weight</i>
Indoxacarb (S)-methyl 7-chloro-2,5-dihydro-2-[[methoxy-carbonyl][4(trifluoromethoxy)phenyl]amino]-carbonyl]indeno[1,2-e][1,3,4]oxadiazine-4a-(3H)-carboxylate	30%
<i>Inert Ingredients</i>	70%
<b>TOTAL</b>	<b>100%</b>

EPA Reg. No. 352-597

Net Contents: \_\_\_\_\_

Establishment No.: 33972-FR-1

**PRECAUTIONARY STATEMENTS**  
**KEEP OUT OF REACH OF CHILDREN**  
**CAUTION**

#### FIRST AID

**IF SWALLOWED:** Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.

**IF IN EYES:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

**IF ON SKIN OR CLOTHING:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

**IF INHALED:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for treatment advice. Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-441-3637 for emergency medical treatment information.

### PRECAUTIONARY STATEMENTS (cont'd)

#### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

**Caution!** Harmful if swallowed. Causes moderate eye irritation. Harmful if absorbed through skin. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling. Harmful if inhaled. Avoid breathing (dust, vapor or spray mist). Remove contaminated clothing and wash clothing before reuse.

#### PERSONAL PROTECTIVE EQUIPMENT

Some materials that are chemical resistant to this product are listed below. If you want more options follow the instructions for Category A on the EPA chemical resistance category selection chart.

**Applicators and other handlers must wear:**

- Long-sleeved shirt and long pants.
- Chemical Resistant Gloves Category A (such as butyl rubber, natural rubber, neoprene rubber or nitrile rubber), all  $\geq 14$  mls.
- Shoes plus socks.

Follow manufacturer's instructions for cleaning/maintaining personal protective equipment (PPE). If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

#### USER SAFETY RECOMMENDATIONS

**USERS SHOULD:** Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

#### ENVIRONMENTAL HAZARDS

This pesticide is toxic to mammals, birds, fish and aquatic invertebrates. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment rinsewater. Do not apply where/when conditions could favor runoff. Runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Cover, incorporate, or clean up granules that are spilled.

This product is highly toxic to bees exposed to direct treatment on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds while bees are actively visiting the treatment area.

## DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

For early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, wear:

- Coveralls over long sleeved shirt and long pants,
- Socks plus chemical resistant footwear,
- Chemical Resistant Gloves Category A (such as butyl rubber, natural rubber, neoprene rubber or nitrile rubber), all  $\geq 14$  mls.

DuPont™ AVAUNT® should be used only in accordance with recommendations on this label or in separate DuPont supplemental labeling available through local dealers.

DuPont will not be responsible for losses or damages resulting from use of this product in any manner not specifically recommended by DuPont. User assumes all risks associated with such non-recommended use.

### GENERAL INFORMATION

AVAUNT® is a water dispersible granule that can be applied as a foliar spray to control many important insects. AVAUNT® is mixed with water for application.

Use only in commercial and farm plantings. Not for use in home plantings. Not for use during any period after a commercial crop site is opened for public entry as a "U-Pick", "Pick Your Own" or similar operation; in no case shall preharvest applications be made after first public entry. (The restricted entry interval(s) stated elsewhere on this label must be followed.

Not for use or sale in Suffolk and Nassau Counties, Long Island, New York.

Do not formulate this product into other End-use products without written permission.

Do not apply this product through any type of irrigation system.

### INTEGRATED PEST MANAGEMENT

This product should be used as part of an Integrated Pest Management (IPM) program which can include biological, cultural, and genetic practices aimed at preventing economic pest damage. Application of this product should be based on IPM principles and practices including field scouting or other detection methods, correct target pest identification, population monitoring, and treating when target pest populations reach locally determined action thresholds. Consult your state cooperative extension service, professional consultants or other qualified authorities to determine appropriate action threshold levels for treating specific pest/crop systems in your area.

### BENEFICIAL ARTHROPODS

AVAUNT® helps conserve certain beneficials (parasites and predators). While these beneficials cannot be relied upon to control pests, they are of potential value and should be monitored along with pests in pest management programs on these crops.

### SCOUTING

Monitor insect populations to determine whether or not there is a need for application of AVAUNT® based on locally determined economic thresholds. More than one treatment of AVAUNT® may be required to control a population of pests.

### RESISTANCE

Some insects are known to develop resistance to products used repeatedly for control. When this occurs, the recommended dosages fail to suppress the pest population below the economic threshold. Because the development of resistance cannot be predicted, the use of this product should conform to resistance management strategies established for the use area. These strategies may include incorporation of cultural and biological control practices, alternation of active classes of insecticides on succeeding generations and targeting the most susceptible life stage. Consult your local agricultural authorities for details.

### APPLICATION

Apply at the recommended rates when insect populations reach locally determined economic thresholds. Consult the cooperative extension service, professional consultants or other qualified authorities to determine appropriate threshold levels for treatment in your area.

Follow-up treatments of AVAUNT® should be applied, as needed, to keep pest populations within threshold limits. On most crops, AVAUNT® should be applied at 5 to 7 day intervals to maintain control. On apple and pear the minimum interval between treatments is 7 days.

Use sufficient water to obtain thorough, uniform coverage.

For aerial application, use a minimum of 5 gals. water per acre (gpa), unless otherwise specified.

Use of Adjuvants: In some situations where coverage is difficult to achieve such as closed canopy, dense foliage, or less than optimum application equipment, an adjuvant may improve performance.

Crops	Insects	DuPont™ AVAUNT® Rate Per Acre		Last Application Days to Harvest	REI
		Lbs. A.I.	Ounces		
Apple	Codling moth*	0.09 - 0.11	5.0 - 6.0	28	12 hrs.
	Oriental fruit moth Lesser appleworm Pandemis leafroller Redbanded leafroller Tufted apple bud moth White apple leafhopper Potato leafhopper Plum curculio Tarnished plant bug Apple maggot Lacanobia fruitworm (except California) European apple sawfly (except California)	0.09 - 0.11	5.0 - 6.0		
<p>Make no more than 3 applications prior to hand-thinning. No hand-thinning after the 4th application.            Make no more than 4 applications per season. Do not apply more than 24 oz. AVAUNT® (0.44 lbs a.i.)            per acre per crop. Do not apply dilute applications of more than 200 gal water per acre. For best results            apply 50 - 150 gal water per acre. The minimum interval between treatments is 7 days.            *West of the Rockies. For use against low to moderate infestations in conjunction with alternate control            measures such as established Mating Disruption blocks.</p>					
Broccoli	Beet Armyworm** Diamondback moth	0.065*	3.5*	3	12 hrs
	Cabbage looper Imported cabbageworm	0.045 - 0.065*	2.5 - 3.5*		
<p>Do not apply more than 14 ounces AVAUNT® (0.26 lbs a.i.) per acre per crop.            The minimum interval between sprays is 3 days.            *Add a wetting agent to improve spray coverage.            **For control of low numbers and not for corrective treatments of higher numbers of larvae.</p>					
Brussels Sprouts	Beet Armyworm** Diamondback moth	0.065*	3.5*	3	12 hrs
	Cabbage looper Imported cabbageworm	0.045 - 0.065*	2.5 - 3.5*		
<p>Do not apply more than 14 ounces AVAUNT® (0.26 lbs a.i.) per acre per crop.            The minimum interval between sprays is 3 days.            *Add a wetting agent to improve spray coverage.            **For control of low numbers and not for corrective treatments of higher numbers of larvae.</p>					
Cabbage (tight headed varieties only)	Beet Armyworm** Diamondback moth	0.065*	3.5*	3	12 hrs
	Cabbage looper Imported cabbageworm	0.045 - 0.065*	2.5 - 3.5*		
<p>Do not apply more than 14 ounces AVAUNT® (0.26 lbs a.i.) per acre per crop.            The minimum interval between sprays is 3 days.            *Add a wetting agent to improve spray coverage.            **For control of low numbers and not for corrective treatments of higher numbers of larvae.</p>					
Cauliflower	Beet Armyworm** Diamondback moth	0.065*	3.5*	3	12 hrs
	Cabbage looper Imported cabbageworm	0.045 - 0.065*	2.5 - 3.5*		
<p>Do not apply more than 14 ounces AVAUNT® (0.26 lbs a.i.) per acre per crop.            The minimum interval between sprays is 3 days.            *Add a wetting agent to improve spray coverage.            **For control of low numbers and not for corrective treatments of higher numbers of larvae.</p>					
Chinese Broccoli	Beet Armyworm** Diamondback moth	0.065*	3.5*	3	12 hrs
	Cabbage looper Imported cabbageworm	0.045 - 0.065*	2.5 - 3.5*		
<p>Do not apply more than 14 ounces AVAUNT® (0.26 lbs a.i.) per acre per crop.            The minimum interval between sprays is 3 days.            *Add a wetting agent to improve spray coverage.            **For control of low numbers and not for corrective treatments of higher numbers of larvae.</p>					
Chinese Cabbage (napa)	Beet Armyworm** Diamondback moth	0.065*	3.5*	3	12 hrs
	Cabbage looper Imported cabbageworm	0.045 - 0.065*	2.5 - 3.5*		
<p>Do not apply more than 14 ounces AVAUNT® (0.26 lbs a.i.) per acre per crop.            The minimum interval between sprays is 3 days.            *Add a wetting agent to improve spray coverage.            **For control of low numbers and not for corrective treatments of higher numbers of larvae.</p>					

Crops	Insects	DuPont™ AVAUNT® Rate Per Acre		Last Application Days to Harvest	REI
		Lbs. A.I.	Ounces		
Chinese Mustard Cabbage	Beet Armyworm** Diamondback moth	0.065*	3.5*	3	12 hrs
	Cabbage looper Imported cabbageworm	0.045 - 0.065*	2.5 - 3.5*		
	Do not apply more than 14 ounces AVAUNT® (0.26 lbs a.i.) per acre per crop. The minimum interval between sprays is 3 days. *Add a wetting agent to improve spray coverage. **For control of low numbers and not for corrective treatments of higher numbers of larvae.				
Corn (sweet)	Fall armyworm European corn borer (except California)	0.045 - 0.065	2.5 - 3.5	3 35 - fodder & stover	12 hrs
	Make no more than 4 applications per season. Whorl application only. Do not apply more than 14 ounces AVAUNT® (0.26 lbs a.i.) per acre per crop. The minimum interval between sprays is 3 days.				
Eggplant	Tomato fruitworm Beet armyworm Southern armyworm	0.065	3.5	3	12 hrs
	Loopers	0.045 - 0.065	2.5 - 3.5		
	Do not apply more than 14 ounces AVAUNT® (0.26 lbs a.i.) per acre per crop. The minimum interval between sprays is 5 days.				
Kohlrabi	Beet Armyworm** Diamondback moth	0.065*	3.5*	3	12 hrs
	Cabbage looper Imported cabbageworm	0.045 - 0.065*	2.5 - 3.5*		
	Do not apply more than 14 ounces AVAUNT® (0.26 lbs a.i.) per acre per crop. The minimum interval between sprays is 3 days. *Add a wetting agent to improve spray coverage. **For control of low numbers and not for corrective treatments of higher numbers of larvae.				
Lettuce (Head and Leaf varieties)	Beet armyworm* Corn earworm	0.065	3.5	3	12 hrs
	Cabbage looper	0.045 - 0.065	2.5 - 3.5		
	Do not apply more than 14 ounces AVAUNT® (0.26 lbs a.i.) per acre per crop. The minimum interval between sprays is 3 days. *For control of low numbers and not for corrective treatments of higher numbers of larvae.				
Pear	Codling moth*	0.09 - 0.11	5.0 - 6.0	28	12 hrs.
	Oriental fruit moth Redbanded leafroller	0.09 - 0.11	5.0 - 6.0		
	Make no more than 3 applications prior to hand-thinning. No hand-thinning after the 4th application. Make no more than 4 applications per season. Do not apply more than 24 oz AVAUNT® (0.44 lbs a.i.) per acre per crop. Do not apply dilute applications of more than 200 gal water per acre. For best results apply 50 - 150 gal water per acre. The minimum interval between treatments is 7 days. *West of the Rockies. For use against low to moderate infestations in conjunction with alternate control measures such as established Mating Disruption blocks.				
Pepper (Bell and Non-Bell)	Tomato fruitworm Beet armyworm Southern armyworm	0.065	3.5	3	12 hrs
	Loopers	0.045 - 0.065	2.5 - 3.5		
	Do not apply more than 14 ounces AVAUNT® (0.26 lbs a.i.) per acre per crop. The minimum interval between sprays is 5 days.				
Tomato	Tomato fruitworm Beet armyworm Tomato pinworm Southern armyworm	0.065	3.5	3	12 hrs
	Loopers Hornworms	0.045 - 0.065	2.5 - 3.5		
	Do not apply more than 14 ounces AVAUNT® (0.26 lbs a.i.) per acre per crop. The minimum interval between sprays is 5 days.				

## SPRAY PREPARATION

Spray equipment must be clean and free of previous pesticide deposits before applying DuPont™ AVAUNT®. Fill spray tank 1/4 to 1/2 full of water. Add AVAUNT® directly to spray tank. Mix thoroughly to fully disperse the insecticide; once dispersed continued agitation is required. Use mechanical or hydraulic means; do not use air agitation. Spray mix should not be stored overnight in spray tank.

**Compatibility** - Since formulations may be changed and new ones introduced, it is recommended that users premix a small quantity of a desired tank mix and observe for possible adverse changes (settling out, flocculation, etc.). Avoid mixtures of several materials and very concentrated spray mixtures.

This product can be mixed with pesticide products labeled for use on crops on this label in accordance with the most restrictive of label limitations and precautions. No label dosage rates should be exceeded. This product cannot be mixed with any product containing a label prohibition against such mixing.

**Tank Mixing Sequence** - Add different formulation types in the sequence indicated below. Allow time for complete mixing and dispersion after addition of each product.

1. Water soluble bags.
2. AVAUNT® and other water dispersible granules.
3. Wettable powders.
4. Water based suspension concentrates.
5. Water soluble concentrates.
6. Oil based suspension concentrates.
7. Emulsifiable concentrates.
8. Adjuvants, surfactants, oils.
9. Soluble fertilizers.
10. Drift retardants.

## SPRAY DRIFT MANAGEMENT

The interaction of many equipment and weather-related factors determines the potential for spray drift. The applicator is responsible for considering all these factors when making application decisions.

**AVOIDING SPRAY DRIFT IS THE RESPONSIBILITY OF THE APPLICATOR.**

### IMPORTANCE OF DROPLET SIZE

The most effective way to reduce drift potential is to apply large droplets (>150 microns). The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. The presence of sensitive species nearby, the environmental conditions, and pest pressure may affect how an applicator balances drift control and coverage. **APPLYING LARGER DROPLETS REDUCES DRIFT POTENTIAL, BUT WILL NOT PREVENT DRIFT IF APPLICATIONS ARE MADE IMPROPERLY OR UNDER UNFAVORABLE ENVIRONMENTAL CONDITIONS! See Wind, Temperature and Humidity, and Surface Temperature Inversions sections of this label.**

### Controlling Droplet Size - General Techniques

- **Volume** - Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets.
- **Pressure** - Use the lower spray pressures recommended for the nozzle. Higher pressure reduces droplet size and does not improve canopy penetration. **WHEN HIGHER FLOW RATES ARE NEEDED, USE A HIGHER-CAPACITY NOZZLE INSTEAD OF INCREASING PRESSURE.**
- **Nozzle Type** - Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles.

### Controlling Droplet Size - Aircraft

- **Number of Nozzles** - Use the minimum number of nozzles with the highest flow rate that provide uniform coverage.
- **Nozzle Orientation** - Orienting nozzles so that the spray is emitted backwards, parallel to the airstream will produce larger droplets than other orientations.
- **Nozzle Type** - Solid stream nozzles (such as disc and core with swirl plate removed) oriented straight back produce larger droplets than other nozzle types.
- Do not apply as a ULV application.

### BOOM LENGTH AND HEIGHT

- **Boom Length (aircraft)** - The boom length should not exceed 3/4 of the wing length, using shorter booms decreases drift potential. For helicopters use a boom length and position that prevents droplets from entering the rotor vortices.
- **Boom Height (aircraft)** - Application more than 10 ft above the canopy increases the potential for spray drift.
- **Boom Height (ground)** Setting the boom at the lowest height which provides uniform coverage reduces the exposure of droplets to evaporation and wind. The boom should remain level with the crop and have minimal bounce.

### WIND

Drift potential increases at wind speeds of less than 3 mph (due to variable direction and inversion potential) or more than 10 mph. However, many factors, including droplet size and equipment type determine drift potential at any given wind speed. **AVOID APPLICATIONS DURING GUSTY OR WINDLESS CONDITIONS.**

**Note:** Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they effect spray drift.

### TEMPERATURE AND HUMIDITY

When making applications in hot and dry conditions, set up equipment to produce larger droplets to reduce effects of evaporation.

## **SURFACE TEMPERATURE INVERSIONS**

Drift potential is high during a surface temperature inversion. Surface inversions restrict vertical air mixing, which causes small suspended droplets to remain close to the ground and move laterally in a concentrated cloud. Surface inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates a surface inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

## **SHIELDED SPRAYERS**

Shielding the boom or individual nozzles can reduce the effects of wind. However, it is the responsibility of the applicator to verify that the shields are preventing drift and not interfering with uniform deposition of the product.

## **AIR ASSISTED (AIR BLAST) FIELD CROP SPRAYERS**

Air assisted field crop sprayers carry droplets to the target via a downward directed air stream. Some may reduce the potential for drift, but if a sprayer is unsuitable for the application and/or set up improperly, high drift potential can result. It is the responsibility of the applicator to determine that a sprayer is suitable for the intended application, is configured properly, and that drift is not occurring.

**Note:** Air assisted field sprayers can affect product performance by affecting spray coverage and canopy penetration. Consult the application equipment section of this label to determine if use of an air assisted sprayer is recommended.

## **AIR ASSISTED (AIR BLAST) TREE AND VINE SPRAYERS**

Air assisted tree and vine sprayers carry droplets into the canopy of trees and vines via a radially or laterally directed air stream.

In addition to the general drift management principles already described, the following specific practices will further reduce the potential for drift.:

- *Adjust deflectors and aiming devices so that spray is only directed into the canopy.*
- *Block off upward pointed nozzles when there is no overhanging canopy.*
- *Use only enough air volume to penetrate the canopy and provide good coverage.*
- *Do not allow spray to go beyond the edge of the cultivated area. Spray the outside row only from outside the planting.*

## **SPRAY TANK CLEANOUT**

Prior to application, start with clean, well maintained application equipment. Immediately following application, thoroughly clean all spray equipment to reduce the risk of forming hardened deposits which might become difficult to remove.

Drain spray equipment. Thoroughly rinse sprayer and flush hoses, boom and nozzles with clean water.

Clean all other associated application equipment. Take all necessary safety precautions when cleaning equipment. Do not clean near wells, water sources or desirable vegetation. Dispose of waste rinse water in accordance with local regulations.

## **CROP ROTATION RESTRICTIONS**

Crops that are on this label and cotton may be planted immediately following harvest. Do not plant for food or feed any other crops not registered for use with indoxacarb for 30 days after last use.

## **STORAGE AND DISPOSAL**

Do not contaminate water, food, or feed by storage or disposal.

**STORAGE:** Store product in original container only in a location inaccessible to children and pets. Do not contaminate water, other pesticides, fertilizer, food or feed in storage. Not for use or storage in or around the home.

**PESTICIDE DISPOSAL:** Do not contaminate water, food, or feed by storage or disposal. Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

### **CONTAINER DISPOSAL: For Plastic**

**Containers:** Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke. **For Fiber Drums With Liners:** Completely empty liner by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application equipment. Then dispose of liner in a sanitary landfill or by incineration if allowed by State and local authorities. If drum is contaminated and cannot be reused, dispose of in the same manner.

**NOTICE TO BUYER**--Purchase of this material does not confer any rights under patents of countries outside of the United States.

The DuPont Oval Logo, DuPont™ and AVAUNT® are trademarks or registered trademarks of E. I. duPont de Nemours and Company.

**SL - 820 - 1 022602 01-11-02**

**LIMITATION OF WARRANTY AND LIABILITY**

NOTICE: Read This Limitation of Warranty and Liability Before Buying or Using This Product. If the Terms Are Not Acceptable, Return the Product at Once, Unopened, and the Purchase Price Will Be Refunded.

It is impossible to eliminate all risks associated with the use of this product. Such risks arise from weather conditions, soil factors, off target movement, unconventional farming techniques, presence of other materials, the manner of use or application, or other unknown factors, all of which are beyond the control of DuPont. These risks can cause: ineffectiveness of the product; crop injury, or; injury to non-target crops or plants.

DuPont does not agree to be an insurer of these risks. WHEN YOU BUY OR USE THIS PRODUCT, YOU AGREE TO ACCEPT THESE RISKS.

DuPont warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for the purpose stated in the Directions for Use, subject to the inherent risks described above, when used in accordance with the Directions for Use under normal conditions.

DUPONT MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR OF MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

IN NO EVENT SHALL DUPONT OR SELLER BE LIABLE FOR ANY INCIDENTAL, CONSEQUENTIAL OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. BUYER'S OR USER'S BARGAINED-FOR EXPECTATION IS CROP PROTECTION. THE EXCLUSIVE REMEDY OF THE USER OR BUYER AND THE EXCLUSIVE LIABILITY OF DUPONT OR SELLER, FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY OR CONTRACT, NEGLIGENCE, TORT OR STRICT LIABILITY), WHETHER FROM FAILURE TO PERFORM OR INJURY TO CROPS OR OTHER PLANTS, AND RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT, OR AT THE ELECTION OF DUPONT OR SELLER, THE REPLACEMENT OF THE PRODUCT.

DuPont or its Ag Retailer must have prompt notice of any claim so that an immediate inspection of buyer's or user's growing crops can be made. Buyer and all users shall promptly notify DuPont or a DuPont Ag Retailer of any claims, whether based on contract, negligence, strict liability, other tort or otherwise or be barred from any remedy.

This Limitation of Warranty and Liability may not be amended by any oral or written agreement.

**For product information call: 1-888-6-DUPONT**

**Internet address: [www.dupont.com/ag/us](http://www.dupont.com/ag/us)**

**© 2000-2002 E. I. du Pont de Nemours and Company, Crop Protection, Wilmington, Delaware 19898.**

**All rights reserved.**

10/10



DuPont Crop Protection

February 27, 2002

Ms. Geri McCann  
Document Processing Desk (NOTIF)  
Office of Pesticide Programs (7504C)  
U.S. Environmental Protection Agency  
Ariel Rios Building  
1200 Pennsylvania Avenue, N.W.  
Washington, DC 20460

Subject: DuPont Avaunt Insecticide (EPA Reg. No. 352-597)  
Submission of Labeling Changes via Notification (SL-820-1 022602 01-11-02)

Dear Ms. McCann:

Enclosed is an Application for Pesticide Amendment Form submitting label changes via notification for DuPont Avaunt Insecticide (SL-820-1 022602 01-11-02). Three copies of the label are enclosed, one with the changes highlighted. I have also enclosed a copy of the currently approved label (H-64181) (SL-820-1 102501 01-11-02) for your reference.

The change to the label is the addition of the following text in the General Information Section:

“Not for use or sale in Suffolk and Nassau Counties, Long Island, New York.”

This notification is consistent with the provisions of PR Notice 95-2 and EPA regulations at 40 CFR 152.46, and no other changes have been made to the labeling or the confidential statements of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the terms of PR Notice 95-2 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.

Please feel free to call me at (302) 451-0829 if you have any questions.

Sincerely,

Rebecca M. Ashley  
Registration Coordinator