

## **PLEASE NOTE**

**This image contains more than one label approved for this product on this date.**

## SUPPLEMENTAL LABELING

**DuPont Agricultural  
Products**

**DUPONT™ OUST XP HERBICIDE  
LOW VOLUME APPLICATION  
ON RAILROAD RIGHTS-OF-WAY**

"..... A Growing Partnership With Nature"

**DUPONT™ OUST® XP HERBICIDE**

EPA Reg. No. 352-601

**LOW VOLUME APPLICATION ON RAILROAD RIGHTS-OF-WAY  
IN THE STATES OF ARIZONA, COLORADO, IDAHO, KANSAS,  
NEBRASKA, NEVADA, NEW MEXICO, NORTH DAKOTA, OKLAHOMA,  
SOUTH DAKOTA, UTAH AND WYOMING**

**DIRECTIONS FOR USE**

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

DuPont™ OUST® XP Herbicide may be applied in 10 to 40 gallons of water per acre by conventional ground application equipment to control various weeds and grasses on railroad rights-of-way.

Refer to the label on the product container for a complete listing of pests controlled and appropriate application rates of the product.


Applications should be made in sufficient volume of water to ensure thorough coverage of the treated site. The volume of water needed to ensure thorough coverage will depend upon several variables including site conditions, such as the size and density of the vegetation, as well as the type and configuration of the application equipment.

**IMPORTANT**

**BEFORE USING DUPONT™ OUST® XP HERBICIDE, READ AND FOLLOW ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA-REGISTERED LABEL.**

This bulletin contains new or supplemental instructions for use of this product which do not appear on the EPA-registered package label. Follow the instructions carefully.

This labeling must be in the possession of the user at the time of pesticide application.




DR184 020402

**NEXT**

**LABEL**

## DuPont Agricultural Products

"..... A Growing Partnership With Nature"



## DUPONT™ OUST® XP HERBICIDE HARDWOODS

# DUPONT™ OUST® XP HERBICIDE

EPA REG. NO. 352-601

## FOR HERBACEOUS WEED CONTROL IN HARDWOOD REFORESTATION AREAS

### DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling

DuPont OUST® XP Herbicide is recommended for herbaceous weed control in commercial reforestation areas where hardwood seedling regeneration is desired following shelterwood seed cuts or where sweetgum (*Liquidambar styraciflua*) plantings are being established. OUST® XP is an effective herbicide providing both foliar and residual activity to control many grasses and broadleaves.

### Hardwood Seedling Regeneration Following Shelterwood Seed Cuts

#### How to use

Apply 2-5 ounces of OUST® XP per acre using appropriate ground equipment and sufficient water, usually 25-50 gallons per acre, for control of hay-scented and New York fern, and germination, growth, and seedhead suppression of short-husk grass. For control of striped maple and beech, tank mixtures with "Roundup"™ at 1-2 quarts per acre are recommended. Calibrate equipment to insure uniform distribution of liquid over the treated area.

Make application from August 1 to September 15 for best results.

Before using OUST® XP in combination, read and carefully observe the cautionary statement and all other information appearing on the product label.

**Note:** Hardwood seedlings present at time of application may be severely injured or killed.

### Selective Sweetgum Release

#### How to Use

Apply 3-5 ounces of OUST® XP per acre using appropriate ground or air (helicopter only) equipment and sufficient water, usually 25-50 gallons per acre by ground and 5-15 gallons per acre by air. Calibrate equipment to insure uniform distribution of liquid over treated area. Apply only after sweetgum seedlings have been transplanted (established) at least one month.

To reduce drift when applying by air, use precise application spray equipment (jet nozzles directed backwards [D4 or larger] or a "Microfoil" boom or equivalent. When compatible with equipment, addition of a spray thickening agent such as "Nalco-Trol" is recommended to further minimize drift. Do not apply by air within 200 feet of any homestead, agricultural land, or other desirable plantings.

Make application from late winter to mid-summer. Best results occur just before or after weeds have germinated. A directed spray may be used to reduce injury to seedlings. When used as described, OUST® XP controls many herbaceous plants such as: crabgrass, dog fennel, fescue, fireweed (willowweed), goldenrod, horseweed, Kentucky bluegrass, nutsedge (yellow), panicums (broadleaf, fall, narrow), pokeweed, ragweed, white snakeroot, and yellow sweetclover.

#### Notes:

Extreme care must be taken to prevent drift or runoff to homesteads, desirable plantings, agricultural land, or any body of water. Do not apply when weather conditions favor drift from treated areas.

Use extreme care when applying in areas adjacent to any body of water. Keep out of lakes, streams, ponds, reservoirs, or any body of water.

OUST® XP is not recommended for use on poorly drained or marshy sites

OUST® XP applications may result in damage to Douglas fir, noble fir, and ponderosa pine.

Use the lower rates on coarse-textured soils (i.e., loamy sands, sandy loams) and higher rates on fine-textured soils (i.e., sandy clay loams and silty clay loams). Do not use on gravelly or rocky soils, exposed subsoils or heavy clay soils (i.e., sandy clay, silty clay, or clay).

### IMPORTANT

**BEFORE USING OUST® XP, READ AND FOLLOW ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA-REGISTERED LABEL.**

This bulletin contains new or supplemental instructions for use of this product which do not appear on the EPA-registered package label. Follow the instructions carefully.

This labeling must be in the possession of the user at the time of pesticide application.

† Trademark of Monsanto Company

DR-200 031802