


352-601

01/24/2007

112



**DuPont Crop
Protection**

DUPONT™ OUST® XP HERBICIDE

EPA Reg. No. 352-601

FOR WEED CONTROL IN HYBRID POPLAR PLANTATIONS IN THE STATE OF NEW MEXICO

DIRECTIONS FOR USE

OUST® XP herbicide controls specific weeds to aid in the establishment and growth of hybrid poplar plantings. It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

OUST® XP USE RATES AND RESTRICTIONS

Site Preparation — Application Before Transplanting
For hybrid poplar, apply 1/2 to 1 1/4 ounces per acre of OUST® XP. Use 1 to 1 1/4 ounces per acre of OUST® XP for heavy weed infestations and where maximum residual control is desired. Use 1/2 to 3/4 ounce per acre of OUST® XP for light weed infestations or when small diameter cuttings have been planted. Allow a minimum of 3 days between application and planting. Limit the first use to a small area to determine the selectivity of OUST® XP on specific clones. OUST® XP must be activated by rainfall or overhead irrigation before weeds become well established. Use of OUST® XP may cause temporary chlorosis (yellowing) or a small reduction in tree height during the year of use.

Release — Application After Transplanting

For hybrid poplar, apply 1/2 to 1 1/4 ounces per acre of OUST® XP. Use 1 to 1 1/4 ounces per acre of OUST® XP for heavy weed infestations and where maximum residual control is desired. Use 1/2 to 3/4 oz per acre of OUST® XP for light weed infestations or when small diameter cuttings have been planted.

Note: Apply only to trees which have been established for a minimum of 1 year. Apply when the trees are dormant and avoid contact of the spray with green buds or tissue as injury to the trees may result. Avoid applications during the period when the hybrid poplar are actively growing; from bud-swell in the spring to leaf drop in the fall. Limit the first use to a small area to determine the selectivity of OUST® XP on specific clones. OUST® XP must be activated by rainfall or overhead irrigation

before weeds become well established.

Use of OUST® XP may cause temporary chlorosis (yellowing) or a small reduction in tree height during the year of use.

TANK MIXES

OUST® XP herbicide can be tank mixed with other products that are registered for use on hybrid poplars and where the labeled method of application and timing of application are the same as for OUST® XP.

Specific Weed Problems

Kochia and Russian Thistle

Since biotypes of kochia and Russian thistle are known to be resistant to OUST® XP, tank mixture combinations with herbicides having different modes of action should be used. To slow the development of resistant biotypes, minimize kochia or Russian thistle forming mature seed.

IMPORTANT PRECAUTIONS

- Applications of OUST® XP made to hybrid poplar trees that are suffering from loss of vigor caused by insects, diseases, drought, winter damage, animal damage, excessive soil moisture, planting shock, previous agricultural practices, or other stresses, may injure or kill the trees.
- Applications of OUST® XP made for release (trees present) should only be made after adequate rainfall has closed the planting slit and settled the soil around the roots following transplanting.
- If a surfactant is used with OUST® XP, allowing the spray to contact tree foliage may injure or kill trees. The user assumes all responsibility for tree injury if a surfactant is used with OUST® XP treatments applied after planting.
- OUST® XP applications may result in damage and mortality to other species of trees when they are present on sites

© 2006 E. I. du Pont de Nemours and Company, Crop Protection, Wilmington, Delaware 19893

BEFORE USING THIS HERBICIDE, READ AND CAREFULLY NOTE THE CAUTIONARY STATEMENTS AND OTHER PROCEDURAL INFORMATION APPEARING ON THE EPA REGISTERED LABEL OR ON OTHER SUPPLEMENTAL LABELS.

This bulletin contains new or supplemental instructions for use of these products in combination which does not appear on the package label. Follow the instructions carefully.

This labeling must be in the possession of the user at the time of pesticide application.

Read the Limitation of Warranty and Liability on the Section 3 Federal product label before buying or using this product. If terms are not acceptable, return the unopened package at once to Seller for full refund of purchase price paid. Otherwise, use by Buyer or any other User constitutes acceptance of the terms of the Limitation of Warranty and Liability on the Section 3 Federal product label.

DR-664 112806

© 2006 E. I. du Pont de Nemours and Company, Crop Protection, Wilmington, Delaware 19898

H -

Page 2 of 2