



352-604

08/05/2003

1/12

|  |  |                                  |  |
|--|--|----------------------------------|--|
|   | U.S. ENVIRONMENTAL PROTECTION AGENCY<br>Office of Pesticide Programs<br>Registration Division (7505C)<br>1200 Pennsylvania Ave., N.W.<br>Washington, D.C. 20460-0001 | EPA Reg. Number:<br>352-604      | Date of Issuance:<br>AUG 5 2003            |
|  | NOTICE OF PESTICIDE:<br><input checked="" type="checkbox"/> Registration<br><input type="checkbox"/> Reregistration<br>(under FIFRA, as amended)                     | Term of Issuance:<br>Conditional | Name of Pesticide Product:<br>DuPont Tanos |
| Name and Address of Registrant (include ZIP Code):<br>DuPont Crop Protection<br>Stine-Haskell Research Center<br>Newark, DE 19714-0030      Attn: Richard A. Carver, Ph.D.   |  |                                  |  |
| Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.  |  |                                  |  |
| <p>On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA). Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.</p> <p>This product is conditionally registered in accordance with FIFRA Section 3(c)(7)(C) provided that you:</p> <ol style="list-style-type: none"> <li>Submit and/or cite all data required for registration of your product under FIFRA Section 3(c)(5) when the Agency requires all registrants of similar products to submit such data; and submit acceptable responses required for reregistration of your product under FIFRA Section 4.</li> <li>Make the following label changes before you release the product for shipment:           <ol style="list-style-type: none"> <li>Revise the EPA Registration Number to read, "EPA Reg. No. 352-604."</li> <li>Revise the statement on page 2: "... is rainfast within one hour ..." to "is rainfast within 12 hours ...".</li> <li>Delete the statement: "Up to 12 oz per acre per application of Tanos may be applied when disease pressure is high, ...".</li> <li>The paragraph: "This product may contaminate water through drift of spray in wind. This product has a potential for runoff ..." should be moved to the Environmental Hazards section of the label under the heading "Surface Water Advisory". In addition, the 25-foot buffer strip is a requirement and label language must be revised to indicate that this 25-foot buffer strip is mandatory.</li> </ol> </li> <li>The Agency recommends that you add Resistance management instructions for cymoxanil as well as famaxodone. Suggested language on how to label a product with two or more active ingredients is listed in the Pesticide Registration Notice 2001-5 on page 8. This is a recommendation, not a requirement.</li> </ol> |  |                                  |  |
| Signature of Approving Official:<br><br>Cynthia Giles-Parker<br>Product Manager (22)<br>Fungicide Branch  |  | Date:<br>AUG 5 2003              |  |

page 2

EPA Reg. No. 352-604

Submit one copy of the revised final printed label before releasing the product for shipment.

If the conditions enumerated above are not complied with, the registration will be subject to cancellation in accordance with FIFRA Section 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Enclosure 1) Stamped Label



DuPont™

Tanos™

fungicide

ACCEPTED  
with COMMENTS  
In EPA Letter Dated:  
AUG 5 2003

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act,  
as amended, for the pesticide  
registered under EPA Reg. No.

352-604

**Dry Flowable**

| <i>Active Ingredients</i> | <i>By Weight</i> |
|---------------------------|------------------|
| Famoxadone                | 25%              |
| Cymoxanil                 | 25%              |
| <i>Inert Ingredients</i>  | <i>50%</i>       |
| TOTAL                     | 100%             |

EPA Reg. No. 352-XXX

**KEEP OUT OF REACH OF CHILDREN**

**CAUTION**

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS  
AND DOMESTIC ANIMALS**

**CAUTION!** Harmful if swallowed. Causes moderate eye irritation. Harmful if absorbed through skin. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling. Harmful if inhaled. Avoid breathing (dust, vapor or spray mist). Remove contaminated clothing and wash clothing before reuse.

**FIRST AID**

**IF INHALED:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.

**IF ON SKIN OR CLOTHING:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

**IF IN EYES:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.

**IF SWALLOWED:** Call poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person. Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-441-3637 for emergency medical treatment information.

**PERSONAL PROTECTIVE EQUIPMENT**

Some materials that are chemical resistant to this product are listed below. If you want more options follow the instructions for Category A on an EPA chemical resistance category selection chart.

**Applicators and other handlers must wear:**

Long-sleeved shirt and long pants.

Chemical Resistant Gloves Category A (such as butyl rubber, natural rubber, neoprene rubber, or nitrile rubber), all  $\geq 14$  mils.

Shoes plus socks.

Follow manufacturer's instructions for cleaning/maintaining personal protective equipment (PPE). If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

**ENGINEERING CONTROL STATEMENTS**

When handlers use enclosed cabs or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR part 170.240 (d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

**USER SAFETY RECOMMENDATIONS**

**USERS SHOULD:** Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

**ENVIRONMENTAL HAZARDS**

This product is toxic to fish and invertebrates. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment wash waters or rinsate. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas.

## DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

**Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.**

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

Coveralls.

Chemical Resistant Gloves Category A (such as butyl rubber, natural rubber, neoprene rubber, or nitrile rubber), all  $\geq 14$  mils.

Shoes plus socks.

DuPont™ TANOS™ fungicide should be used only in accordance with recommendations on this label or in separate published DuPont recommendations or supplemental labels.

DuPont will not be responsible for losses or damages resulting from use of this product in any manner not specifically recommended by DuPont. User assumes all risks associated with such non-recommended use.

Do not formulate this product into other end-use products without written permission from DuPont.

## GENERAL INFORMATION

TANOS™ is a broad-spectrum protectant fungicide, recommended for control of many important plant diseases. It has curative and locally systemic activities against downy mildew and late blight diseases.

TANOS™ must be applied in a regularly scheduled protective spray program in rotation with other fungicides. See directions below for specific crop/disease recommendations.

TANOS™ can be applied with ground, air or chemigation equipment, except as otherwise directed, using sufficient water to obtain thorough coverage of plants. Use only in commercial or farm plantings. Not for use in home plantings or if a commercial crop is turned into U-Pick, Pick-Your-Own or similar operation.

Rainfastness: TANOS™ rapidly penetrates into plant tissues and is rainfast within one hour after application.

This product may contaminate water through drift of spray in wind. This product has a potential for runoff for several months or more after application. Poorly draining soils and soils with shallow water tables are more prone to produce runoff that contains this product. A level, well maintained 25-foot buffer strip between areas to which this product is applied and surface water features such as ponds, streams and springs will reduce the potential for contamination of water from rainfall-runoff. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 48 hours. Sound erosion control practices will reduce this product's contribution to surface water contamination.

## INTEGRATED PEST MANAGEMENT

DuPont recommends the use of Integrated Pest Management (IPM) programs to control pests. This product may be used as part of an Integrated Pest Management (IPM) program which can include biological, cultural, and genetic practices aimed at preventing economic pest damage. Application of this product should be based on IPM principles and practices including field scouting or other detection methods, correct target pest identification, population monitoring, and treating when disease forecasting models reach locally determined action levels. Consult your state cooperative extension service, professional consultants or other qualified authorities to determine the appropriate management, cultural practice and treatment threshold levels for the specific crop, geography and diseases.

## RESISTANCE MANAGEMENT

Repeated use of products for control of specific plant pathogens may lead to selection of resistant strains of fungi and result in a reduction of disease control. Famoxate, one of the active ingredients in TANOS™, is one of EPA's Target Site of Action Group 11 fungicides, which also includes all strobilurins and fenamidone. A disease management program that includes rotation between TANOS™ and other non-Group 11 fungicides is essential to reduce the risk of fungicide resistance development. Tank-mixing TANOS™ with a protectant (contact) fungicide that has a different mode of action is recommended for optimum performance and to further reduce the potential for resistance development. For guidance on the particular crop and disease control situation, consult your state extension specialist or official state recommendations.

## APPLICATION INFORMATION

### PESTICIDE HANDLING

- Calibrate sprayers only with clean water away from the well site.
- Make scheduled checks of spray equipment.
- Ensure accurate measurement of pesticides by all operation employees.
- Mix only enough product for the job at hand.
- Avoid overfilling of spray tank.
- Do not discharge excess material on the soil at a single spot in the field/grove or mixing/loading station.

- Dilute and agitate excess solution and apply at labeled rates/uses.
- Avoid storage of pesticides near well sites.
- When triple rinsing the pesticide container, be sure to add the rinsate to the spray mix.

**MIXING INSTRUCTIONS**

1. Fill the tank 1/4 to 1/3 full of water.
2. While agitating, add the required amount of DuPont™ TANOS™.
3. Continue agitation until the TANOS™ is fully dispersed, at least 5 minutes.
4. Once the TANOS™ is fully dispersed, maintain agitation and continue filling tank with water. TANOS™ must be thoroughly mixed with water before adding any other materials.
5. As the tank is filling, add tank mix partner(s), and any desired adjuvants following the sequence listed in the TANK MIXTURES COMPATIBILITY section of this label. See tank mix partners labels for recommended adjuvants.
6. If the mixture is not continuously agitated, settling will occur. If settling occurs, thoroughly re-agitate before using.
7. Apply TANOS™ spray mixture within 12 hours of mixing to avoid product degradation. If the pH of the spray solution is above 7, either add a buffering agent to lower the pH to below 7 or apply to spray solution immediately.
8. If TANOS™ and a tank-mix partner(s) are to be applied in multiple loads, pre-slurry the TANOS™ in clean water prior to adding to the tank. This will prevent the tank-mix partner(s) from interfering with the dissolution of TANOS™.

**TANK MIXTURES/COMPATIBILITY**

TANOS™ is compatible with many commonly used fungicides, liquid fertilizers, herbicides, insecticides, adjuvants, and biological control products. Optimum disease control is achieved when TANOS™ is applied in a tank-mix with fungicides that have a different mode-of-action (non-Group 11 fungicides). Refer to tank-mix partner label(s) for information on fungal diseases controlled, application information and conditions, and use restrictions. Unless specified on this label or a TANOS™ supplemental label, follow the label guidelines that are most restrictive.

The physical compatibility of TANOS™ with tank-mix partner(s) must be evaluated before use. To determine the physical compatibility, the recommended proportions of products must be added into a suitable container of water in the following sequence:

1. TANOS™ and other water dispersible granules
2. Wettable powders
3. Liquid flowables
4. Emulsifiable concentrates
5. Adjuvants

Mix thoroughly and allow to stand for at least 20 minutes. If the combination remains mixed or can be re-mixed readily, it is considered physically compatible.

The crop safety of all potential tank-mixes, including additives and other pesticides, on all crops has not been tested. Before applying any tank-mixture not specifically recommended on this label or other DuPont supplemental labeling, the safety to the target crop must be confirmed. To test for crop safety, apply the combination to a small area of the target crop in accordance with the label instructions to ensure that a phytotoxic response will not occur.

**CROP ROTATION RESTRICTIONS**

The following rotational intervals must be observed when using TANOS™ fungicide:

| Crop   | Rotational Interval in Days |
|--|-----------------------------|
| Cucurbits, Head Lettuce,<br>Peppers, Potatoes and Tomatoes | Anytime                     |
| All other crops  | 30                          |

6/12

### USE RATES AND APPLICATION INSTRUCTIONS

| Crop   | Target Diseases   | Use Rate   | Remarks  |
|--|---|--|--|
| Cucurbits<br>Including:<br>Cantaloupe,<br>Cucumber,<br>Honeydew Melon,<br>Muskmelon,<br>Watermelon,<br>Pumpkin,<br>Summer Squash,<br>Winter Squash,<br>and other cucurbits | Alternaria Leaf Blight<br><i>(Alternaria cucumerina)</i><br>Anthracnose<br><i>(Colletotrichum, spp.)</i><br>Downy mildew<br><i>(Pseudoperonospora cubensis)</i> | 8 oz/acre/<br>application*<br><br>Seasonal<br>maximum -<br>48 oz/acre<br><br>Annual<br>maximum -<br>72 oz/acre | <p><b>Resistance Management:</b> Do not make more than one application of TANOS™ before alternating with a fungicide that has a different mode of action, such as mancozeb, chlorothalonil, etc. Do not make more than four (4) applications of TANOS™ or other Group 11 fungicides (all strobilurins or fenamidone) per crop per acre per season. Do not alternate or tank mix with other Group 11 fungicides. Do not alternate or tank mix with fungicides to which resistance has developed.</p> <p><b>Application Directions</b></p> <ul style="list-style-type: none"> <li>• Make preventive applications on a 5- to 7-day schedule. TANOS™ applications should begin prior to disease development, following the resistance management instructions, above.</li> <li>• TANOS™ must be tank-mixed with an appropriate contact fungicide that has a different mode of action, such as mancozeb, chlorothalonil, copper, etc. Follow all tank-mix partner label restrictions using at least the minimum labeled rates of each fungicide.</li> <li>• For best results, alternate each TANOS™ application with a chlorothalonil application.</li> </ul> <p><b>Minimum Application Volume:</b></p> <ul style="list-style-type: none"> <li>• For ground application, apply a minimum of 20 gallons of spray volume per acre, increasing the spray volume as plants mature to ensure thorough coverage of foliage, blooms and fruit.</li> <li>• For aerial application, apply a minimum of 5 gallons per acre.</li> </ul> <p>Do not use TANOS™ for the control of Gummy Stem Blight or Powdery Mildew.<br/>           Minimum Pre-Harvest Interval (PHI) is 3 days.<br/>           Reentry interval is 12 hours.</p> |
| Head Lettuce   | Downy Mildew<br><i>(Bremia lactucae)</i>  | 8 oz/acre/<br>application*<br><br>Seasonal maximum -<br>36 oz/acre<br><br>Annual maximum -<br>72 oz/acre       | <p><b>Resistance Management:</b> Do not make more than one application of TANOS™ before alternating with a fungicide that has a different mode of action, such as maneb, etc. Do not make more than three (3) applications of TANOS™ or other Group 11 fungicides (all strobilurins or fenamidone) per crop per acre per season. Do not alternate or tank mix with other Group 11 fungicides. Do not alternate or tank mix with fungicides to which resistance has developed.</p> <p><b>Application Directions:</b></p> <ul style="list-style-type: none"> <li>• TANOS™ applications should begin prior to disease development, following the resistance management instructions, above. (cont'd next page)</li> </ul>   |

| Crop                       | Target Diseases   | Use Rate  | Remarks   |
|----------------------------|---|---|---|
| Head Lettuce (Cont'd)      |   |   | <ul style="list-style-type: none"> <li>• Make preventive applications on a 5- to 7-day schedule. TANOS™ must be tank-mixed with an appropriate contact fungicide that has a different mode of action, such as maneb or "Aliette". Follow all tank-mix partner label restrictions using at least the minimum labeled rates of each fungicide.</li> </ul> <p><b>Minimum Application Volume:</b></p> <ul style="list-style-type: none"> <li>• For ground application, apply a minimum of 20 gallons of spray volume per acre, increasing the spray volume as plants mature to ensure thorough coverage of foliage.</li> <li>• For aerial application, apply a minimum of 5 gallons per acre.</li> </ul> <p>Minimum Pre-Harvest Interval (PHI) is 3 days.<br/>Reentry interval is 12 hours.</p>   |
| Peppers<br>(all varieties) | Phytophthora blight<br>( <i>Phytophthora capsici</i> )<br>Foliar and fruit phase ONLY | 8 - 10 oz/acre/<br>application*<br><br>Seasonal maximum –<br>72 oz/acre<br><br>Annual maximum –<br>72 oz/acre | <p><b>Resistance Management:</b> Do not make more than one application of TANOS™ before alternating with a fungicide that has a different mode of action, such as maneb, copper, etc. Do not alternate or tank mix with other Group 11 fungicides (all strobilurins or fenamidone). Do not alternate or tank mix with fungicides to which resistance has developed.</p> <p><b>Application Directions:</b></p> <ul style="list-style-type: none"> <li>• TANOS™ applications should begin prior to disease development, following the resistance management instructions, above.</li> <li>• Make preventive applications on a 5- to 7-day schedule. TANOS™ applications should begin prior to disease development, following the resistance management instructions, above.</li> <li>• TANOS™ must be tank-mixed with an appropriate contact fungicide that has a different mode of action, such as maneb, copper, etc. Follow all tank-mix partner label restrictions using at least the minimum labeled rates of each fungicide.</li> <li>• For best results, tank-mix TANOS™ with a copper-containing fungicide for control of Phytophthora blight.</li> </ul> <p><b>Minimum Application Volume:</b></p> <ul style="list-style-type: none"> <li>• For ground application, apply a minimum of 20 gallons of spray volume per acre, increasing the spray volume as plants mature to ensure thorough coverage of foliage, blooms and fruit.</li> <li>• For aerial application, apply a minimum of 5 gallons per acre.</li> </ul> <p>Minimum Pre-Harvest Interval (PHI) is 3 days.<br/>Reentry interval is 12 hours.</p> |

| Crop     | Target Diseases                              | Use Rate   | Remarks  |
|----------|--|--|--|
| Potatoes | Early Blight<br>( <i>Alternaria solani</i> ) | 6 - 8 oz /acre /application*<br><br>Seasonal maximum – 72 oz/acre<br><br>Annual maximum – 72 oz/acre | <b>Resistance Management:</b> Do not make more than one application of TANOS™ before alternating with a fungicide that has a different mode of action, such as mancozeb, chlorothalonil, etc. Do not make more than six (6) applications of TANOS™ or other Group 11 fungicides (all strobilurins or fenamidone) per crop per acre per season. Do not alternate or tank mix with other Group 11 fungicides. Do not alternate or tank mix with fungicides to which resistance has developed.  |
|          | Late Blight                                  | 8 oz/acre/ application*<br><br>Seasonal maximum – 72 oz/acre<br><br>Annual maximum – 72 oz/acre      | <p><b>Application Directions:</b></p> <ul style="list-style-type: none"> <li>TANOS™ applications should begin prior to disease development, following the resistance management instructions, above.</li> <li>For early blight control, make fungicide applications on a 7- to 10-day interval. Use higher rates and shorter intervals when disease is present in the area or if weather conditions favor disease development.</li> <li>For late blight control, make preventive fungicide applications on a 7-day interval prior to disease development. When weather conditions favor late blight development or late blight is present in the area, make fungicide applications on a 5- to 7- day interval.</li> <li>TANOS™ must be tank-mixed with an appropriate contact fungicide that has a different mode of action, such as mancozeb, chlorothalonil, etc. Follow all tank-mix partner label restrictions using at least the minimum labeled rates of each fungicide.</li> </ul> <p><b>Minimum Application Volume:</b></p> <ul style="list-style-type: none"> <li>For ground application, apply a minimum of 20 gallons of spray volume per acre, increasing the spray volume as plants mature to ensure thorough coverage of foliage.</li> <li>For aerial application, apply a minimum of 5 gallons per acre.</li> </ul> <p>Minimum Pre-Harvest Interval (PHI) is 14 days.<br/>Reentry interval is 12 hours.</p> |



| Crop     | Target Diseases  | Use Rate  | Remarks   |
|----------|--|---|---|
| Tomatoes | Early Blight<br><i>(Alternaria solani)</i><br>Septoria Leaf Spot<br><i>(Septoria lycopersici)</i>  | 6-8 oz/acre /application*<br><br>Seasonal maximum – 72 oz/acre<br><br>Annual maximum – 72 oz/acre | <p><b>Resistance Management:</b> Do not make more than one application of TANOS™ before alternating with a fungicide that has a different mode of action, such as mancozeb, chlorothalonil, copper, etc. Do not alternate or tank mix with other Group 11 fungicides (all strobilurins or fenamidone). Do not alternate for tank mix with fungicides to which resistance has developed.</p> <p><b>Application Directions:</b></p> <ul style="list-style-type: none"> <li>TANOS™ applications should begin prior to disease development, following the resistance management instructions, above.</li> <li>Make preventive applications on a 5- to 7-day schedule.</li> <li>For Early blight and Septoria leaf spot control, use higher rates when disease is present in the area or if weather conditions favor disease development.</li> <li>TANOS™ must be tank-mixed with an appropriate contact fungicide that has a different mode of action, such as mancozeb, chlorothalonil, copper, etc. Follow all tank-mix partner label restrictions using at least the minimum labeled rates of each fungicide.</li> <li>For best results, tank-mix TANOS™ with a full rate of a copper-containing fungicide for suppression of bacterial diseases.</li> </ul> <p><b>Minimum Application Volume:</b></p> <ul style="list-style-type: none"> <li>For ground application, apply a minimum of 20 gallons of spray volume per acre, increasing the spray volume as plants mature to ensure thorough coverage of foliage, blooms and fruit.</li> <li>For aerial application, apply a minimum of 5 gallons per acre.</li> </ul> <p>Minimum Pre-Harvest Interval (PHI) is 3 days.<br/>Reentry interval is 12 hours.</p> |
|          | Anthracnose<br><i>(Colletotrichum spp.)</i><br>Gray Leaf Mold<br><i>(Cladosporium fulvum)</i><br>Late Blight<br><i>(Phytophthora infestans)</i><br>Target Spot<br><i>(Corynespora cassicola)</i> | 8 oz/acre /application*<br><br>Seasonal maximum – 72 oz/acre<br><br>Annual maximum – 72 oz/acre   |   |
|          | <p><b>Disease Suppression:</b></p> Bacterial Spot<br><i>(Xanthomonas campestris)</i><br>Bacterial Speck<br><i>(Pseudomonas syringae)</i><br>Buckeye Rot<br><i>(Phytophthora spp.)</i>            | 8 oz/acre /application*<br><br>Seasonal maximum – 72 oz/acre<br><br>Annual maximum – 72 oz/acre   |   |

\* Up to 12 oz per acre per application of TANOS™ may be applied when disease pressure is high, based on supplemental DuPont labeling, local recommendations for specific pathogens from your state cooperative extension service, professional consultants, or other qualified authorities.

## CHEMIGATION

Apply DuPont™ TANOS™ only through sprinkler irrigation systems (such as center pivot, lateral move, end tow, side (wheel) roll, traveler, big gun, solid set or hand move irrigation systems). Do not apply this product through any other type of irrigation system.

Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water.

If you have questions about calibration, you should contact State Extension Service Specialists, equipment manufacturers or other experts.

Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place.

A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

### *Specific Instructions for Sprinkler Irrigation Systems:*

1. The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.
2. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
3. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
4. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
5. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
6. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
7. Do not apply when wind speed favors drift beyond the area intended for treatment.
8. Good agitation is required in the injection tank.
9. In moving systems, apply specified dosage of TANOS™ as a continuous injection. In nonmoving systems inject TANOS™ for 15 to 30 minutes at end of cycle. Use the least amount of water possible consistent with uniform coverage.

10. Mix the amount of TANOS™ needed for acreage to be treated into the quantity of water determined during prior calibration. For moving systems inject into the system continuously for one complete revolution of the field. For nonmoving systems inject into system for the time established during calibration.

11. Stop injection equipment after treatment is completed and continue to operate irrigation equipment until all TANOS™ is flushed from system.

### *Specific Instructions for Public Water Systems:*

1. Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.
2. Chemigation systems connected to public water systems must contain a functional, reduced-pressure zone, backflow preventer (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.
3. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
4. The pesticide injection pipeline must contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
5. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected.
6. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
7. Do not apply when wind speed favors drift beyond the area intended for treatment.

## SPRAY DRIFT MANAGEMENT

The interaction of many equipment and weather-related factors determines the potential for spray drift. The applicator is responsible for considering all these factors when making application decisions.

**AVOIDING SPRAY DRIFT IS THE RESPONSIBILITY OF THE APPLICATOR.**

**IMPORTANCE OF DROPLET SIZE**

The most effective way to reduce drift potential is to apply large droplets (>150 - 200 microns). The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. The presence of sensitive species nearby, the environmental conditions, and pest pressure may affect how an applicator balances drift control and coverage. **APPLYING LARGER DROPLETS REDUCES DRIFT POTENTIAL, BUT WILL NOT PREVENT DRIFT IF APPLICATIONS ARE MADE IMPROPERLY OR UNDER UNFAVORABLE ENVIRONMENTAL CONDITIONS!** See **Wind, Temperature and Humidity**, and **Temperature Inversions** sections of this label.

**Controlling Droplet Size - General Techniques**

- **Volume** - Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets.
- **Pressure** - Use the lower spray pressures recommended for the nozzle. Higher pressure reduces droplet size and does not improve canopy penetration. **WHEN HIGHER FLOW RATES ARE NEEDED, USE A HIGHER-CAPACITY NOZZLE INSTEAD OF INCREASING PRESSURE.**
- **Nozzle Type** - Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles.

**Controlling Droplet Size - Aircraft**

- **Number of Nozzles** - Use the minimum number of nozzles with the highest flow rate that provide uniform coverage.
- **Nozzle Orientation** - Orienting nozzles so that the spray is emitted backwards, parallel to the air stream will produce larger droplets than other orientations.
- **Nozzle Type** - Solid stream nozzles (such as disc and core with swirl plate removed) oriented straight back produce larger droplets than other nozzle types.

**BOOM LENGTH AND HEIGHT**

- **Boom Length (aircraft)** - The boom length should not exceed 3/4 of the wing length, using shorter booms decreases drift potential. For helicopters use a boom length and position that prevents droplets from entering the rotor vortices.
- **Boom Height (aircraft)** - Application more than 10 ft above the canopy increases the potential for spray drift.
- **Boom Height (ground)** Setting the boom at the lowest height which provides uniform coverage reduces the exposure of droplets to evaporation and wind. The boom should remain level with the crop and have minimal bounce.

**WIND**

Drift potential increases at wind speeds of less than 3 mph (due to variable direction and inversion potential) or more than 10 mph. However, many factors, including droplet size and equipment type determine drift potential at any given wind speed. **AVOID GUSTY OR WINDLESS CONDITIONS.**

**Note:** Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they effect spray drift.

**TEMPERATURE AND HUMIDITY**

When making applications in hot and dry conditions, set up equipment to produce larger droplets to reduce effects of evaporation.

**SURFACE TEMPERATURE INVERSIONS**

Drift potential is high during a surface temperature inversion. Surface inversions restrict vertical air mixing, which causes small suspended droplets to remain close to the ground and move laterally in a concentrated cloud. Surface inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates a surface inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

**SHIELDED SPRAYERS**

Shielding the boom or individual nozzles can reduce the effects of wind. However, it is the responsibility of the applicator to verify that the shields are preventing drift and not interfering with uniform deposition of the product.

**AIR ASSISTED (AIR BLAST) FIELD CROP SPRAYERS**

Air assisted field crop sprayers carry droplets to the target via a downward directed air stream. Some may reduce the potential for drift, but if a sprayer is unsuitable for the application and/or set up improperly, high drift potential can result. It is the responsibility of the applicator to determine that a sprayer is suitable for the intended application, is configured properly, and that drift is not occurring.

**Note:** Air assisted field sprayers can affect product performance by affecting spray coverage and canopy penetration. Consult the application equipment section of this label to determine if use of an air assisted sprayer is recommended.

**SPRAY TANK CLEANOUT**

Prior to application, start with clean, well maintained application equipment. Immediately following application, thoroughly clean all spray equipment to reduce the risk of forming hardened deposits which might become difficult to remove.

Drain spray equipment. Thoroughly rinse sprayer and flush hoses, boom and nozzles with clean water.

Clean all other associated application equipment. Take all necessary safety precautions when cleaning equipment. Do not clean near wells, water sources or desirable vegetation. Dispose of waste rinse water in accordance with local regulations.

**STORAGE AND DISPOSAL**

**Storage:** Store product in original container only. Do not contaminate water, other pesticides, fertilizer, food or feed in storage. Store in a cool, dry place.

**Product Disposal:** Do not contaminate water, food, or feed by disposal. Waste resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**Container Disposal: For Plastic Containers:** Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke. **For Fiber Sacks:** Completely empty fiber sack by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into manufacturing or application equipment. Then dispose of sack in a sanitary landfill or by incineration if allowed by State and local authorities. **For Fiber Drums With Liners:** Completely empty liner by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application equipment. Then dispose of liner in a sanitary landfill or by incineration if allowed by State and local authorities. If drum is contaminated and cannot be reused, dispose of in the same manner. **For Bags Containing Water Soluble Packets:** Do not reuse the outer box or the re-sealable plastic bag. When all water-soluble packets are used, the outer packaging should be clean and may be disposed of in a sanitary landfill or by incineration, or if allowed by State and local authorities, by open burning. If burned, stay out of smoke. If the re-sealable plastic bag contacts the formulated product in any way, the bag must be triple-rinsed with clean water. Add the rinsate to the spray tank and dispose of the outer wrap as described above. **For Metal Containers (non aerosol):** Triple rinse (or equivalent) the container. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities. **For Paper and Plastic Bags:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

**NOTICE TO BUYER--**Purchase of this material does not confer any rights under patents of countries outside of the United States.

The DuPont Oval Logo, DuPont™, and TANOS™ are trademarks or registered trademarks of DuPont or its affiliates

"Aliette" is a registered trademark of Bayer CropScience

**D - 670 060603**

**Net Wt. \_\_\_\_\_**

**LIMITATION OF WARRANTY AND LIABILITY**

**NOTICE:** Read This Limitation of Warranty and Liability Before Buying or Using This Product. If the Terms Are Not Acceptable, Return the Product at Once, Unopened, and the Purchase Price Will Be Refunded.

It is impossible to eliminate all risks associated with the use of this product. Such risks arise from weather conditions, soil factors, off target movement, unconventional farming techniques, presence of other materials, the manner of use or application, or other unknown factors, all of which are beyond the control of DuPont. These risks can cause: ineffectiveness of the product; crop injury, or; injury to non-target crops or plants.

DuPont does not agree to be an insurer of these risks. **WHEN YOU BUY OR USE THIS PRODUCT, YOU AGREE TO ACCEPT THESE RISKS.**

DuPont warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for the purpose stated in the Directions for Use, subject to the inherent risks described above, when used in accordance with the Directions for Use under normal conditions.

**DUPONT MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR OF MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.**

**IN NO EVENT SHALL DUPONT OR SELLER BE LIABLE FOR ANY INCIDENTAL, CONSEQUENTIAL OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. BUYER'S OR USER'S BARGAINED-FOR EXPECTATION IS CROP PROTECTION. THE EXCLUSIVE REMEDY OF THE USER OR BUYER AND THE EXCLUSIVE LIABILITY OF DUPONT OR SELLER, FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY OR CONTRACT, NEGLIGENCE, TORT OR STRICT LIABILITY), WHETHER FROM FAILURE TO PERFORM OR INJURY TO CROPS OR OTHER PLANTS, AND RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT, OR AT THE ELECTION OF DUPONT OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

DuPont or its Ag Retailer must have prompt notice of any claim so that an immediate inspection of buyer's or user's growing crops can be made. Buyer and all users shall promptly notify DuPont or a DuPont Ag Retailer of any claims, whether based on contract, negligence, strict liability, other tort or otherwise or be barred from any remedy.

This Limitation of Warranty and Liability may not be amended by any oral or written agreement.

For product information call: 1-888-6-DUPONT

Internet address: <http://cropprotection.dupont.com/>

© 2003 E. I. du Pont de Nemours and Company, Wilmington, Delaware 19898.

All rights reserved.