


352-607

5-8-2002

SUPPLEMENTAL LABELING ^{1/2}

**DuPont Agricultural
Products**

"..... A Growing Partnership With Nature"



**DUPONT™ GLYPHOSATE
HERBICIDE**

DUPONT™ GLYPHOSATE HERBICIDE

EPA Reg. No. 352-607

**CHEMICAL FALLOW, PREPLANT FALLOW BEDS, PREPLANT, PRE-
EMERGENCE, AT-PLANTING, HOODED SPRAYERS, WIPER APPLICA-
TORS IN ROWS, AND POST-HARVEST TREATMENTS IN
STEVIA CROPS**

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

This label must be in the possession of the user at the time of herbicide application.

AVOID CONTACT OF HERBICIDE WITH FOLIAGE, GREEN STEMS, EXPOSED NON-WOODY ROOTS OR FRUIT OF CROPS, DESIRABLE PLANTS AND TREES, BECAUSE SEVERE INJURY OR DESTRUCTION MAY RESULT.

See the "GENERAL INFORMATION", "MIXING", "ANNUAL WEEDS RATE TABLE", and "PERENNIAL WEEDS RATE TABLE" sections of the label booklet for DuPont GLYPHOSATE herbicide for essential product performance information.

Treatment Recommendations for Stevia Crops

TYPES OF APPLICATIONS: Chemical Fallow, Preplant Fallow beds, Preplant, Preemergence, At-Planting, hooded Sprayers in rows, Wiper Applicators in rows, and Post-harvest Treatments in Stevia crops.

Avoid contact of herbicide with foliage, green shoots or stems, bark or exposed roots (including those emerging from plastic mulch) of crops because severe injury or destruction may result. Applications made at emergence will result in injury or death to emerged seedlings. Hooded sprayers and wiper equipment capable of preventing all crop contact with herbicide solutions may be used only in row middles when such equipment can be operated without leakage of spray mists or dripping onto crop. Treatments with wipers and hooded sprayers must be made at least 14 days prior to harvest. Post harvest or fallow applications must be made at least 30 days prior to planting any non-labeled crop.

See the "Application Equipment and Techniques" section of the label for additional information.

USE INSTRUCTIONS: This product may be applied during fallow intervals preceding planting, prior to planting or transplanting, at-planting, or preemergent to Stevia crops. Post-directed hooded sprayers may be used in mulched or unmulched row middles after crop establishment. Wiper applicators may be used to control weeds in row middles where any dripping or leaking will not contact crop foliage. Apply 1 pint to 5 quarts per acre according to the rates specified in the "ANNUAL WEEDS RATE TABLE" AND "PERENNIAL WEEDS RATE TABLE" SECTION OF THE Glyphosate Herbicide label. Repeat applications may be made up to a maximum of 8 quarts per acre per year. The maximum use rates stated throughout this product's labeling apply to this product combined with the use of all other herbicides containing glyphosate or sulfosate as the active ingredient whether applied as mixtures or separately. Calculate the application rates and ensure that the total use of this and other glyphosate or sulfosate containing products does not exceed stated maximum use rate.

Hooded sprayers must be fully enclosed including top, sides, front and back. Use only hoods designed to minimize excessive dripping or run-off down the insides of the hoods. A single, low pressure/low drift flat fan nozzle with an 80 to 95 degree spray angle positioned at the top center of the hood as recommended. Spray volume should be 20 to 30 gallons per acre. Hoods should be run so that the sides contact the ground, eliminating escape of spray droplets.

PRECAUTIONS AND RESTRICTIONS

Avoid contact of herbicide with foliage, green shoots or stems, bark or exposed roots (including those emerging from plastic mulch) of crops because severe injury or destruction may result. Applications made at emergence will result in injury or death to

2/2

emerged seedlings. Hooded sprayers and wiper equipment capable of preventing all crop contact with herbicide solutions may be used only in row middles when such equipment can be operated without leakage of spray mists or dropping onto crop. treatments with wipers and hooded sprayers must be made at least 14 days prior to harvest. Post harvest or fallow applications must be made at least 30 days prior to planting any non-labeled crop.

When making preemergence and at planting applications, apply before seed germination in coarse sandy soils to minimize the risk of injury.

Residues of this product remaining on plastic mulch could cause injury in transplanted crops. Take care to remove such residues prior to transplanting. Residues can be removed by a single 0.5 inch application of water, either by natural rainfall or via a sprinkler system.

Crop injury is possible with these applications and shall be the sole responsibility of the applicator.

**IMPORTANT
BEFORE USING THESE PRODUCTS, READ
AND FOLLOW ALL APPLICABLE DIRECTIONS,
RESTRICTIONS AND PRECAUTIONS
ON THE EPA-REGISTERED LABEL.**

This bulletin contains new or supplemental instructions for use of this product which do not appear on the EPA-registered package label. Follow the instructions carefully.

This labeling must be in the possession of the user at the time of pesticide application.

DR-128 013002-3

