

PLEASE NOTE

**This image contains more than one label
approved for this product on this date.**



352-610

3-28-2002

1/1

**DuPont Agricultural
Products**



SUPPLEMENTAL LABELING

**DUPONT™ ALLY® EXTRA HERBICIDE
DUPONT™ CANVAS® XP HERBICIDE
PLUS STARANE*1 + SWORD®2
HERBICIDES**

"..... A Growing Partnership With Nature"

DUPONT™ ALLY® EXTRA HERBICIDE

Reg. No. 352-610

DUPONT™ CANVAS® XP HERBICIDE

EPA Reg. No. 352-610

PLUS STARANE* + SWORD® HERBICIDES

DuPont™ ALLY® EXTRA or DuPont™ CANVAS® XP herbicide can be tank mixed with Starane* + Sword® herbicides for improved control of broadleaf weeds in wheat, barley, and fallow.

For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

DIRECTIONS FOR USE

For improved control of Kochia (2-4" tall) Russian thistle, mustard species and wild buckwheat, ALLY® EXTRA or CANVAS® XP herbicide may be tank mixed with 3/4 to 2 3/4 pints per acre of Starane* + Sword®. Refer to the DuPont herbicide label, and the Starane* + Sword® label for information regarding use restrictions, labeled crops, rotational cropping recommendations, sprayer cleanup, use precautions and other information. The most restrictive provisions on either label will apply. Do not use the tank mix if any restrictions on the Starane* + Sword® label conflict with recommendations on the DuPont herbicide label.

Other suitable registered herbicides, fungicides, and insecticides registered for use on cereal grains or fallow may be tank mixed or used sequentially with this mixture. Read and follow all manufacturer's label recommendations for the companion herbicide. The most restrictive provisions on either label will apply.

It is a violation of federal law to use this product in a manner inconsistent with its labeling.


IMPORTANT

BEFORE USING DUPONT ALLY® EXTRA OR CANVAS® XP HERBICIDE, READ AND FOLLOW ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA-REGISTERED LABEL.

This bulletin contains new or supplemental instructions for the use of this product which does not appear on the EPA-registered package label. Follow these instructions carefully. This labeling must be in the possession of the user at the time of application.

¹ Starane is a trademark of Dow AgroSciences LLC

² Sword is a trademark of Platte Chemical Co.



DR-189 030402

For product information call 1-888-6-DUPONT
© 2002 E. I. du Pont de Nemours and Company, Crop Protection, Wilmington, Delaware 19898

NEXT

LABEL



352-610

3-28-2002

1/1

**DuPont Agricultural
Products**



SUPPLEMENTAL LABELING

**DUPONT™ ALLY® EXTRA HERBICIDE
DUPONT™ CANVAS® XP HERBICIDE
PLUS STARANE™ + SALVO®
HERBICIDES**

"..... A Growing Partnership With Nature"

DUPONT™ ALLY® EXTRA HERBICIDE

EPA Reg. No. 352-610

DUPONT™ CANVAS® XP HERBICIDE

EPA Reg. No. 352-610

PLUS STARANE™ + SALVO® HERBICIDES

DuPont™ ALLY® EXTRA or DuPont™ CANVAS® XP herbicide can be tank mixed with Starane™ + Salvo® herbicides for improved control of broadleaf weeds in wheat, barley, and fallow.

For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

DIRECTIONS FOR USE

For improved control of Kochia (2-4" tall), Russian thistle, mustard species and wild buckwheat, ALLY® EXTRA or CANVAS® XP herbicide may be tank mixed with 2/3 to 2 2/3 pints per acre of Starane™ + Salvo®. Refer to the DuPont herbicide label, and the Starane™ + Salvo® label for information regarding use restrictions, labeled crops, rotational cropping recommendations, sprayer cleanup, use precautions and other information. The most restrictive provisions on either label will apply. Do not use the tank mix if any restrictions on the Starane™ + Salvo® label conflict with recommendations on the DuPont herbicide label.

Other suitable registered herbicides, fungicides, and insecticides registered for use on small grains or fallow may be tank mixed or used sequentially with this mixture. Read and follow all manufacturer's label recommendations for the companion herbicide. The most restrictive provisions on either label will apply.


It is a violation of federal law to use this product in a manner inconsistent with its labeling.

IMPORTANT

BEFORE USING DUPONT ALLY® EXTRA OR CANVAS® XP HERBICIDE, READ AND FOLLOW ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA-REGISTERED LABEL.

This bulletin contains new or supplemental instructions for the use of this product which does not appear on the EPA-registered package label. Follow these instructions carefully. This labeling must be in the possession of the user at the time of application.

- 1 Starane is a trademark of Dow AgroSciences LLC
- 2 Salvo is a trademark of Platte Chemical Co.



DR-191 030402

For product information call 1-888-6-DUPONT

© 2002 E. I. du Pont de Nemours and Company, Crop Protection, Wilmington, Delaware 19898

NEXT

LABEL



352-610

3-28-2002

1/1

**DuPont Agricultural
Products**



SUPPLEMENTAL LABELING

**DUPONT™ ALLY® EXTRA HERBICIDE
DUPONT™ CANVAS® XP HERBICIDE
PLUS STARANE¹ HERBICIDE**

"..... A Growing Partnership With Nature"

DUPONT™ ALLY® EXTRA HERBICIDE

EPA Reg. No. 352-610

DUPONT™ CANVAS® XP HERBICIDE

EPA Reg. No. 352-610

PLUS STARANE^{*} HERBICIDE

DuPont™ ALLY® EXTRA or DuPont™ CANVAS® XP herbicide can be tank mixed with Starane^{*} herbicide for improved control of broadleaf weeds in wheat, barley, and fallow.

For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

DIRECTIONS FOR USE

For improved control of Kochia (2-4" tall), Russian thistle, mustard species, and wild buckwheat ALLY® EXTRA or CANVAS® XP herbicide may be tank mixed with 1/3 to 1 1/3 pints per acre of Starane^{*}. Refer to the DuPont herbicide label, and the Starane^{*} label for information regarding use restrictions, labeled crops, rotational cropping recommendations, sprayer cleanup, use precautions and other information. The most restrictive provisions on either label will apply. Do not use the tank mix if any restrictions on the Starane^{*} label conflict with recommendations on the DuPont herbicide label.

Other suitable registered herbicides, fungicides, and insecticides registered for use on small grains or fallow may be tank mixed or used sequentially with this mixture. Read and follow all manufacturer's label recommendations for the companion herbicide. The most restrictive provisions on either label will apply.


It is a violation of federal law to use this product in a manner inconsistent with its labeling.

IMPORTANT

BEFORE USING DUPONT ALLY® EXTRA OR CANVAS® XP HERBICIDE, READ AND FOLLOW ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA-REGISTERED LABEL.

This bulletin contains new or supplemental instructions for the use of this product which does not appear on the EPA-registered package label. Follow these instructions carefully. This labeling must be in the possession of the user at the time of application.

¹ Starane is a trademark of Dow AgroSciences LLC



DR-187 030402

For product information call 1-888-6-DUPONT
© 2002 E. I. du Pont de Nemours and Company, Crop Protection, Wilmington, Delaware 19898

NEXT

LABEL

**DuPont Agricultural
Products**

DRAFT

**ALLY® EXTRA HERBICIDE
CANVAS® XP HERBICIDE**
ROTATION INTERVAL TO
GRAIN SORGHUM, STS™ SOYBEANS,
IR CORN, PROSO MILLET

1/2

"..... A Growing Partnership With Nature"

ACCEPTED

MAR 28 2002

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 352-610

DUPONT ALLY® EXTRA HERBICIDE

EPA Reg. No. 352-610

DUPONT CANVAS® XP HERBICIDE

EPA Reg. No. 352-610

**ROTATION INTERVAL TO GRAIN SORGHUM,
STS® SOYBEAN¹, IR CORN², PROSO MILLET**

DIRECTIONS FOR USE

DuPont ALLY® EXTRA Herbicide and DuPont CANVAS® XP Herbicide are dry flowable granule formulations containing 71.25% active ingredient by weight.

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

CROP ROTATION

Minimum Rotation Intervals

Minimum rotation intervals* are determined by the rate of breakdown of ALLY® EXTRA or CANVAS® XP applied. ALLY® EXTRA or CANVAS® XP breakdown in the soil is affected by soil pH, soil temperature, soil microorganisms, and soil moisture. Low soil pH, high soil temperature, and high soil moisture increase ALLY® EXTRA or CANVAS® XP breakdown in soil, while high soil pH, low soil temperature, and low soil moisture slow ALLY® EXTRA or CANVAS® XP breakdown.

Of these three factors, only soil pH remains relatively constant. Soil temperature, and to a greater extent, soil moisture, can vary significantly from year to year and from area to area. For this reason, soil temperatures and soil moisture should be monitored regularly when considering rotating to other crops.

*The minimum rotation interval represents the period of time from the last application to the anticipated date of the next planting. Rotation intervals are based on normal precipitation/irrigation amounts. See the EPA approved ALLY® EXTRA or CANVAS® XP label for additional details on crop rotation recommendations and restrictions.

Soil pH Limitations

ALLY® EXTRA or CANVAS® XP should not be used on fields having a soil pH above 7.9, as extended soil residual activity could extend crop rotation intervals beyond those specified in the rotation table, and under certain conditions, could injure wheat or barley. In addition, other crops planted in high-pH soils can be extremely sensitive to low concentrations of ALLY® EXTRA or CANVAS® XP.

Before using ALLY® EXTRA or CANVAS® XP, determine the soil pH of the field. To obtain a representative pH value, take several samples from different areas of the field between 0" and 4" deep and analyze them separately. Consult local extension publications for additional information on recommended soil sampling procedures. Before using ALLY® EXTRA or CANVAS® XP, carefully consider your crop rotation plans and options. For rotational flexibility, do not treat all of your wheat, barley, or fallow acres at the same time.

Rotation Intervals are shown below for Non-Cereal Crops (Grain Sorghum, STS Soybeans, IR Corn, Proso Millet) on Irrigated/Non Irrigated Land following wheat, barley or fallow treated with ALLY® EXTRA or CANVAS® XP at the maximum use rate.

Areas	Rotational Crop	Soil pH	Rotation Interval (months)*
All Areas of TX, OK, KS, NE and CO	Grain Sorghum, STS Soybeans, IR Corn, Proso Millet	7.9 or lower	4

2/2

IMPORTANT PRECAUTIONS

- Injury to or loss of desirable trees or vegetation may result from failure to observe the following:
 - Do not apply, drain or flush equipment on or near desirable trees or other plants, or on areas where their roots may extend, or in locations where the chemical may be washed or moved into contact with their roots.
 - Do not use on lawns, walks, driveways, tennis courts or similar areas.
 - Prevent drift of spray to desirable plants.
- Do not contaminate any body of water. Thoroughly clean application equipment immediately after use
- ALLY® EXTRA and CANVAS® XP are non-corrosive, non-flammable, non-volatile, and do not freeze in storage.
- Under certain conditions (such as drought, prolonged cold weather, pH variability in the fields) temporary discoloration and/or crop injury may occur to sorghum, STS soybeans, IR corn, and Proso Millet planted after ALLY® EXTRA or CANVAS® XP applications.
- This supplemental label is not applicable to crops grown for seed.

IMPORTANT

BEFORE USING ALLY® EXTRA OR CANVAS® XP, READ AND FOLLOW ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA-REGISTERED LABEL.

This bulletin contains new or supplemental instructions for use of this product which do not appear on the EPA-registered package label. Follow the instructions carefully.

This labeling must be in the possession of the user at the time of pesticide application.

¹ Sulfonylurea Tolerant Soybeans

² Imidazolinone Resistant Corn

DR-143 011002

NEXT

LABEL

**DuPont Agricultural
Products**

DRAFT

**ALLY® EXTRA AND
CANVAS® XP
HERBICIDES**

"..... A Growing Partnership With Nature"

DUPONT ALLY® EXTRA HERBICIDE

EPA Reg. No. 352-610

DUPONT CANVAS® XP HERBICIDE

EPA Reg. No. 352-610

REDUCED ROTATION INTERVAL FOR CONDIMENT MUSTARD AND CHICKPEAS (GARBANZO BEANS) IN THE STATES OF IDAHO, OREGON, AND WASHINGTON

DIRECTIONS FOR USE

It is violation of federal law to use this product in a manner inconsistent with its labeling.

DuPont ALLY® EXTRA Herbicide and DuPont CANVAS® XP Herbicide are selective postemergence herbicides recommended for control of broadleaf weeds in wheat, barley, and fallow.

HOW TO USE

In wheat, barley and fallow where up to 0.4 oz per acre of ALLY® EXTRA or DuPont CANVAS® XP has been applied, the following minimum crop rotation interval* can be followed:

Soil pH	Minimum precipitation between application and planting	Rotation interval in months*
7.3 or lower	10 inches	10
7.4 or higher	28 inches	34

* The rotation interval represents the period of time from the last application to the anticipated date of the next planting. Rotation intervals must be extended 1 crop season if drought conditions prevail after application and before the rotational crop is planted, unless a successful field bioassay has been completed (see the ALLY® EXTRA and/or CANVAS® XP label for details).

RESTRICTIONS

Wherever ALLY® EXTRA or CANVAS® XP is used on land previously treated with GLEAN® FC, FINESSE®, Amber¹, Assert², or other longer residual herbicides having the same mode of action, read the rotational guidelines on both labels and follow the one with the longest interval stated for your situation before choosing to rotate to crops other than wheat or barley.

IMPORTANT


BEFORE USING ALLY® EXTRA OR CANVAS® XP READ AND FOLLOW ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA-REGISTERED LABEL.

This bulletin contains new or supplemental instructions for use of this product which do not appear on the EPA-registered package label. Follow the instructions carefully.

This label must be in the possession of the user at the time of pesticide application.

¹ Registered trademark of Novartis

² Registered trademark of BASF



DR-138 020902

NEXT

LABEL

**DuPont Agricultural
Products**

DRAFT

**ALLY® EXTRA AND
CANVAS® XP
HERBICIDES**

"..... A Growing Partnership With Nature"

DUPONT ALLY® EXTRA HERBICIDE

EPA Reg. No. 352-610

DUPONT CANVAS® XP HERBICIDE ✓

EPA Reg. No. 352-610

REDUCED ROTATION INTERVAL FOR CONDIMENT MUSTARD AND CHICKPEAS (GARBANZO BEANS) IN THE STATES OF IDAHO, OREGON, AND WASHINGTON

DIRECTIONS FOR USE

It is violation of federal law to use this product in a manner inconsistent with its labeling.

DuPont ALLY® EXTRA Herbicide and DuPont CANVAS® XP Herbicide are selective postemergence herbicides recommended for control of broadleaf weeds in wheat, barley, and fallow.

HOW TO USE

In wheat, barley and fallow where up to 0.4 oz per acre of ALLY® EXTRA or DuPont CANVAS® XP has been applied, the following minimum crop rotation interval* can be followed:

Soil pH	Minimum precipitation between application and planting	Rotation interval in months*
7.3 or lower	10 inches	10
7.4 or higher	28 inches	34

* The rotation interval represents the period of time from the last application to the anticipated date of the next planting. Rotation intervals must be extended 1 crop season if drought conditions prevail after application and before the rotational crop is planted, unless a successful field bioassay has been completed (see the ALLY® EXTRA and/or CANVAS® XP label for details).

RESTRICTIONS

Wherever ALLY® EXTRA or CANVAS® XP is used on land previously treated with GLEAN® FC, FINESSE®, Amber¹, Assert², or other longer residual herbicides having the same mode of action, read the rotational guidelines on both labels and follow the one with the longest interval stated for your situation before choosing to rotate to crops other than wheat or barley.

IMPORTANT


BEFORE USING ALLY® EXTRA OR CANVAS® XP READ AND FOLLOW ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA-REGISTERED LABEL.

This bulletin contains new or supplemental instructions for use of this product which do not appear on the EPA-registered package label. Follow the instructions carefully.

This label must be in the possession of the user at the time of pesticide application.

¹Registered trademark of Novartis

²Registered trademark of BASF



DR-138 020902