

352-643

05-25-2010

1/9



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY  
Washington, D.C. 20460

OFFICE OF  
CHEMICAL SAFETY AND  
POLLUTION PREVENTION

MAY 25 2010

S.K. Theodorakis  
E.I. du Pont de Nemours and Company  
1007 Market Street  
Wilmington, DE 19898

Subject: Label Amendment (container disposal)  
DuPont TranXit Herbicide  
EPA Reg. No. 352-643  
Application Dated April 26, 2010

Dear Mr. Theodorakis:

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, is acceptable, provided you make the following changes:

1. Change the heading from "Inert Ingredients" to "**Other Ingredients**".
2. Change the Precautionary Statement to read "Causes **moderate** eye irritation."
3. Change the PPE phrase to read "If no such instructions for washables **exist**,"
4. Under User Safety Recommendations, the statement "Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing." should be added.
5. Change the Environmental Hazards sentence to read "Do not contaminate water **when disposing of equipment washwater or rinsate**."
6. Change the Directions for Use statement to read "It is a violation of Federal law to use this product in a manner inconsistent with **its labeling**."
7. Change the heading from "GENERAL INFORMATION" to "**PRODUCT INFORMATION**".
8. On page 3, change the heading from "PRECAUTIONS" to "**RESTRICTIONS**".
9. On pages 4 & 5, remove all instances of the heading "General Comments and Precautions".

- 10. Change the Pesticide Disposal sentence to read “**Wastes** resulting from the use of this product **must** be disposed of on site or at an approved waste disposal facility.”
- 11. Remove the warranty statement “DuPont does not agree to be an insurer of these risks.”
- 12. Add the phrase “**TO THE EXTENT CONSISTENT WITH APPLICABLE LAW**” directly in front of the following warranty statements:
  - “DUPONT MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY...”
  - “IN NO EVENT SHALL DUPONT OR SELLER BE LIABLE...”
  - “THE EXCLUSIVE REMEDY OF THE USER OR BUYER...”
  - “DuPont or its distributor must have prompt notice of any claim...”

A stamped copy of your label is enclosed for your records. This label supercedes all previously accepted labels. You must submit one (1) copy of the final printed label before you release the product for shipment. Products shipped after eighteen (18) months from the date of this letter or the next printing of the label, whichever occurs first, must bear the new revised label. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA §6(e). Your release for shipment of the product constitutes acceptance of these conditions.

Sincerely,

*Mindy Ondrich for*

Jim Tompkins  
Product Manager 25  
Herbicide Branch  
Registration Division (7505P)

# DuPont™ TranXit®

## HERBICIDE

### Professional Products

ACCEPTED  
with COMMENTS  
in EPA Letter Dated

MAY 25 2010

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No.

352-643

#### Dry flowable

For Postemergence Control of Weeds in Warm Season Turf.

For Use on Sodfarms\*, Seed Farms\*, Golf Courses, Professionally Managed College and Professional Sports Fields, Industrial and Commercial Lawns and Other Similar Nonresidential Areas.

Not for use in areas where children can contact treated turf.

\*Except in Arizona

Active Ingredient	By Weight
Rimsulfuron: N-((4,6-dimethoxypyrimidin-2-yl)aminocarbonyl)-3-(ethylsulfonyl)-2-pyridinesulfonamide	25.0%
Inert Ingredients	75.0%
TOTAL	100.0%

EPA Reg. No. 352-643      EPA Est. No. \_\_\_\_\_

Nonrefillable Container

Net: \_\_\_\_\_

OR

Refillable Container

Net: \_\_\_\_\_

### KEEP OUT OF REACH OF CHILDREN CAUTION FIRST AID

**IF IN EYES:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for further treatment advice.

**IF ON SKIN OR CLOTHING:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 to 20 minutes. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-441-3637 for emergency medical treatment information.

### PRECAUTIONARY STATEMENTS

#### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

**CAUTION!** Causes eye irritation. Harmful if absorbed through skin. Avoid contact with skin, eyes, or clothing. Avoid breathing dust or spray mist.

Some materials that are chemical-resistant to this product are listed below. If you want more options follow the instructions for Category A on an EPA chemical-resistance category selection chart.

#### PERSONAL PROTECTIVE EQUIPMENT (PPE)

*Applicators and other handlers must wear:*

Long-sleeved shirt and long pants.

Chemical-resistant gloves made of any waterproof material such as butyl rubber, natural rubber, or neoprene rubber ≥ 14 mils.

Shoes plus socks.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

#### USER SAFETY RECOMMENDATIONS

**USERS SHOULD:** Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.

#### ENVIRONMENTAL HAZARDS

Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water by cleaning of equipment or disposal of wastes.

## DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with the terms of this label.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency in your State responsible for pesticide regulation.

### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry intervals. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 4 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

Coveralls.

Chemical-resistant gloves made of any waterproof material, such as butyl rubber, natural rubber, or neoprene rubber  $\geq$  14 mils.

Shoes plus socks.

Sod farms are within the scope of the Worker Protection Standard.

### NON AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are not within the scope of the Worker Protection Standard for agricultural pesticides, 40 CFR part 170.

The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Professional applications to golf courses, industrial and commercial lawns and sports fields are not within the scope of the Worker Protection Standard.

Do not enter or allow others to enter the treated area until sprays have dried.

## GENERAL INFORMATION

DuPont™ TRANXIT® is noncorrosive to equipment, nonflammable, and nonvolatile. Control of weeds with TRANXIT® requires approximately 3 to 4 weeks, but weed growth ceases soon after application.

TRANXIT® temporarily suppresses growth of Tifway bermudagrass. This is a growth regulator

effect. Delayed green-up of bermudagrass in the spring has not been observed. This is typically followed by a flush of growth within 3 to 4 weeks. Some slight yellowing of bermudagrass may occur and last for approximately 7 days.

TRANXIT® is absorbed through the roots and foliage of plants, rapidly inhibiting the growth of susceptible weeds. Rainfall or sprinkler irrigation is needed to move TRANXIT® into the soil. In some cases, susceptible weeds may germinate and emerge a few days after application; but, growth then ceases and leaves become chlorotic 3 to 10 days after emergence. Death of leaf tissue and growing point will follow in some species; in other species, the seedling may remain green, but be stunted in growth and remain noncompetitive.

One to three weeks after postemergence application to weeds, leaves of susceptible plants appear chlorotic, and the growing point subsequently dies. In warm moist conditions, the expression of herbicide symptoms is accelerated. In cold dry conditions, expression of herbicide symptoms may be delayed. Death of leaf tissue and growing point will follow in some species; in other species, the seedling may remain green, but be stunted in growth and remain noncompetitive.

The herbicidal action of TRANXIT® may be less effective on susceptible species stressed from environmental conditions (such as extreme temperature or moisture), abnormal soil conditions (saturated or waterlogged soils) or cultural practices. In addition, weeds hardened off by drought stress are less susceptible to TRANXIT®. Under cold conditions, TRANXIT® activity is delayed and takes longer to control weeds. In order to achieve faster control in cold conditions, increase the rate to 2 ounces per acre.

Lateral movement and tracking have been observed in heavier clay soils which are characterized by low water infiltration rates. These problems can be significantly reduced by use of short, frequent irrigation cycles. A minimum of three irrigation cycles is recommended in order to move TRANXIT® from the turf and weed canopy into the soil. These irrigation cycles should be applied prior to allowing foot or equipment traffic into treated areas. The use of a soil wetting agent may be beneficial in both clay and sandy soils. Where slopes are severe, mechanical aerification prior to spraying may help water penetrate into the soil and move TRANXIT® into the soil profile.

Never depend solely upon rainfall to move TRANXIT® into the soil. Rainfall is unpredictable and, if heavy enough, will move TRANXIT® laterally. Some foliar uptake may have to be sacrificed when foot and/or equipment traffic is imminent soon after TRANXIT® application. Begin irrigation regime 1 hour following TRANXIT® application.

Extreme caution should be used when applying this product to slopes of heavy or clay soils that drain onto bentgrass greens, overseeded greens, fairways, or tees. Do not apply to slopes that drain directly onto soil or "push up" bentgrass greens. Tracking and lateral movement onto bentgrass putting greens rarely results in death to bentgrass. Some off color and growth

9/4

5/9

regulatory effects have been observed for approximately 14 days. This effect may last longer on non-USGA (sand greens) or during periods of cool weather or on greens maintained at low nitrogen levels. Fertilization with a liquid fertilizer can help speed bentgrass recovery. The addition of a product containing gibberellic acid may also be beneficial. Application of activated charcoal around and/or on the perimeter of bentgrass putting greens has also been effective in reducing injury potential.

**PRECAUTIONS**

- Do not apply to residential lawns.
- Do not apply to areas where children can contact treated turf.
- Do not apply an organophosphate insecticide or nematicide within 7 days of a DuPont™ TRANXIT® application as injury potential to the desired grass may increase.
- Do not apply to newly sprigged or sodded bermudagrass.
- Do not apply if wind speed becomes excessive; spray drift can occur at wind speeds greater than 10 mph. If sensitive species are downwind, extreme caution must be used. If conditions for spray drift exist, use a spray shield.
- Do not apply if winds are gusty.

**CHEMIGATION**

Do not apply this product through any type of irrigation system.

**SPRAY DRIFT MANAGEMENT**

The interaction of many equipment and weather related factors determine the potential for spray drift. The applicator is responsible for considering all these factors when making application decisions.

**AVOIDING SPRAY DRIFT IS THE RESPONSIBILITY OF THE APPLICATOR.**

**IMPORTANCE OF DROPLET SIZE**

The most effective way to reduce drift potential is to apply large droplets (>150 – 200 microns). The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. The presence of sensitive species nearby, the environmental conditions, and pest pressure may affect how an applicator balances drift control and coverage. **APPLYING LARGER DROPLETS REDUCES DRIFT POTENTIAL, BUT WILL NOT PREVENT DRIFT IF APPLICATIONS ARE MADE IMPROPERLY OR UNDER UNFAVORABLE ENVIRONMENTAL CONDITIONS.** See Wind, Temperature and Humidity, and Temperature Inversions sections of this label.

**CONTROLLING DROPLET SIZE**

- Volume - Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets.
- Pressure - Do not exceed the nozzle manufacturers recommended pressures. For many nozzle types, lower pressure produces larger droplets. When higher flow rates are needed, use higher flow rate nozzles instead of increasing pressure.

- Number of nozzles - Use the minimum number of nozzles that provide uniform coverage.
- Nozzle Orientation - Orienting nozzles so that the spray is released parallel to the airstream produces larger droplets than other orientations and is the recommended practice. Significant deflection from horizontal will reduce droplet size and increase drift potential.
- Nozzle Type - Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles. Solid stream nozzles oriented straight back produce the largest droplets and the lowest drift.

**APPLICATION HEIGHT**

Make applications at the lowest height that is safe in order to reduce exposure of droplets to evaporation and wind.

**SWATH ADJUSTMENT**

When applications are made with a crosswind, the swath will be displaced downwind. Swath adjustment distance should increase, with increasing drift potential (higher wind, smaller drops, etc.)

**WIND**

Drift potential is lowest between wind speeds of 2-10 mph. However, many factors, including droplet size and equipment type determine drift potential at any given speed. Application should be avoided below 2 mph due to variable wind direction and high inversion potential. If wind speeds are above 10 mph, a spray shield is recommended. NOTE: Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they affect spray drift.

**TEMPERATURE AND HUMIDITY**

When making applications in low relative humidity, set up equipment to produce larger droplets to compensate for evaporation. Droplet evaporation is most severe when conditions are both hot and dry.

**TEMPERATURE INVERSIONS**

Applications should not occur during a temperature inversion because drift potential is high. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain in a concentrated cloud. This cloud can move in unpredictable directions due to the light variable winds common during inversions. Temperature inversions are characterized by increasing temperatures with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

**SENSITIVE AREAS**

The pesticide should only be applied when the potential for drift to adjacent sensitive areas (e.g. residential areas, bodies of water, known habitat for threatened or endangered species, non-target crops) is minimal (e.g. when wind is blowing away from the sensitive areas.)

**SPRAYER CLEANUP**

To avoid subsequent injury to sensitive species, thoroughly clean all mixing and spray equipment immediately after application of DuPont™ TRANXIT® as follows:

- 1) Drain tank and thoroughly rinse spray tank, boom and hoses with clean water. Loosen and physically remove any visible deposits.
- 2) Fill the tank with clean water and 1 gallon of household ammonia for every 100 gallons of water. Circulate the cleaning solution through the pump and bypass for several minutes. Turn on boom(s) and spray for several minutes. Drain the cleaning solution.
- 3) Remove the nozzle tips and screens, and clean separately in a bucket containing household ammonia and water. Rinse with water and replace screens and nozzle tips in the boom(s).
- 4) Repeat Step 2.
- 5) Repeat Step 2 with water only.

**USE DIRECTIONS:**

**I. Control of Annual Bluegrass Infesting Non-Overseeded Bermudagrass on Sod Farms\*, Seed Farms\*, Golf Courses, Professionally Managed College and Professional Sports Fields, Industrial & Commercial Lawns and Other Similar Nonresidential Areas (not including areas where children can contact treated turf)**

**\*Except in Arizona**

**General Comments and Precautions:** Do not allow spray droplet drift to contact desirable ryegrasses, bentgrasses, bluegrasses, fescues, or ornamental shrubs, trees, or flowers.

Annual bluegrass (*Poa annua* var. *annua*) is controlled with TRANXIT®. Degree of control for the perennial (*Poa annua* var. *reptans*) has not been fully determined. The perennial biotypes/ecotypes are primarily confined to bentgrass putting greens where TRANXIT® cannot be used.

**Non-putting green bermudagrass not overseeded:** Apply in November/December and again in February/March if necessary at a rate of 1 to 2 ounces TRANXIT®/acre. Add a nonionic surfactant at 0.25% v/v. Apply in 15 to 60 gallons of water per acre. Use the higher rate on areas with large plants and high annual bluegrass populations.

**Putting green bermudagrass not overseeded:** Apply at a rate of 1 ounce TRANXIT®/acre to annual bluegrass that is no larger than one inch in diameter. Bermudagrass that is not fully dormant may show some yellowing. Repeat applications as needed on a 3 week schedule, but the number of applications must not exceed a total of 4.

**II. Removal of Perennial Ryegrass and *Poa trivialis* (Rough Bluegrass) from Bermudagrass on Sod Farms\*, Seed Farms\*, Golf Courses, Professionally Managed College and Professional Sports Fields, Industrial & Commercial Lawns and Other Similar Nonresidential Areas (not including areas where children can contact treated turf)**

**\*Except in Arizona**

**General Precautions and Comments:** Perennial ryegrass is more sensitive to TRANXIT® than *Poa trivialis* (Rough Bluegrass).

Repeat applications made at lower rates are more efficacious than a single large application.

Perennial ryegrass and *Poa trivialis* growing in high density seedlings and maintained at low mowing heights are more sensitive to TRANXIT® than individual plants growing as unmowed clumps.

Putting green bermudagrass cultivars that have shown tolerance to TRANXIT® include: Tifdwarf, TifEagle, Floradwarf, Mississippi Supreme, and Champion.

**Non-putting green bermudagrass:** Apply 0.5 to 2 ounces TRANXIT®/acre in the spring months 3 to 4 weeks before desired date for overseed removal. Add a nonionic surfactant at 0.25% v/v. Apply in 15 to 60 gallons of water per acre. The higher rate and repeat applications may be necessary for complete removal of *Poa trivialis*.

**Putting green bermudagrass:** Apply 0.5 to 1 ounce of TRANXIT®/acre 3 weeks before desired date for overseed removal. Add a nonionic surfactant at 0.25% v/v. Apply in 15 to 60 gallons of water per acre. The lower rate of TRANXIT® can be used for slower transition. Three applications of the 0.5 ounce/acre rate or 2 applications of the 1 ounce/acre rate are generally required for removal of *Poa trivialis*.

**III. Annual bluegrass Control Prior to Overseeding Bermudagrass on Sod Farms\*, Seed Farms\*, Golf Courses, Professionally Managed College and Professional Sports Fields, Industrial & Commercial Lawns and Other Similar Nonresidential Areas with Perennial Ryegrass or *Poa trivialis* (not including areas where children can contact treated turf)**

**\*Except in Arizona**

**General Comments and Precautions:** Annual bluegrass control will be reduced where "early overseeding" is practiced. Also, applying TRANXIT® earlier than 2 weeks prior to normal overseeding times will result in reduced annual bluegrass control.

Avoid mechanical disturbance (aeration, verti-cutting, etc.) of the turf, thatch and/or soil layer after TRANXIT® application or annual bluegrass control may be reduced.

Avoid application to wet and/or waterlogged putting greens. Allow at least 72 hours for drying on waterlogged putting greens before applying TRANXIT®.

Stressed bermudagrass turf growing in shaded areas, waterlogged soil, or under other environmental stress (such as nematodes) may exhibit more discoloration or chlorosis following application of DuPont™ TRANXIT®.

**DO NOT APPLY AFTER OVERSEEDING, EXCEPT AS DIRECTED FOR REMOVAL OF OVERSEEDED GRASSES.**

**Non-putting green bermudagrass:** Apply 1 to 2 ounces of TRANXIT®/acre 10 to 14 days prior to overseeding perennial ryegrass and/or Poa trivialis. Add a nonionic surfactant at 0.25% v/v.

Apply in 15 to 60 gallons of water per acre. Occasional stunting of the overseeded perennial ryegrass and/or Poa trivialis may occur, but symptoms disappear in approximately 7 days. Cultural practices and favorable environmental conditions that allow for maximum germination of annual bluegrass prior to application of TRANXIT® increase the degree of control.

Apply 3 to 5 light irrigation cycles approximately 2 to 4 hours after application of TRANXIT® to dislodge TRANXIT® from the turf canopy. Apply enough irrigation water to penetrate the soil but do not allow the water to sheet or move lateral onto sensitive species.

**Putting green Bermudagrass:**

Apply 2 ounces of TRANXIT®/acre without an adjuvant 10 to 14 days prior to overseeding golf putting greens for control of emerged annual bluegrass. Irrigate within 2 to 4 hours of application and continue routine irrigation schedule. For putting greens growing in the shade, under waterlogged conditions, or other environmental stresses, such as nematodes, and to help reduce potential of bermudagrass injury, apply 1 ounce of TRANXIT® 3 weeks prior to overseeding and apply another ounce 1 week prior to overseeding.

**IV. Weed Control in Centipedegrass and Zoysiagrass on Sod Farms\*, Seed Farms\*, Golf Courses, Professionally Managed College and Professional Sports Fields, Industrial & Commercial Lawns and Other Similar Nonresidential Areas (not including areas where children can contact treated turf).**

**\*Except in Arizona**

**General Comments and Precautions:** Emerald, Zenith and Meyer zoysiagrass have shown tolerance to TRANXIT® similar to that of bermudagrass. Centipedegrass have exhibited moderate tolerance.

Tolerance of the majority of cultivars of these turfgrasses to TRANXIT® have not been fully investigated. The effects of TRANXIT® on these turfgrasses during transition has not been fully evaluated.

**Dormant and non-dormant turf:** Apply 1 to 2 ounces of TRANXIT®/acre for weed control. Some chlorosis of the turf may occur following application. Use the lower rate with repeat applications for weed control in centipedegrass.

**Cool-season weed species controlled:** Annual bluegrass, blue-eyed grass, wild carrots, little barley, ryegrass, tall fescue, rough bluegrass, common chickweed, shepherd's purse, cutleaf evening primrose, henbit and field pansy.

**Warm-season species controlled:** Spotted spurge, dollarweed, and suppression of some sedge species.

**STORAGE AND DISPOSAL**

**PESTICIDE STORAGE:** Store product in original container only. Do not contaminate water, other pesticides, fertilizer, food or feed in storage.

**PESTICIDE DISPOSAL:** Do not contaminate water, food or feed by disposal. Waste resulting from the use of this product may be disposed of on-site or at an approved waste disposal facility.

**CONTAINER HANDLING:** Refer to the Net Contents section of this product's labeling for the applicable "Nonrefillable Container" or "Refillable Container" designation.

**Nonrefillable Plastic and Metal Containers**

**(Capacity Equal to or Less Than 50 Pounds):**

Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then, for Plastic Containers, offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration. Do not burn, unless allowed by state and local ordinances. For Metal Containers, offer for recycling if available or reconditioning if appropriate, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities

**Nonrefillable Plastic and Metal Containers**

**(Capacity Greater Than 50 Pounds):**

Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then, for Plastic Containers, offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration. Do not burn, unless allowed by state and local ordinances. For Metal Containers, offer for recycling if available or reconditioning if appropriate, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

**Nonrefillable Plastic and Metal Containers, e.g., Intermediate Bulk Containers [IBC] (Size or Shape Too Large to be Tipped, Rolled or Turned Upside Down):** Nonrefillable container. Do not reuse or refill this container. Clean container promptly after emptying the contents from this container into application equipment or mix tank and before final disposal using the following pressure rinsing procedure. Insert a lance fitted with a suitable tank cleaning nozzle into the container and ensure that the water spray thoroughly covers the top, bottom and all sides inside the container. The nozzle manufacturer generally provides instructions for the appropriate spray pressure, spray duration and/or spray volume. If the manufacturer's instructions are not available, pressure rinse the container for at least 60 seconds using a minimum pressure of 30 PSI with a minimum rinse volume of 10% of the container volume. Drain, pour or pump rinsate into application equipment or rinsate collection system. Repeat this pressure rinsing procedure two more times. Then, for Plastic Containers, offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration. For Metal Containers, offer for recycling if available or reconditioning if appropriate, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

**Nonrefillable Paper or Plastic Bags, Fiber Sacks including Flexible Intermediate Bulk Containers (FIBC) or Fiber Drums With Liners:** Nonrefillable container. Do not reuse or refill this container. Completely empty paper or plastic bag, fiber sack or drum liner by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application or manufacturing equipment. Then offer for recycling if available or dispose of empty paper or plastic bag, fiber sack or fiber drum and liner in a sanitary landfill, or by incineration. Do not burn, unless allowed by state and local ordinances.

**Refillable Fiber Drums With Liners: Refillable container (fiber drum only).** *Refilling Fiber Drum:* Refill this fiber drum with DuPont™ TRANXIT® containing rimsulfuron only. Do not reuse this fiber drum for any other purpose. Cleaning before refilling is the responsibility of the refiller. Completely empty liner by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application or manufacturing equipment. *Disposing of Fiber Drum and/or Liner:* Do not reuse this fiber drum for any other purpose other than refilling (see preceding). Cleaning the container (liner and/or fiber drum) before final disposal is the responsibility of the person disposing of the container. Offer the liner for recycling if available or dispose of liner in a sanitary landfill, or by incineration. Do not burn, unless allowed by state and local ordinances. If drum is contaminated and cannot be reused, dispose of it in the manner required for its liner. To clean the fiber drum before final disposal, completely empty the fiber drum by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application or manufacturing equipment. Then offer the fiber drum for recycling if available or dispose of in a sanitary landfill, or by incineration. Do not burn, unless allowed by state and local ordinances.

**All Other Refillable Containers:** Refillable container. *Refilling Container:* Refill this container with TRANXIT® containing rimsulfuron only. Do not reuse this container for any other purpose. Cleaning before refilling is the responsibility of the refiller. Prior to refilling, inspect carefully for damage such as cracks, punctures, abrasions, worn out threads and closure devices. If damage is found, do not use the container, contact DuPont at the number below for instructions. Check for leaks after refilling and before transporting. If leaks are found, do not reuse or transport container, contact DuPont at the number below for instructions. *Disposing of Container:* Do not reuse this container for any other purpose other than refilling (see preceding). Cleaning the container before final disposal is the responsibility of the person disposing of the container. To clean the container before final disposal, use the following pressure rinsing procedure. Insert a lance fitted with a suitable tank cleaning nozzle into the container and ensure that the water spray thoroughly covers the top, bottom and all sides inside the container. The nozzle manufacturer generally provides instructions for the appropriate spray pressure, spray duration and/or spray volume. If the manufacturer's instructions are not available, pressure rinse the container for at least 60 seconds using a minimum pressure of 30 PSI with a minimum rinse volume of 10% of the container volume. Drain, pour or pump rinsate into application equipment or rinsate collection system. Repeat this pressure rinsing procedure two more times. Then, for Plastic Containers, offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration. Do not burn, unless allowed by state and local ordinances. For Metal Containers, offer for recycling if available or reconditioning if appropriate, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

**Outer Foil Pouches of Water Soluble Packets (WSP):** Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available or, dispose of the empty outer foil pouch in the trash as long as WSP is unbroken. If the outer pouch contacts the formulated product in any way, the pouch must be triple rinsed with clean water. Add the rinsate to the spray tank and dispose of the outer pouch as described previously.

Do not transport if this container is damaged or leaking. If the container is damaged, leaking or obsolete, or in the event of a major spill, fire or other emergency, contact DuPont at 1-800-441-3637, day or night.

**NOTICE TO BUYER:** Purchase of this material does not confer any rights under patents of countries outside of the United States.

The DuPont Oval Logo, DuPont™ and TRANXIT® are trademarks or registered trademarks of E. I. duPont de Nemours and Company

D - 1557 032410

8/1/8

**LIMITATION OF  
WARRANTY AND LIABILITY**

**NOTICE:** Read This Limitation of Warranty and Liability Before Buying or Using This Product. If the Terms Are Not Acceptable, Return the Product at Once, Unopened, and the Purchase Price Will Be Refunded.

It is impossible to eliminate all risks associated with the use of this product. Such risks may arise from factors such as weather conditions, soil factors, off target movement, unconventional technique, presence of other materials, the manner of use or application, or other unknown factors, all of which are beyond the control of DuPont or Seller. These risks can cause: ineffectiveness of the product or other unintended consequences.

DuPont does not agree to be an insurer of these risks. **WHEN YOU BUY OR USE THIS PRODUCT, YOU AGREE TO ACCEPT THESE RISKS.**

DuPont warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for the purpose stated in the Directions for Use, subject to the inherent risks described above, when used in accordance with the Directions for Use under normal conditions.

**DUPONT MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR OF MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.**

**IN NO EVENT SHALL DUPONT OR SELLER BE LIABLE FOR ANY INCIDENTAL, CONSEQUENTIAL OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. THE EXCLUSIVE REMEDY OF THE USER OR BUYER AND THE EXCLUSIVE LIABILITY OF DUPONT OR SELLER, FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY OR CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHER LEGAL THEORY) RESULTING FROM THE STORAGE, USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT, OR AT THE ELECTION OF DUPONT OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

DuPont or its distributor must have prompt notice of any claim so that an immediate investigation of buyer's or user's claim can be made. Buyer and all users shall promptly notify DuPont or a DuPont distributor of any claims, whether based on contract, negligence, strict liability, other tort or otherwise or be barred from any remedy.

This Limitation of Warranty and Liability may not be amended by any oral or written agreement.

**For product information call: 1-888-6-DUPONT**

**© 2005 E. I. du Pont de Nemours and Company, Wilmington, Delaware 19898.**

**All rights reserved.**