

352-650

02/09/2005

1/18



U.S. ENVIRONMENTAL PROTECTION AGENCY
Office of Pesticide Programs
Registration Division (H7505C)
401 "M" St., S.W.
Washington, D.C. 20460

EPA Reg. Number:
352-650

Date of Issuance:
FEB - 9 2005

NOTICE OF PESTICIDE:
 Registration
 Reregistration

Term of Issuance:
Conditional

Name of Pesticide Product:
DuPont Synchrony XP
(mp) Herbicide

(under FIFRA, as amended)

Name and Address of Registrant (include ZIP Code):

E.I. DuPont de Nemours & Company
DuPont Crop Protection
Stine-Haskell Research Center
P.O. Box 30
Newark, DE 19714-0030

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA section 3(c)(7)(A) provided that you:

1. Submit/cite all data required for registration/reregistration of your product when the Agency requires all registrants of similar products to submit such data.
2. Add the phrase "EPA Registration No.352-650" before you release the product for shipment.
3. Submit two (2) copies of your final printed labeling before you release the product for shipment.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of this product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Signature of Approving Official:

Date:

Walter R. ...

2/9/05



DuPont™
Synchrony® XP (mp)
herbicide

DRAFT LABEL



“..... A Growing Partnership With Nature”

DUPONT™ SYNCHRONY® XP (mp)

Highlights

- SYNCHRONY® XP (mp) is a unit pack product that provides selective postemergence weed control of a large spectrum of broadleaf weeds.
- SYNCHRONY® XP (mp) may be tank mixed with DuPont™ ASSURE® II, or other products for increased weed control.
- Include a crop oil concentrate or a nonionic surfactant and ammonium-based nitrogen fertilizer where required.
- SYNCHRONY® XP (mp) may be applied by ground (broadcast or band) or by air.
- For ground application, apply in 10-25 gallons of water at 25-60 psi. Use flat fan nozzles to optimize SYNCHRONY® XP (mp) performance.
- Apply to actively growing weeds at the recommended sizes. See Rate.
- Certain crop rotations apply. See Rotational Crop Guidelines.
- Certain environmental conditions, such as cool and dry, or hot and humid weather, affect the performance of SYNCHRONY® XP (mp). See Environmental Conditions.
- Consult label text for complete instructions. Always read and follow label directions for use.

TABLE OF CONTENTS

Directions For Use 2

Application Information - All Uses..... 2

 Timing To Crop Stage 2

 Timing To Weeds 2

 Tank Mix Restrictions 2

 Cultivation 2

SYNCHRONY® XP (mp) at 0.75 oz per acre - in season use on "STS" soybeans only

 Rate and Weeds Controlled 3

 Sequential Applications 3

 Spray Additives 3

 Tank Mixes for 0.75 oz per acre in season use..... 3

SYNCHRONY® XP (mp) at 0.75 oz per acre - Burndown Application

 Application Timing 4

 Spray Additives 4

 Weeds Controlled - Burndown 4

 Weeds Controlled - Limited Residual 4

SYNCHRONY® XP (mp) at 0.375 oz per acre - Reduced Rate Uses

 Postemergence Application Information 5

 Timing to Weeds..... 5

 Spray Additives..... 5

 - "STS" soybeans 5

 - Non-"STS" soybeans..... 5

 Weeds Controlled by 0.375 oz per acre SYNCHRONY® XP (mp)..... 5

 Tank Mixes with 0.375 oz per acre SYNCHRONY® XP (mp)..... 5

 Weeds Controlled by 0.375 oz per acre SYNCHRONY® XP (mp) preceded by DuPont™ AUTHORITY®..... 5

 Burndown tankmix with AUTHORITY®..... 5

 Precautions for 0.375 oz per acre SYNCHRONY® XP (mp) 5

 Mixing Instructions 5

 Application Equipment 6

 Ground Application 6

 Aerial Application 6

 Environmental Conditions and Biological Activity 6

Rotational Crop Guidelines 6

 The Importance of Soil pH..... 8

 Sprayer Preparation and Cleanup 8

 Spray Drift Management 8

 Important Precautions 9

 Information on Resistant Weeds 9

 Integrated Pest Management 10

 Storage and Disposal 10

 Notice of Warranty 11

4/18



DuPont™

Synchrony® XP (mp)


herbicide

A unit pack product for use on soybeans.

Dispersible Granules

Active Ingredients	By Weight
Chlorimuron ethyl Ethyl 2-[[[(4-chloro-6-methoxypyrimidin-2-yl)amino]carbonyl]amino]sulfonyl]benzoate	21.5%
Thifensulfuron methyl Methyl 3-[[[(4-methoxy-6-methyl-1,3,5-triazin-2-yl)amino]carbonyl]amino]sulfonyl]-2-thiophenecarboxylate	6.9%
Inert Ingredients	71.6%
TOTAL	100.0%

EPA Reg. No. 352 - XXX
EPA Est. No. _____
Net Weight: 3.75 ounces



KEEP OUT OF REACH OF CHILDREN

CAUTION

FIRST AID

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-441-3637 for emergency medical treatment information.

PRECAUTIONARY STATEMENTS

CAUTION

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION! Causes moderate eye irritation. Harmful if absorbed through skin.

Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling.

PERSONAL PROTECTIVE EQUIPMENT

Some materials that are chemical resistant to this product are listed below. If you want more options follow the instructions for Category A on an EPA chemical resistance category selection chart.

Applicators and other handlers must wear:

Long-sleeved shirt and long pants.

Chemical Resistant Gloves Category A (such as butyl rubber, natural rubber, neoprene rubber, or nitrile rubber), all ≥14 mls.

Shoes plus socks.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Control Statement: When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

Important: When reduced PPE is worn because a closed system is being used, handlers must be provided all PPE specified above for "Applicators and Other Handlers" and have such PPE immediately available for use in an emergency, such as a spill or equipment breakdown.

USER SAFETY RECOMMENDATIONS

USERS SHOULD: Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment washwaters. Do not apply where/when conditions favor runoff.

GENERAL INFORMATION

DuPont™ SYNCHRONY® XP (mp) is a convenient unit pack that is mixed in water and applied as a spray. Open package completely and empty contents of both compartments into spray tank.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

Coveralls.

Chemical Resistant Gloves Category A (such as butyl rubber, natural rubber, neoprene rubber, or nitrile rubber), all ≥14 mls.

Shoes plus socks.

Use only in the geographies identified in the "Rotational Crop Guidelines" section of this label.

SYNCHRONY® XP (mp) herbicide must be used only in accordance with recommendations on this label or in separately published DuPont recommendations.

Do not apply this product through any type of irrigation system.

DuPont will not be responsible for losses or damages resulting from the use of this product in any manner not specifically recommended by DuPont.

APPLICATION INFORMATION - ALL USES

- SYNCHRONY® XP (mp) herbicide is a dispersible granule formulation which readily dissolves in water.
- SYNCHRONY® XP (mp) may be used in conventional, no-till, or conservation tillage soybean production.

Timing To Crop Stage

- SYNCHRONY® XP (mp), for in-season use, may be applied any time after the first trifoliolate has opened but no later than 60 days before soybean maturity.

- SYNCHRONY® XP (mp), for burndown use, may be applied up to 30 days before planting, or prior to soybean emergence after planting.

Timing To Weeds

- Apply SYNCHRONY® XP (mp) when weeds are young and actively growing (after the first true leaves have expanded, but before the weeds exceed the size indicated below).
- Applications made to weeds larger than the sizes indicated below, or to weeds under stress, may result in unsatisfactory control.
- Applications made after July 15 may result in less than satisfactory weed control due to the large sizes of the weeds and increased weather stress.

Tank Mix Restrictions

When tank mixing SYNCHRONY® XP (mp) with any other approved soybean pesticide, always read and follow all use directions, restrictions and precautions of both SYNCHRONY® XP (mp) and the tank mix partner(s). When tank mixing, the most restrictive labeling applies.

Weed control and crop safety resulting from the use of tank mixtures not specifically noted on this label, or in separately published DuPont recommendations, are the responsibility of the user.

Cultivation

Do not cultivate within 7 days of application. Cultivation may put weeds under stress by pruning roots, thus diminishing control.

Cultivation approximately 14 days after application will help control suppressed weeds.

SYNCHRONY® XP (MP) AT 0.75 - 1.125 OZ PER ACRE - IN SEASON USE ON "STS" SOYBEANS ONLY

FOR USE ONLY ON SOYBEAN VARIETIES DESIGNATED AS "STS"

The normal use rate of SYNCHRONY® XP (mp) on STS soybeans is 0.75 oz oz per acre.

SYNCHRONY® XP (mp) may be used at 1.125 oz per acre for preserving the STS trait in STS soybean seed production. Consult ROTATIONAL CROP GUIDELINES for recrop intervals when using the 1.125 oz application rate.

- Application to soybean varieties not designated as "STS" will result in severe crop injury and/or yield loss.
- DuPont will not warrant the safety of this treatment to seed saved from previous year's production (bin run seed).
- These "STS" varieties must be purchased from an authorized seed supplier.
- The "STS" designation indicates the soybean variety contains a proprietary trait that enhances the soybean's natural tolerance to DuPont soybean sulfonyleurea herbicides. Information on "STS" soybean varieties may be obtained from your seed supplier or DuPont representative.

6/18

DuPont™ SYNCHRONY® XP (mp) is a unit pack product which will treat up to 5 acres of soybeans as a broadcast application. Refer to the following table for the number of acres treated by 0.75 or 1.125 ounces per acre of SYNCHRONY® XP (mp).

	One (1) 3.75 oz. Unit Pack Will Treat:	Three (3) 3.75 oz. Unit Packs Will Treat:	Five (5) 3.75 oz. Unit Packs Will Treat:
SYNCHRONY® XP (mp)			
Application Rate			
0.75 oz/A	5 acres	15 acres	25 acres
1.125 oz/A	3 1/3 acres	10 acres	16 2/3 acres

Weeds Controlled at 0.75 oz/acre rate in season

Apply SYNCHRONY® XP (mp) at a rate of 0.75 ounce per acre for selective postemergence control of the following broadleaf weeds:

Weed	Maximum Height (in inches)
Beggarticks (Bidens sp)	6
Bristly Starbur	3
Burcucumber*	3
Cocklebur	8
Common Milkweed (above ground portion)	6
Cowpea	5
Dandelion (above ground portion)	4
Florida Beggarweed	5
Hemp Sesbania	5
Lambsquarters	4
Jerusalem Artichoke (above ground portion)	6
Jimsonweed	5
Marestail	5
Morningglory* (annual)	
Entireleaf	3
Ivyleaf	3
Pitted	3
Smallflower	3
Tall	3
Mustard	5**
Pigweeds	
Redroot (Rough)	8
Others	8
Ragweed	
Common	4
Giant*	4
Sicklepod*	3
Smartweed	
Ladysthumb	8
Pennsylvania	8
Sunflower	8
Wild Poinsettia	2
Yellow Nutsedge	3
Velvetleaf	8

* See Sequential Applications section.

** Diameter

When applied as directed, SYNCHRONY® XP (mp) will suppress the following weeds:

Weed	Maximum Height (in inches)
Buffalobur	6
Kochia	3
Pokeweed	6
Purple Nutsedge	4
Sowthistle, perennial (rosette stage)	6
Spurred Anoda	3
Thistle, Canadian	4
Thistle, Russian	3
Venice Mallow	3

Sequential Applications: 0.75 oz/acre rate in season

A follow up application of DuPont™ CLASSIC® herbicide or SYNCHRONY® XP (mp) herbicide may be made 2-3 weeks after a SYNCHRONY® XP (mp) application to control weeds with multiple germination flushes or weeds under stress such as burcucumber, cowpea, giant ragweed, morningglory, and sicklepod. See Rotational Crop Guidelines for intervals following sequential applications.

Spray Additives: 0.75 oz/acre rate in season

Applications of SYNCHRONY® XP (MP) must include either a crop oil concentrate or a nonionic surfactant. Crop oil concentrate is the required adjuvant system unless tank mixing with a product that precludes use of crop oil concentrate. An ammonium nitrogen fertilizer may also be required. Products that combine ammonium fertilizers with surfactants or crop oils must meet all of the surfactant/crop oil and ammonium nitrogen fertilizer requirements.

Consult local DuPont fact sheets, technical bulletins, and service policies prior to using other adjuvant systems. If another herbicide is tank mixed with SYNCHRONY® XP (mp), select adjuvants authorized for use with both products. Adjuvants must contain only EPA-exempt ingredients (40 CFR 1001).

Crop Oil Concentrate (COC) - Petroleum or Modified Seed Oil (MSO)

- Apply at 1% v/v (1 gal per 100 gal spray solution) or 2% under arid conditions.
- Oil adjuvants must contain at least 80% high quality, petroleum (mineral) or modified vegetable seed oil with at least 15% surfactant emulsifiers.

Nonionic Surfactant (NIS)

- Apply at 0.25% v/v (1 qt per 100 gal spray solution) or 0.5% under arid conditions.
- Surfactant products must contain at least 60% nonionic surfactant with a hydrophilic/lipophilic balance (HLB) greater than 12.

Ammonium Nitrogen Fertilizer

- Use 2 qt/acre of a high-quality urea ammonium nitrate (UAN), such as 28%N or 32%N, or 2 lb/acre of a spray grade ammonium sulfate (AMS). Use 4 qt/acre UAN or 4 lb/acre AMS under arid conditions.
- Do not use liquid nitrogen fertilizer as the total carrier solution.

Special Adjuvant Types

- Combination adjuvant products may be used at doses that provide the required amount of NIS, COC, MSO and/or ammonium nitrogen fertilizer. Consult product literature for use rates and restrictions.
- In addition to the adjuvants specified above, other adjuvant types may be used if they provide the same functionality and have been evaluated and approved by DuPont Product Management. Consult separate DuPont technical bulletins for detailed information before using adjuvant types not specified on this label.

DUPONT™ SYNCHRONY® XP (MP) TANK MIXTURES

Other than the exceptions noted, and in addition to the tank mix partners and rates indicated below, SYNCHRONY® XP (mp) may be tank mixed or followed with sequential applications of other products registered for use in soybeans. DuPont™ SYNCHRONY® XP (mp) may be applied in tank mix combinations with full or reduced rates of other products provided:

- The tank mix product is labeled for the same timing, method of application, adjuvants, and use restrictions as SYNCHRONY® XP (mp).
- The tank mixture is not specifically prohibited on the label of the tank mix product.
- The tank mix combination is compatible as determined by a "jar test" described in the TANK MIX COMPATIBILITY TESTING section below.

Tank Mixes for 0.75 - 1.125 oz per acre rate in season

SYNCHRONY® XP and glyphosate products on RR/STS stacked-trait soybeans:

- SYNCHRONY® XP (mp) at 0.75 oz per acre may be tank mixed with glyphosate-containing herbicides for improved control of broadleaf weeds not controlled by either product alone.
- SYNCHRONY® XP (mp) at 1.125 oz per acre may be tank mixed with glyphosate-containing herbicides for preserving the STS trait in RR/STS stacked-trait soybean seed production fields.
- Always include 0.25% v/v non-ionic surfactant. The addition of 1 - 2 percent dry ammonium sulfate by weight or 8.5 - 17 pounds per 100 gallons of water may increase performance of these tank mixtures.

SYNCHRONY® XP (mp) and "Flexstar", "Reflex", "Blazer", or "Cobra" Herbicides:

For control of up to 4" waterhemp, up to 2" eastern black nightshade and for improved common ragweed control, SYNCHRONY® XP (mp) may be tankmixed with:

- 0.75 - 1.25 pt/acre "Flexstar"
- 0.75 - 1.5 pt/acre "Reflex"
- 0.5 - 1.5 pt/acre "Blazer", or
- 4 - 6 fluid oz/acre "Cobra"

Refer to the "Flexstar", "Reflex", "Blazer" and "Cobra" labels for the appropriate rate based on the weed sizes to be controlled. Nonionic surfactant or crop oil concentrate must be added.

- **For best results with SYNCHRONY® XP (mp) plus "Reflex" or "Flexstar"**, use a methylated seed oil-based or petroleum oil-based crop oil concentrate at 8 pt per 100 gallon spray solution (1% v/v).

- **For best results with SYNCHRONY® XP (mp) plus "Blazer"**, use nonionic surfactant at 2 pt per 100 gallon spray solution. Use of crop oil concentrate is not recommended, as severe injury may occur.
- **For best results with SYNCHRONY® XP (mp) plus "Cobra"**, use crop oil concentrate at 4 pt per 100 gallon spray solution (0.5% v/v).

For control of Prickly Sida and Hemp Sesbania, tank mix 0.75 oz SYNCHRONY® XP (mp) with 8-12.5 fl oz "Cobra".

Use the higher "Cobra" rate when Prickly Sida or Hemp Sesbania are heavy or if Prickly Sida and Hemp Sesbania approach the maximum size of 1" or 4", respectively. Include a nonionic surfactant at 1-2 pt per 100 gallons of spray solution (.125-.25% v/v). Do not use crop oil concentrate when tankmixing SYNCHRONY® XP (mp) and "Cobra" at these rates.

- **Precautions for tankmixes of SYNCHRONY® XP (mp) plus "Flexstar", "Reflex", "Blazer", or "Cobra"**
Tank mix applications of SYNCHRONY® XP (mp) plus "Flexstar", "Reflex", "Blazer", or "Cobra" may not control weeds listed on the SYNCHRONY® XP (mp) label as completely as applications of SYNCHRONY® XP (mp) alone.

SYNCHRONY® XP (mp) and Postemergence Grass Herbicides:

SYNCHRONY® XP (mp) and SYNCHRONY® XP (mp) tankmixes may be tank mixed with postemergence grass herbicides such as DuPont™ ASSURE® II herbicide. For best results, apply SYNCHRONY® XP (mp) seven days before or one day after the grass herbicide. Refer to the grass herbicide label for precautions and specific use information.

SYNCHRONY® XP (mp) and "FirstRate" herbicide:

For improved Ragweed and Cocklebur control, add between 0.075 - 0.3 oz per acre to 0.75 oz per acre SYNCHRONY® XP (mp). These tankmixes will control up to 10" Cocklebur or Common Ragweed and up to 12" Giant Ragweed. Use a lower amount of "FirstRate" when weeds are less than the maximum size and under good growing conditions. Use a higher amount of "FirstRate" when weeds are approaching the maximum size and/or under unfavorable growing conditions.

A good quality petroleum-based or methylated seed oil-based crop oil concentrate must be added to the tank mix at the rate of 8 pints per 100 gallons of spray solution or at 1% v/v. An ammonium nitrogen fertilizer may be added as directed under the "Spray Additives" section.

SYNCHRONY® XP (mp) and 2,4-DB:

- In soybeans at least 8" tall, 0.75 oz SYNCHRONY® XP (mp) may be tank mixed with 1-2 fl oz per acre of 2,4-DB for improved control of giant ragweed up to 4" tall, annual morningglory and other broadleaf weeds.
- Applications of SYNCHRONY® XP (mp) and 2,4-DB must include a nonionic surfactant at the rate of 2 pints per 100 gal or crop oil concentrate at 8 pints per 100 gal of spray solution (1.0% v/v). See Spray Additives.
- Apply a tank mix of SYNCHRONY® XP (mp) and 2,4-DB by ground only.
- In Kansas and Missouri (except the bootheel area), when conditions are excessively hot and dry (> 90°F and < 30% relative humidity), make applications at the rate of 2 fl oz of 2,4-DB and SYNCHRONY® XP (mp).

8/18

- The use of crop oil concentrate may increase temporary injury to soybeans.

Some crop response may occur 5-7 days after application of DuPont™ SYNCHRONY® XP (mp) and 2,4-DB to soybeans under stress. Temporary yellowing, leaf crinkling, and/or soybean growth retardation may occur following application of SYNCHRONY® XP (mp) and 2,4-DB. Under favorable growing conditions, the crop will quickly recover.

SYNCHRONY® XP (mp) and DuPont™ CLASSIC® herbicide:

A tankmix of 0.75 oz/A SYNCHRONY® XP (mp) herbicide, plus 1/8 oz/A CLASSIC® herbicide is recommended for control of 4-6" giant ragweed in STS soybeans.

SYNCHRONY® XP (MP) BURNDOWN APPLICATION

Tank Mixtures with glyphosate

SYNCHRONY® XP (mp) at 0.75 oz per acre may be tank mixed with glyphosate-containing herbicides registered for soybeans for burndown of existing summer and winter annual weeds and limited residual control of certain summer annual broadleaf weeds.

Application Timing - Burndown

SYNCHRONY® XP (mp) tank mixtures may be applied up to 30 days before planting or prior to soybean emergence after planting.

Spray Additives - Burndown

- Always include 0.25% non-ionic surfactant.
- The addition of 1 - 2 percent dry Ammonium Sulfate by weight or 8.5 - 17 pounds per 100 gallons of water may increase performance of this tank mix.
- For enhanced burndown control, 1 pint of 2,4-D LVE (0.5 lbs a.e./A) may be added up to seven days prior to planting.

Weeds Controlled - Burndown

SYNCHRONY® XP (mp) tank mixed with glyphosate applied prior to planting will provide burndown control of the weeds listed on page three of this label as well as control of most winter and summer annual broadleaf and grass weeds. Consult the glyphosate specimen labels for specific use instructions including all weeds controlled or suppressed, the suggested use rates, maximum weed size at application, restrictions, limitations and precautions. The addition of SYNCHRONY® XP (mp) to glyphosate-containing herbicides will increase the burndown control of the following weeds versus application of glyphosate alone:

Dandelion, Common	Nutsedge, Yellow
Dock, Curly	Primrose, cutleaf evening
Hemp Sesbania	Ragweed, Common and Giant
Henbit	Sicklepod
Marestail	Smartweed, Annual
Morningglory Spp., Annual	Velvetleaf

Weeds Controlled - Limited Preemergence

When used according to this label, SYNCHRONY® XP (mp) at 0.75 oz/A can provide limited preemergence control of the weeds listed below to contribute to a clean seedbed at planting.

For season-long control, a planned PRE or POST sequential program is required.

Jimsonweed	Pigweeds (Redroot, Smooth, Palmer)
Lambsquarter	Smartweeds, Annual
Marestail	Ragweeds
Nutsedge, yellow	

Due to heavy weed pressure, delayed planting or adverse environmental conditions, additional burndown control measures may be required at planting.

Tank mixture with DuPont™ EXPRESS® XP herbicide in the States of WI and MN

Tank mixes of SYNCHRONY® XP (mp) at 0.375 oz per acre plus EXPRESS® or EXPRESS® XP herbicides are recommended for fall burndown of 3-inch dandelion in conservation tillage or no-till soybean production systems.

Apply 0.375 oz/acre SYNCHRONY® XP (mp) -plus- 0.125-0.25 oz/acre EXPRESS® or EXPRESS® XP -plus- 1% v/v Crop Oil Concentrate (1 gallon per 100 gallons spray solution) -plus- 1 pt/acre 2, 4-D LVE.

SYNCHRONY® XP (mp) + EXPRESS or EXPRESS® XP may be applied to no-till or conservation tillage fields anytime after the fall harvest, up to 45 days prior to soybean planting.

SYNCHRONY® XP (MP) 0.375 OZ PER ACRE - REDUCED RATE POSTEMERGENCE USES (NON-"STS" SOYBEANS)

- SYNCHRONY® XP (mp) at 0.375 oz/acre may be applied to any soybean type (which includes non-"STS", "STS" and "Roundup Ready") for broadleaf weed control.
- SYNCHRONY® XP (mp) at 0.375 oz/acre may be used:
 - postemergence, or
 - tankmixed with DuPont™ AUTHORITY® preplant or preemergence for improved burndown control, or
 - in a planned program following a preemergence, preplant incorporated or preplant burndown application of AUTHORITY®.

SYNCHRONY® XP (mp) at 0.375 oz per acre - postemergence application information

Timing to Weeds

Apply SYNCHRONY® XP (mp) to 1 - 4" weeds that are actively growing. Applications made to weeds that are in the cotyledon stage, larger than the size indicated, or to weeds under stress may result in unsatisfactory control.

Spray Additives with 0.375 oz/acre SYNCHRONY® XP (mp)

"STS" soybeans: When using SYNCHRONY® XP (mp) in soybeans designated as "STS", add crop oil concentrate at 8 pt per 100 gal of spray solution (1.0% v/v).

Non-"STS" Soybeans: When using SYNCHRONY® XP (mp) in soybeans that are NOT designated as "STS", add nonionic surfactant at a rate of 2 pt per 100 gal of spray solution (0.25% v/v) unless directed by other supplemental labeling.

Ammonium nitrogen fertilizer: Use a high quality, liquid nitrogen fertilizer such as:

- 28-0-0 or 30-0-0 4 - 8 pts per acre or
- 10-34-0 2 - 4 pts per acre or
- AMS: sprayable grade ammonium sulfate
(21-0-0) at 2 - 4 lb per acre

Weeds controlled by 0.375 oz per acre DuPont™ SYNCHRONY® XP (mp) alone:

- Cocklebur
- Pigweed species (including triazine resistant)
- Wild sunflower

And suppress growth of the following weed species:

- Lambsquarters
- Jimsonweed
- Milkweed, common (above ground portions only)
- Ragweed, common
- Smartweeds, annual
- Velvetleaf

Tankmixes with 0.375 oz/acre SYNCHRONY® XP (mp)

Add 0.75 - 1.25 pt/acre "Flexstar" for control of emerged 4" waterhemp and 2" eastern black nightshade, or improved control of velvetleaf and common ragweed.

Add up to 0.15 oz/acre "FirstRate" for improved control of emerged common ragweed and velvetleaf.

Add up to 0.042 oz/acre DuPont™ HARMONY® GT for improved control of emerged lambsquarters, common milkweed, annual smartweeds and velvetleaf.

Refer to the "Flexstar", "FirstRate" and HARMONY® GT labels for other weeds controlled and for the appropriate rate based on the weed sizes. Read and follow all use directions, restrictions and precautions of both SYNCHRONY® XP (mp) and the tankmix partners. When tankmixing, the most restrictive labeling applies.

Weeds controlled by 0.375 oz per acre SYNCHRONY® XP (mp) preceded by 4 - 5.3 oz per acre AUTHORITY®:

- Cocklebur
- Jimsonweed
- Kochia
- Lambsquarters
- Milkweed, Common (above ground portions only)
- Morningglory, Ivyleaf
- Nightshade, Eastern Black
- Pigweed species (including triazine resistant)
- Ragweed, Common
- Smartweeds, Annual
- Velvetleaf
- Waterhemp, Common and Tall
- Wild sunflower

SYNCHRONY® XP (mp) at 0.375 oz per acre - Burndown tankmix with DuPont™ AUTHORITY®

For improved burndown with AUTHORITY®, a tankmix of 0.375 oz/acre SYNCHRONY® XP (mp) + AUTHORITY® + 1% v/v COC may be applied to burndown broadleaf weeds up to 3" in height and annual grasses up to 1" in height. See the Pre-plant Burndown section of the AUTHORITY® label for weeds burned down by AUTHORITY®. Addition of 0.375 oz/acre SYNCHRONY® XP (mp) will improve burndown of

dandelion, marestail, prickly lettuce, giant ragweed, common ragweed, pennycress, small-flowered bittercress, shepherd's purse and tansy mustard. One pint of 2,4-D LVE (0.5 lbs a.e./acre) may be tankmixed with AUTHORITY® + 0.375 oz/acre SYNCHRONY® XP (mp) up to 7 days prior to planting.

Precautions for 0.375 oz SYNCHRONY® XP (mp) uses

- A temporary crop response may occur following an application of SYNCHRONY® XP (mp) to soybeans not designated "STS" in the variety name.
- Applications made when the crop and weeds are under stress from moisture, cold, heat, high humidity, disease, insect pressure and prior herbicide stress may result in excessive crop response and/or reduced weed control effectiveness.
- Do not tankmix "Poast Plus" with 0.375 oz/acre SYNCHRONY® XP (mp) unless the soybean is designated as an "STS" variety.

TANK MIX COMPATIBILITY TESTING

Perform a jar test prior to tank mixing to ensure compatibility of SYNCHRONY® XP (mp) and other pesticides. Use a clear glass quart jar with lid and mix the tank mix ingredients in their relative proportions. Invert the jar containing the mixture several times and observe the mixture for approximately 1/2 hour. If the mixture balls-up, forms flakes, sludges, gels, oily film or layers, or other precipitates, it is not compatible.

MIXING INSTRUCTIONS

1. Fill the tank 1/4 to 1/3 full of water.
2. While agitating, add the required amount of SYNCHRONY® XP (mp).
3. Continue adequate agitation.
4. SYNCHRONY® XP (mp) should be thoroughly mixed with water in the spray tank before adding any other material. As the tank is filling, add (in order): other herbicide(s), the required spray adjuvant, and the nitrogen fertilizer where required.
5. Apply SYNCHRONY® XP (mp) spray preparation within 24 hours of mixing to avoid product degradation.
6. If the mixture has settled, thoroughly reagitrate before using.

APPLICATION EQUIPMENT

Ground Application

(See Also Spray Drift Management)

Broadcast Application

- Postemergence in soybeans, use a minimum of 10 gal water per acre. Under heavy weed pressure or dense crop foliage, increase minimum spray volume to 15-25 gal per acre.
- Use flat fan nozzles at 25-60 psi for SYNCHRONY® XP (mp) applications.
- For burndown applications of existing vegetation, flat fan nozzles and a minimum of 20 gallons per acre are recommended so as to thoroughly wet the foliage to be burned down.
- Do not use flood, rain drop, whirl chamber, or controlled droplet applicator (CDA) type nozzles. Unacceptable crop injury, excessive spray drift, or poor weed control may result.

10/18

- For proper spray coverage adjust the boom and nozzle height according to the specifications listed by the nozzle manufacturer.

Band Application

- Because band applicators spray a narrower area than broadcast applicators, use proportionately less spray solution for band applications.
- Carefully calibrate the band applicator to not exceed the labeled rate.
- Flat fan nozzles are recommended.
- Carefully follow the nozzle manufacturer's instructions for nozzle orientation, distance of the nozzles from the crop and weeds, spray volumes, calibration, and spray pressure for band applications.
- For additional information on row banders, see DuPont bulletin, "Application Accuracy - Row Banders."

**Aerial Application
(See Also Spray Drift Management)**

- Use nozzle types and arrangements that will provide optimum spray distribution and maximum coverage at 3-5 gal per acre.
- Use a minimum of 3 gal water per acre. Under heavy weed pressure or dense crop foliage, increase the minimum spray volume to 5 gal per acre.
- Do not apply during a temperature inversion, when winds are gusty, or when other conditions could produce poor coverage and/or off-target spray movement.

**ENVIRONMENTAL CONDITIONS
AND BIOLOGICAL ACTIVITY**

DuPont™ SYNCHRONY® XP (mp) rapidly inhibits the growth of susceptible weeds. Leaves of susceptible plants yellow 3-5 days after application, followed, in controlled plants, by the death of the growing point. SYNCHRONY® XP (mp) will provide complete control of susceptible weeds in 7-21 days. Suppressed plants may remain green but will be stunted and noncompetitive.

SYNCHRONY® XP (mp) will provide best results when applied to young, actively growing weeds. Degree of control depends on: weed spectrum; weed size (if weeds are large, use higher spray volume); growing conditions at and following treatment; soil moisture; precipitation; and spray adjuvants. Treating weeds under stress or large weeds may result in only partial control. Stress may be caused by:

- abnormal weather (hot or cold)
- mechanical injury from cultivation
- drought
- water-saturated soil
- disease
- insect injury
- prior herbicide injury

Stress affects some weeds, such as pigweed, more than others. Delay application until stress passes and weeds start to grow again.

Severe stress (drought, disease, insect damage, or nutrient deficiency such as iron chlorosis) following application may also result in poor weed control.

Do not apply SYNCHRONY® XP (mp) if rain is expected within

1 hour or weed control may decrease.

ROTATIONAL CROP GUIDELINES

Crop rotation intervals noted in the table below are based on crops grown under favorable growing conditions. Crops grown under unfavorable environmental conditions, such as drought, nutrient deficiency, high salts, disease and insect pressure may demonstrate reduced tolerance to crop protection chemicals. When deciding on a particular crop to replant in your fields, carefully consider your particular soil and other field conditions (see IMPORTANCE OF SOIL pH section of this label).

- Important: Crops other than soybeans following a SYNCHRONY® XP (mp) application can vary in their sensitivity to low concentrations of SYNCHRONY® XP (mp) remaining in the soil. Rotational crop guidelines must be followed.

Region A: The states of Delaware, Illinois, Indiana, Iowa (fields located outside the boundaries of the Clarion-Nicollet-Webster and Hamburg-Ida-Monona soil associations and fields located outside the historic flood plain of the Missouri River), Kansas, Kentucky, Maryland, Michigan (fields south of Interstate 96 or per supplemental labeling), Missouri (except the Bootheel), Nebraska (fields south of Route 30 and east of Route 281), New Jersey, Ohio, Pennsylvania and West Virginia.

Region B: The states of Alabama (except the "Black Belt" where soil pH must be less than 7.0), Arkansas, Florida, Georgia, Louisiana, Missouri (Bootheel region only), Mississippi (except the "Black Belt" where soil pH must be less than 7.0), North Carolina, Oklahoma, South Carolina, Tennessee, Texas (fields east of Route 183), and Virginia.

Region C: The states of Minnesota, South Dakota, Wisconsin, Iowa (fields inside the boundaries of the Clarion-Nicollet-Webster and Hamburg-Ida-Monona soil associations and fields located inside the historic floodplain of the Missouri River) and Nebraska (fields north of route 30 and west of Route 281). **Do not exceed 0.375 oz/acre per season in region C states.**

Follow Recrop Interval 1 if:

- The field is located in a "Region A" state (all pH soils) AND, EITHER
- A maximum of 1.125 oz/acre of SYNCHRONY® XP (mp) was applied for the use season, OR
- A maximum of 0.75 oz/acre of SYNCHRONY® XP (mp) in sequence with 1/3 oz/acre of DuPont™ CLASSIC® was applied for the use season.

Follow Recrop Interval 1 if:

- The field is located in a "Region A" state with soil pH 7.0 or less AND
- A maximum of 0.75 oz/acre of SYNCHRONY® XP (mp) was applied in sequence with a maximum rate of 3/4 oz/acre of CLASSIC®, or a maximum of two applications of SYNCHRONY® XP (mp) at a rate of 0.75 oz/acre per application was applied or a single maximum application of 1.125 oz.

Follow Recrop Interval 2 if:

- The field is located in a "Region B" state (all pH soils except those with pH greater than 7.0 in Virginia and the Black Belt

region of Alabama and Mississippi)
AND

- A maximum of 0.75 oz/acre of DuPont™ SYNCHRONY® XP (mp) was applied in sequence with 3/4 oz/acre of CLASSIC®, or a maximum of two applications of SYNCHRONY® XP (mp) at a rate of 0.75 oz/acre per application was applied or a single maximum application of 1.125 oz.

Follow Recrop Interval 3 if:

- The field is located in a “Region A, B or C” state
AND
- A maximum of 0.375 oz/acre SYNCHRONY® XP (mp) was applied during the use season.

**Rotational Interval (months)
Following the Use of
SYNCHRONY® XP (mp)***

Crop	Interval 1	Interval 2	Interval 3
Soybeans	Anytime	Anytime	Anytime
Cereal Grains, Pasture Grasses (such as Fescue and Ryegrass)	3	3	3
Dry Beans, Kidney Beans			
Peas, Snap Beans	9	9	9
Field Corn (IR)	8	7	8
Field Corn**(States in Regions A & C)	9	---	9
Field Corn**(States of AR, MO [Bootheel only], NC, OK, TN, and TX)	---	8	---
Field Corn**(States of AL, FL, GA, LA, MS, and SC)	---	7	---
Popcorn	9	9	9
Sorghum	9	9	9
Tobacco (transplant)	9	9	9
Tomato (transplant)	9	9	9
Peanuts	15	6	6
Rice	15	9	9
Cotton	9	8	9
Alfalfa	12	9	9
Clover	12	9	9
Cabbage	18	18	18
Canola (Rapeseed)	18	18	18
Cucumber	18	18	9
Flax	18	18	18
Lentils	18	18	18
Mustard	18	18	18
Pumpkins	18	18	9
Sunflower	18	18	9
Sweet Corn	18	18	9†
Watermelon	18	18	9
Carrots	30	30	30
Onions	30	30	30
Potatoes (including Sweet Potatoes)	30	30	30
Potatoes, irish	---	8††	---
Sugar Beets	30	30	30
Any crop not listed	30	30	30

* If SYNCHRONY® XP (mp) or the latter part of a sequential treatment containing chlorimuron ethyl (such as DuPont™ CLASSIC®) is applied after August 1, extend rotational crop intervals 2 months for alfalfa, clover, corn (non-IR), cotton, popcorn, rice, sorghum, tobacco, and tomato.

**The term “Field Corn” is defined to include only that corn grown for grain or silage or for seed corn relative to the Rotational Crop Guidelines section of this label.

† Rotational crop intervals are for processing Sweet Corn varieties only. The rotational crop interval for other Sweet Corn varieties is 18 months.

††States of NC and VA in soils with organic matter greater than 1%.

THE IMPORTANCE OF SOIL PH

Soil pH varies greatly, even within the same field. pH variations as much as 2 pH units are common. Composite soil samples taken across an entire field, such as those samples taken for soil fertility recommendations, may not detect areas of high pH. Sub-sampling is recommended for areas likely to have pH values higher than the field average. The following is a non-inclusive list of potential high pH areas where subsampling is recommended.

- Where different soil types are evident within a field, sample soil types separately.
- Where conditions vary within a field, sample areas separately, such as:
 - areas bordered by limestone gravel roads,
 - river bottoms subject to flooding,
 - low areas in hardpan soils where evaporative ponds may occur,
 - eroded hillsides,
 - along drain tile lines, and
 - areas where drainage ditch spoil has been spread.
- Where lime has not been deeply incorporated, soil may exhibit significantly higher pH values in the upper 3 inches of soil. Composite soil samples taken at a 6-8 inch depth may not reflect the elevated pH near the surface. In these cases shallow sampling, the upper 3 inches, is advised.

Determine soil pH by laboratory analysis using a 1:1 soil:water suspension.

SPRAYER PREPARATION AND CLEANUP

Prior to application of SYNCHRONY® XP (mp), start with clean, well maintained application equipment. Immediately following application, thoroughly clean all application equipment. Postponing action, even for a few hours, only makes effective cleanup more difficult. Failure to clean spraying equipment thoroughly may result in injury to subsequently sprayed crops.

When spraying multiple loads of SYNCHRONY® XP (mp) over an extended period of time, rinse the equipment with clean water at the end of the day. Leave water in the equipment overnight to prevent deposits from drying on surfaces.

When applications of SYNCHRONY® XP (mp) are completed and prior to using the sprayer and associated equipment for other products or for crops other than soybeans, thoroughly clean the equipment using the procedure below.

STEP 1. Drain spray equipment. Thoroughly rinse sprayer, and flush hoses, boom and nozzles with clean water. Loosen and physically remove visible deposits.

STEP 2. Fill the sprayer with clean water and add household ammonia (one gallon of 3% active for every 100 gallons of water) or correct amount of a DuPont approved cleaner*. Flush hoses, boom and nozzles. Turn off the boom and top off the tank with clean water. Circulate through the spraying system for 15 minutes. Flush the hoses, boom and nozzles with the cleaning solution. Drain the tank.

STEP 3. Remove and clean nozzle, screens and strainers in a bucket of fresh cleaner and water.

STEP 4. Repeat STEP 2.

12/18

STEP 5. Thoroughly rinse the sprayer, hoses, boom and nozzles with clean water, several times.

Clean all other associated application equipment. Take all necessary safety precautions when cleaning equipment. Do not clean near wells, water sources or near desirable vegetation. Dispose of waste rinse water in accordance with local regulations.

* For additional information on sprayer cleanup and a listing of DuPont-approved cleaners, see DuPont Bulletin "A Guide To Application Equipment Cleanout For DuPont Sulfonylurea Herbicides".

SPRAY DRIFT MANAGEMENT

The interaction of many equipment and weather-related factors determines the potential for spray drift. The applicator is responsible for considering all these factors when making application decisions.

AVOIDING SPRAY DRIFT IS THE RESPONSIBILITY OF THE APPLICATOR.

IMPORTANCE OF DROPLET SIZE

The most effective way to reduce drift potential is to apply large droplets (>150 - 200 microns). The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. The presence of sensitive species nearby, the environmental conditions, and pest pressure may affect how an applicator balances drift control and coverage. **APPLYING LARGER DROPLETS REDUCES DRIFT POTENTIAL, BUT WILL NOT PREVENT DRIFT IF APPLICATIONS ARE MADE IMPROPERLY OR UNDER UNFAVORABLE ENVIRONMENTAL CONDITIONS!** See **Wind, Temperature and Humidity, and Temperature Inversions** sections of this label.

Controlling Droplet Size - General Techniques

- **Volume** - Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets.
- **Pressure** - Use the lower spray pressures recommended for the nozzle. Higher pressure reduces droplet size and does not improve canopy penetration. **WHEN HIGHER FLOW RATES ARE NEEDED, USE A HIGHER-CAPACITY NOZZLE INSTEAD OF INCREASING PRESSURE.**
- **Nozzle Type** - Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles.

Controlling Droplet Size - Aircraft

- **Number of Nozzles** - Use the minimum number of nozzles with the highest flow rate that provide uniform coverage.
- **Nozzle Orientation** - Orienting nozzles so that the spray is emitted backwards, parallel to the airstream will produce larger droplets than other orientations.
- **Nozzle Type** - Solid stream nozzles (such as disc and core with swirl plate removed) oriented straight back produce larger droplets than other nozzle types.
- **Boom Length** - The boom length should not exceed 3/4 of the wing or rotor length - longer booms increase drift potential.
- **Application Height** - Application more than 10 ft above the canopy increases the potential for spray drift.

BOOM HEIGHT

Setting the boom at the lowest labeled height (if specified) which provides uniform coverage reduces the exposure of droplets to evaporation and wind. For ground equipment, the boom should remain level with the crop and have minimal bounce.

WIND

Drift potential increases at wind speeds of less than 3 mph (due to inversion potential) or more than 10 mph. However, many factors, including droplet size and equipment type determine drift potential at any given wind speed. **AVOID GUSTY OR WINDLESS CONDITIONS.**

Note: Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they affect spray drift.

TEMPERATURE AND HUMIDITY

When making applications in hot and dry conditions, set up equipment to produce larger droplets to reduce effects of evaporation.

TEMPERATURE INVERSIONS

Drift potential is high during a temperature inversion. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain close to the ground and move laterally in a concentrated cloud. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

SHIELDED SPRAYERS

Shielding the boom or individual nozzles can reduce the effects of wind. However, it is the responsibility of the applicator to verify that the shields are preventing drift and not interfering with uniform deposition of the product.

AIR ASSISTED (AIR BLAST) FIELD CROP SPRAYERS

Air assisted field crop sprayers carry droplets to the target via a downward directed air stream. Some may reduce the potential for drift, but if a sprayer is unsuitable for the application and/or set up improperly, high drift potential can result. It is the responsibility of the applicator to determine that a sprayer is suitable for the intended application, is configured properly, and that drift is not occurring. **Note:** Air assisted field sprayers can affect product performance by affecting spray coverage and canopy penetration. Consult the application equipment section of this label to determine if use of an air assisted sprayer is recommended.

13/18

AIR ASSISTED (AIR BLAST) TREE AND VINE SPRAYERS

Air assisted tree and vine sprayers carry droplets into the canopy of trees and vines via a radially or laterally directed air stream. These sprayers are not suitable for applying herbicides.

IMPORTANT PRECAUTIONS

Do not graze treated fields or harvest for forage or hay.

Do not apply this product through any type of irrigation equipment.

Injury to or loss of desirable trees or vegetation may result from failure to observe the following:

- Do not apply DuPont™ SYNCHRONY® XP (mp) or drain or flush equipment on or near desirable trees or other plants, on areas where their roots may extend, or in locations where the chemical may be washed or moved into contact with their roots.
- Do not use on lawns, walks, driveways, tennis courts or similar areas.
- Many crops are sensitive to SYNCHRONY® XP (mp). All direct or indirect contact (such as spray drift) with crops other than soybeans should be avoided.
- Thoroughly clean all application equipment immediately after use and prior to spraying crops other than soybeans.

RESISTANCE

When herbicides that affect the same biological site of action are used repeatedly over several years to control the same weed species in the same field, naturally-occurring resistant biotypes may survive a correctly applied herbicide treatment, propagate, and become dominant in that field. Adequate control of these resistant weed biotypes cannot be expected. If weed control is unsatisfactory, it may be necessary to retreat the problem area using a product affecting a different site of action.

To better manage herbicide resistance through delaying the proliferation and possible dominance of herbicide resistant weed biotypes, it may be necessary to change cultural practices within and between crop seasons such as using a combination of tillage, retreatment, tank-mix partners and/or sequential herbicide applications that have a different site of action. Weed escapes that are allowed to go to seed will promote the spread of resistant biotypes.

It is advisable to keep accurate records of pesticides applied to individual fields to help obtain information on the spread and dispersal of resistant biotypes. Consult your agricultural dealer, consultant, applicator, and/or appropriate state agricultural extension service representative for specific alternative cultural practices or herbicide recommendations available in your area.

INTEGRATED PEST MANAGEMENT

DuPont recommends the use of Integrated Pest Management (IPM) programs to control pests. This product may be used as part of an Integrated Pest Management (IPM) program which can include biological, cultural, and genetic practices aimed at preventing economic pest damage. Application of this product should be based on IPM principles and practices including field scouting or other detection methods, correct target pest identification, population monitoring, and treating when target pest populations reach locally determined action

thresholds. Consult your state cooperative extension service, professional consultants or other qualified authorities to determine appropriate action treatment threshold levels for treating specific pest/crop or site systems in your area.

PESTICIDE STORAGE AND DISPOSAL

Pesticide Storage: Store product in original container only. Do not contaminate water, other pesticides, fertilizer, food or feed in storage.

Pesticide Disposal: Do not contaminate water, food, or feed by disposal. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: Completely empty bag into application equipment. Then dispose of bag in a sanitary landfill, or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

Notice to Buyer: Purchase of these materials does not confer any rights under patents of countries outside of the United States. Use of this quantity of purchased DuPont™ SYNCHRONY® XP (MP) herbicide is permitted under claim 24 of U.S. Patent 5,084,082.

The DuPont Oval Logo, DuPont™, SYNCHRONY®, ASSURE®, CANOPY®, CANOPY XL®, CLASSIC®, EXPRESS®, STEADFAST® and HARMONY® are trademarks or registered trademarks of E.I. duPont de Nemours and Company

AUTHORITY® is a registered trademark of FMC Corp.

"Blazer" and "Poast Plus" are registered trademarks of BASF Corporation

"Cobra" is a registered trademark of Valent USA Corporation

"Flexstar", "Reflex", are trademarks or registered trademarks of Syngenta Crop Protection Inc.

"FirstRate" is a registered trademark of Dow AgroSciences

D - 986 112904

LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read This Limitation of Warranty and Liability Before Buying or Using This Product. If the Terms Are Not Acceptable, Return the Product at Once, Unopened, and the Purchase Price Will Be Refunded.

It is impossible to eliminate all risks associated with the use of this product. Such risks arise from weather conditions, soil factors, off target movement, unconventional farming techniques, presence of other materials, the manner of use or application, or other unknown factors, all of which are beyond the control of DuPont. These risks can cause: ineffectiveness of the product; crop injury, or; injury to non-target crops or plants.

DuPont does not agree to be an insurer of these risks. **WHEN YOU BUY OR USE THIS PRODUCT, YOU AGREE TO ACCEPT THESE RISKS.**

DuPont warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for the purpose stated in the Directions for Use, subject to the inherent risks described above, when used in accordance with the Directions for Use under normal conditions.

DUPONT MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR OF MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

TO THE FULLEST EXTENT PERMITTED BY LAW; IN NO EVENT SHALL DUPONT OR SELLER BE LIABLE FOR ANY INCIDENTAL, CONSEQUENTIAL OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. BUYER'S OR USER'S BARGAINED-FOR EXPECTATION IS CROP PROTECTION. THE EXCLUSIVE REMEDY OF THE USER OR BUYER AND THE EXCLUSIVE LIABILITY OF DUPONT OR SELLER, FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY OR CONTRACT, NEGLIGENCE, TORT OR STRICT LIABILITY), WHETHER FROM FAILURE TO PERFORM OR INJURY TO CROPS OR OTHER PLANTS, AND RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT, OR AT THE ELECTION OF DUPONT OR SELLER, THE REPLACEMENT OF THE PRODUCT.

DuPont or its Ag Retailer must have prompt notice of any claim so that an immediate inspection of buyer's or user's growing crops can be made. Buyer and all users shall promptly notify DuPont or a DuPont Ag Retailer of any claims, whether based on contract, negligence, strict liability, other tort or otherwise or be barred from any remedy.

This Limitation of Warranty and Liability may not be amended by any oral or written agreement.

For product information call: 1-888-6-DUPONT

Internet address: www.dupont.com/ag/us

© 2004 E. I. du Pont de Nemours and Company, Crop Protection, Wilmington, Delaware 19898.

All rights reserved.

15/14

DuPont Agricultural Products

SUPPLEMENTAL LABELING

"..... A Growing Partnership With Nature"

DUPONT SYNCHRONY® XP (MP) HERBICIDE- EXPANDED USE RATES FOR PREEMERGENCE AND BURNDOWN APPLICATIONS IN SOYBEANS

DUPONT SYNCHRONY® XP (MP) HERBICIDE

EPA Reg. No. 352-XXX

EXPANDED USE RATES FOR PREEMERGENCE AND BURNDOWN APPLICATIONS IN SOYBEANS

Directions For Use

SYNCHRONY® XP (mp) at 1 to 3 oz/acre may be used for burndown and residual weed control in all states located in SYNCHRONY® XP (mp) Rotational Regions A and B on the main Federal Section 3 SYNCHRONY XP (mp) product label, excluding the state of Florida (refer to Rotational Crop Guidelines).

Application Timing - SYNCHRONY® XP (mp) may be applied to no-till or conservation tillage fields 45 days before planting soybeans to just before soybean emergence. Do not apply to frozen ground.

Application Rates:

Medium and Fine Soils, 1.5 - 4% Organic Matter	Rate
Region A States:	
No pH restriction*	1 oz/acre
Region B States:	
<u>Composite soil pH of 7 or less</u>	<u>1.25 - 3 oz/acre</u>
No pH restriction	1 - 1.5 oz/acre
Composite soil pH of 7 or less	greater than 1.5 up to 3 oz/acre

* In Michigan, do not apply the 1 oz/acre rate to soils exceeding pH 7.6. In all other states, the soil pH is unrestricted for 1 oz/acre rate.

16/13

For season-long control of all grass and broadleaf weeds following 1 – 3 oz/acre applications of SYNCHRONY® XP (mp), a planned sequential program is required. Use higher rates of SYNCHRONY® XP (mp) where longer residual control is desired.

Weeds Controlled

Burndown Control of existing winter and summer annual weeds

SYNCHRONY® XP (mp) applications will provide burndown control of certain broadleaf weeds which are no greater than 3 inches in height.

To obtain burndown of the weed species listed below:

- Addition of Crop Oil Concentrate at 1% v/v (1 gallon per 100 gallons of final spray volume) is required.
- Use a minimum of 20 gallons per acre with spray nozzles that provide thorough spray coverage of the weeds.
- 2,4-D LVE may be added for enhanced burndown control.

Bittercress, small-flowered	Henbits	Pigweed	Tansy mustard
Bushy wallflower	Lambsquarters	Ragweed, common	Thistle - Canadian (above ground portion)
Buttercup, smallflower	Lettuce, prickly	Ragweed, giant	Velvetleaf
Butterweed	Marestail*	Shepherdspurse	Whitlowgrass
Dandelion	Mustard wild	Smartweed, annual	Yellow rocket
Deadnettle (purple, red)	Pennycress	Speedwell (field, Purselane)	
Garlic, wild*	Pepperweed	Sunflower	

* Addition of 1 pt/acre 2,4-D LVE is required for the 1 oz/acre rate of SYNCHRONY® XP (mp), and recommended for all other rates.

To burndown annual grasses and broadleaf weeds listed above when they exceed the recommended height, SYNCHRONY® XP (mp) may be tank mixed with other herbicides registered for preemergence or burndown application prior to soybean emergence. When tank mixing with glyphosate-containing products, replace the Crop Oil Concentrate with nonionic surfactant at 0.25% v/v (1 quart per 100 gallons final spray volume) and follow the manufacturer's instructions for ammonium sulfate addition.

Preemergence Residual Control

Applications of 1.25 – 3 oz/acre SYNCHRONY® XP (mp) will provide preemergence control or partial control (suppression) of the following weeds:

<u>Control</u>	<u>Suppression</u>
Cocklebur	Annual grasses*
Lambsquarters	(foxtails, barnyardgrass, crabgrass, panicum)
Marestail	Chickweed, common
Pigweeds (redroot, smooth)	Jimsonweed
Purselane	Morningglory, annual
Speedwell	Nutsedge, yellow
Ragweed, common	Prickly Sida (teaweed)
Smartweeds, annual	Ragweed, giant
Velvetleaf	

If applying 1 oz/acre of SYNCHRONY® XP (mp), heavy weed pressure, delayed planting, or adverse environmental conditions may require additional burndown control measures at planting. For enhanced residual control, products such as 2-4 oz/acre "Sencor" may be tank mixed with 1 oz/acre SYNCHRONY® XP (mp).

Early spring applications of 1 oz/acre SYNCHRONY® XP (mp) will provide limited residual control of the above-listed weeds to contribute to a clean seed at planting.

17/18

Planned Sequential Programs

SYNCHRONY® XP (mp) will not provide adequate season-long preemergence control of annual grasses and all broadleaf weeds. For season-long control, follow SYNCHRONY® XP (mp) with postemergence herbicide applications based on targeted weeds. To insure maximal rotation flexibility when considering a sequential program of SYNCHRONY® XP (mp) followed by CLASSIC® or SYNCHRONY® XP(mp), carefully consider: the soil pH, the recommendations below, the rotational information in this section, and the Rotational Crop Guidelines on product labeling.

Applications of 1 oz/acre SYNCHRONY® XP (mp) [Region A and B states] to soils with pH greater than 7: Do not apply additional chlorimuron-ethyl-containing herbicides (CLASSIC®, SYNCHRONY® XP) except in the states of AL, AR, GA, LA, MO bootheel, MS, NC, OK, SC, TN, and TX, where up to 0.5 oz/acre SYNCHRONY® XP (mp) may be applied.

Applications of 1.5 oz/acre SYNCHRONY® XP (mp) [Region B states] to soils with pH greater than 7: Do not apply additional chlorimuron-ethyl-containing herbicides (CLASSIC®, SYNCHRONY® XP)

Applications of 1-3 oz SYNCHRONY® XP (mp) to soils with pH less than 7 may be followed with a single postemergence application of CLASSIC® or SYNCHRONY® XP. See table below for rates that may be applied:

Expanded Application Rate SYNCHRONY® XP (mp) oz/acre	CLASSIC® oz/acre	SYNCHRONY® XP (mp) oz/acre
up to 2	up to 3/4	up to 1/2
2.1 - 2.5	up to 2/3	up to 1/2
2.6 - 3	up to 1/4	none

Refer to the sequential herbicide labels for specific information regarding use rates, application timing, crop rotations and other restrictions and precautions.

Rotational Crop Information

For rotational crop information following 1 oz/acre SYNCHRONY® XP (mp) in Region A states, and up to 1.5 oz/acre applications in Region B states, use Recrop Interval 1 or 2 under the Section 'Rotational Crop Guidelines' on the main Federal Section 3 SYNCHRONY XP (mp) product label, depending on whether the use was in a Region A or Region B state.

For all other applications of SYNCHRONY® XP (mp) under this supplemental labeling, follow the recropping intervals given in the table below. Crop rotation intervals noted in the table below are based on crops grown under favorable growing conditions. Crops grown under unfavorable environmental conditions, such as drought, nutrient deficiency, high salts, disease and insect pressure may demonstrate reduced tolerance to crop protection chemicals. When deciding on a particular crop to replant in your fields, carefully consider your particular soil and other field conditions.

K/R

<u>Crop</u>	<u>Recropping Interval in Months</u>
Soybeans	anytime
Cereal grains, pasture, grasses	4
Peanuts	8
Alfalfa	10
Cotton, Rice	10
Tobacco and Tomato transplants	10
Field Corn*	10**
Clover, Sorghum	12
Dry Beans, Kidney Beans, Snap Beans, Peas	12
Cucumber, Flax, Pumpkin	18
Sunflower, Sweet Corn, Watermelon	18
Cabbage, Canola, Lentils, Mustard	18
Carrot, Onion, Potato (all types), Sugarbeets and any other crop not listed	30†

* Field Corn is defined to include only that corn grown for grain, silage, popcorn, and seed corn. However, because seed corn inbred lines may vary in their sensitivity to trace amounts of herbicide carryover, DuPont cannot warrant that seed corn can be recropped without damage or yield loss. Users should seek the advice of their seed corn company agronomists regarding inbred sensitivity to herbicides prior to planting any inbred lines.

**In the states of DE, KY, MD, MO bootheel, NJ, NC, SC, TN, VA, and WV, field corn may be recropped after 9 months if the SYNCHRONY® XP (mp) rate does not exceed 2.5 oz/acre.

† Carrots, onions, potato (all types), sugarbeets, and any other crop not listed may be recropped after 18 months in the states of AL, AR, DE, GA, LA, MD, MS, MO bootheel, NJ, NC, SC, TN, VA, and WV.

**IMPORTANT
BEFORE USING SYNCHRONY XP (MP), READ AND FOLLOW ALL APPLICABLE DIRECTIONS,
RESTRICTIONS, AND PRECAUTIONS ON THE EPA-REGISTERED LABEL.**

This bulletin contains new or supplemental instructions for use of this product which do not appear on the EPA-registered package label. Follow the instructions carefully.

This labeling must be in the possession of the user at the time of pesticide application.

Read the Limitation of Warranty and Liability on the Section 3 Federal product label before buying or using THIS product. If terms are not acceptable, return the unopened package at once to Seller for full refund of purchase price paid. Otherwise, use by Buyer or any other User constitutes acceptance of the terms of the Limitation of Warranty and Liability on the Section 3 Federal product label.