

352-669

7/1/2009

4/7



U.S. ENVIRONMENTAL PROTECTION AGENCY
Office of Pesticide Programs
Registration Division (7505C)
1200 Pennsylvania Ave., N.W.
Washington, D.C. 20460

EPA Reg. Number:

352-669

Date of Issuance:

JUL - 1 2009

NOTICE OF PESTICIDE:

Registration
 Reregistration

(under FIFRA, as amended)

Term of Issuance:

Unconditional

Name of Pesticide Product:

DuPont Cimarron X-Tra

Name and Address of Registrant (include ZIP Code):

E.I. DuPont de Nemours & Company, Inc
DuPont Crop Protection
Stine-Haskell Research Center
P. O. Box 30
Newark, DE 19714-0030

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

Based on your response to the Reregistration Eligibility Document, EPA has reregistered the product listed above provided you make the following changes to your label.

1. Reorder your first aid statements to read: "If on skin or clothing", "If in Eyes", "If Swallowed".
2. Based on the acute toxicology review, the Hazards to Humans and Domestic Animals must be revised to include the statement "prolonged or frequently repeated skin contact may cause allergic reactions in some individuals."

Signature of Approving Official:

James A. Tompkins, Product Manager (25)
Herbicide Branch, Registration Division (7505P)

Date:

JUL - 1 2009

3. The PPE section must be revised to include the text (PPE) immediately following the Personal Protective Equipment heading. In addition add “chemical resistant gloves made of any waterproof material such as polyethylene or polyvinylchloride” to the list of required PPE.

4. Revise the second sentence in the paragraph above Engineering Controls to read “If no such instructions for washables **exist**, use detergent and hot water.

5. A Non-Agricultural Use Requirements box must be added to the label with the following entry restriction text:

“NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are not within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Do not enter or allow others to enter until sprays have dried.”

6. On page 2, revised the first paragraph at the top of the second column to read “DuPont Cimarron X-tra should be used only in accordance with **directions** on this label or in separate published DuPont **directions**. DuPont will not be responsible for losses or damages resulting from the use of this product in any manner not specifically **directed** by DuPont.

7. On page 2, under General Information, revise the first sentence by deleting the word “recommended”. The sentence may be modified as necessary.

8. On page 3, under Application Information for Conservation Reserve Program (CRP), revise the first sentence by deleting the word “recommended”. The sentence may be modified as necessary.

9. On page 5, under Musk Thistle, revise the next to last sentence to read “Consult with you local DuPont representative, dealer or applicator for specific use rate and tank mix **information** for your area.

10. On page 6, under Spray Adjuvants, revise the first sentence to read “Unless otherwise **stated**, applications if Cimarron X-tra...”

11. On page 6, under Herbicide Tank Mixtures for Pastures and Rangeland:, revise the second and third sentence to read “Read and follow all manufacturer’s label **instructions** for the companion herbicide. If those **instructions** conflict with this label, do not tank mix...”

12. On page 7, under With Liquid Nitrogen Solution Fertilizer, revise the last sentence in the fifth paragraph to read “Consult your agricultural dealer, consultant, fieldman, or DuPont representative for specific **guidance** before adding an adjuvant to these tank mixtures.”

13. On page 7, under the Grazing and haying section, remove the statement “Coveralls and shoes plus socks must be worn if cutting within 4 hours of treatment.

Page 3

EPA Registration No. 352-669

Enclosed is a copy of your label stamped "Accepted With Comments". This action is taken under the authority of section 4(g)(2)© of the Federal Insecticide, Fungicide, and Rodenticide Act as amended. Reregistration under this section does not eliminate the need for continual reassessment of pesticides. EPA may require submission of data at any time to maintain registration of your product.

Submit one (1) copy of the final printed label before you release the product for shipment. Products shipped after 12 months from the date of this Notice or the next printing whichever occurs first, must bear the new revised label. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.



The miracles of science™

H-

4
17

DuPont™ Cimarron®

X-tra

herbicide

DRAFT LABEL



The miracles of science™

DuPont™ Cimarron® X-tra

herbicide

*For Use on Pastures, Rangeland or Established
Grasses on Acres Enrolled in the Conservation
Reserve Program*

<i>Active Ingredient</i>	<i>By Weight</i>
Metsulfuron Methyl Methyl 2-[[[(4-methoxy-6-methyl -1,3,5-triazin-2-yl)amino]carbonyl] amino]sulfonyl]benzoate	30.0%
Chlorsulfuron 2-Chloro-N-[(4-methoxy-6-methyl- 1,3,5-triazin-2-yl)aminocarbonyl] benzenesulfonamide	37.5%
<i>Inert Ingredients</i>	32.5%
TOTAL	100.0%

EPA Reg. No. 352-669

EPA Est. No. _____

Net Weight: _____

**ACCEPTED
with COMMENTS
In EPA Letter Dated:**

JUL 1 - 2009

**Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under EPA Reg. No.
352-669**

KEEP OUT OF REACH OF CHILDREN

CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand this label, find someone to explain it to you in detail.)

FIRST AID

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

IF SWALLOWED: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything to an unconscious person.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-441-3637 for emergency medical treatment information.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION! Causes moderate eye irritation. Harmful if swallowed or absorbed through skin. Avoid contact with skin, eyes, or clothing. Avoid breathing dust or spray mist. Remove and wash contaminated clothing before reuse.

PERSONAL PROTECTIVE EQUIPMENT

Some of the materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for Category A on an EPA chemical-resistance category selection chart.

Mixers, loaders, applicators, and other handlers must wear:

Long-sleeved shirt and long pants

Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Control Statement: When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet. Users should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate.

IMPORTANT INFORMATION

PESTICIDE HANDLING

- Calibrate sprayers only with clean water away from the well site.
- Make scheduled checks of spray equipment.
- Assure accurate measurement of pesticides by all operation employees.
- Mix only enough product for the job at hand.
- Avoid overfilling of spray tank.
- Do not discharge excess material on the soil at a single spot in the field or mixing/loading station.
- Dilute and agitate excess solution and apply at labeled rates/uses.
- Avoid storage of pesticides near well sites.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 4 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

Coveralls.

Chemical-resistant gloves made of any waterproof material

Shoes plus socks.

DuPont™ CIMARRON® X-tra should be used only in accordance with recommendations on this label or in separate published DuPont recommendations. DuPont will not be responsible for losses or damages resulting from the use of this product in any manner not specifically recommended by DuPont.

Do not apply this product through any type of irrigation system.

GENERAL INFORMATION

CIMARRON® X-tra is recommended for use on land primarily dedicated to the production of pasture, rangeland, or established grasses in the Conservation Reserve Program (CRP). This product may also be used on selected uncultivated areas (fence rows, farmyards, and rights-of-way) directly adjacent to, or which transect or pass through, treated pastures, rangeland, or CRP, where grazing or harvesting for animal feed of those uncultivated areas may occur.

A spray adjuvant must be used in the spray mix unless otherwise specified on this label.

CIMARRON® X-tra is a broad-spectrum herbicide recommended for use on pastures, rangeland or CRP in most states. Check with your state extension or Dept. of Agriculture before use to be certain CIMARRON® X-tra is registered in your state. Do not use CIMARRON® X-tra in the following counties of Colorado: Alamosa, Conejos, Costilla, Rio Grande, and Saquache.

CIMARRON® X-tra controls weeds by preemergence and postemergence activity. For best results, apply CIMARRON® X-tra to young, actively growing weeds. Weeds hardened off by cold weather or drought stress may not be controlled. The use rate depends upon the weed spectrum and size of weeds at application. The degree and duration of control may depend on the following factors:

- weed spectrum and infestation intensity
- weed size at application
- environmental condition at and following treatment

It is permissible to apply CIMARRON® X-tra to floodplains where surface water is not present, terrestrial areas of deltas, and low lying areas where water is drained but may be isolated in pockets due to uneven or unlevel conditions.

Environmental Conditions and Biological Activity

CIMARRON® X-tra is absorbed through the foliage and roots of broadleaf weeds, rapidly inhibiting their growth. Leaves of susceptible plants appear chlorotic from 1 to 3 weeks after application and the growing point subsequently dies. The final effects on annual weeds are evident about 4 to 6 weeks after application. The ultimate effects on perennial weeds and woody plants occur in the growing seasons following application.

One to two inches of rainfall or sprinkler irrigation (enough to wet the top 2-3 inches of soil profile) is needed to move DuPont™ CIMARRON® X-tra into the weed root zone before the next flush of weeds emerge. The amount of moisture required for sufficient activation increases with crop or weed residue and for finer textured soils. Without sufficient rainfall or sprinkler irrigation to move CIMARRON® X-tra into the weed root zone, weeds that germinate after treatment will not be controlled.

Application of CIMARRON® X-tra provides the best control in vigorously growing grasses that shade competitive weeds. Weed control in areas of thin grass may not be as satisfactory. However, a grass canopy that is too dense at application can intercept spray and reduce weed control.

CIMARRON® X-tra is safe to desirable grass species under normal conditions. However, grasses that are stressed from adverse environmental conditions (such as extremes in temperatures or moisture), abnormal soil conditions, or cultural practices, may be injured by applications of CIMARRON® X-tra. In addition, different species of grass crops may be sensitive to treatment with CIMARRON® X-tra under otherwise normal conditions. Application of CIMARRON® X-tra to these species may result in injury.

In warm, moist conditions, the expression of herbicide symptoms is accelerated in weeds and brush; in cold and/or dry conditions, expression of herbicide symptoms is delayed. In addition, weeds and brush hardened-off by drought stress are less susceptible to CIMARRON® X-tra. Weed and brush control or suppression may be reduced if rainfall, sprinkler irrigation or snowfall occurs within 4 hours after application.

APPLICATION INFORMATION FOR PASTURES AND RANGELAND

Application Timing—Pastures and Rangeland

CIMARRON® X-tra may be used on established native grasses such as bluestems, blue grama, buffalograss and other pasture grasses such as bermudagrass, bluegrass, orchardgrass, brome grass (except Matua brome grass) and fescue. Specific application information on several of these pasture grasses follows.

Do not use on bentgrass or susceptible grass pastures such as timothy, carpetgrass, Matua brome grass or St. Augustine grass.

Applications of CIMARRON® X-tra may cause severe injury to and/or loss of Pensacola bahiagrass, ryegrass (Italian or perennial) and Garrison's creeping foxtail pastures.

Pasture Grass	Minimum time from grass establishment to CIMARRON® X-tra application
Bermudagrass	2 months
Bluegrass, brome grass (except Matua brome grass and orchardgrass)	6 months
Fescue	24 months

Buffalograss Precautions:

Do not use CIMARRON® X-tra on buffalograss that has been established for less than one year or on stands grown for seed production. Do not apply more than 1 oz/acre of CIMARRON® X-tra to buffalograss.

Fescue Precautions:

Note that CIMARRON® X-tra may temporarily stunt fescue, cause it to turn yellow, or cause seedhead suppression. To minimize these symptoms, take the following precautions:

- use only 0.5 oz/acre of CIMARRON® X-tra
- use a non-ionic surfactant at 1/2 to 1 pt per 100 gal of spray solution (1/16 to 1/8% v/v). Do not use a spray adjuvant other than non-ionic surfactant.
- make application later in the spring after the new growth is 5 to 6 inches tall, or in the fall
- do not use surfactant when liquid nitrogen is used as a carrier

The first cutting yields may be reduced due to seedhead suppression resulting from treatment with CIMARRON® X-tra.

Other Pasture and Rangeland Grasses: Varieties and species of forage grasses differ in their tolerance to herbicides. When using CIMARRON® X-tra on a particular grass for the first time, limit use to a small area. If no injury occurs throughout the season, larger acreage may be treated the following season. Broadleaf pasture species, such as alfalfa and clover, are highly sensitive to CIMARRON® X-tra and will be severely stunted or injured by CIMARRON® X-tra.

Application Information for Conservation Reserve Program (CRP)

CIMARRON® X-tra is recommended for the control or suppression of broadleaf weeds in established stands (planted previous year, or earlier) of the following perennial native or improved grasses grown on land enrolled in the Conservation Reserve Program (CRP):

- | | |
|--|---|
| Blue Grama | Orchardgrass |
| Bluestems - big little plains sand WW spar | Sideoats grama |
| Buffalograss | Switchgrass - blackwell |
| Green sprangletop | Wheatgrasses - crested intermediate pubescent slender streambank tall |
| Indiangrass | thickspike western |
| Kleingrass | Wildrye grass - Russian |
| Lovegrasses - atherstone sand weeping wilman | |

Because newly planted CRP grass stands do not sufficiently compete with weeds, and because weed pressure in CRP fields is often severe, performance from DuPont™ CIMARRON® X-tra may not always be satisfactory. An additional herbicide application or mowing may be needed.

Application Timing and Use Rates

CIMARRON® X-tra may be applied postemergence at 0.5 oz/acre or 1.0 oz/acre to labeled grasses listed above that were planted the previous season and are fully tillered.

Weeds and Brush Controlled or Suppressed in Pastures, Rangeland or CRP

Unless otherwise directed in the Specific Weed Problem section of this label, treat when weeds are less than 4" tall or in diameter and are actively growing. Before using CIMARRON® X-tra, carefully consider your crop rotation plans and options. For rotational flexibility, do not treat all of your pasture, rangeland or CRP acres at the same time.

CIMARRON® X-tra at 0.5 oz/acre

Annual marshelder	(common, slimleaf)
Annual sunflower*	Marestail
Bitter sneezeweed	Mayweed chamomile
Blackeyed-Susan	Miners lettuce
Blue/purple mustard*	Mountain snowberry*
Broomweed, common	Musk thistle*
Buckbrush‡	Pigweed (redroot, smooth, tumble)
Bur buttercup (testiculate)	Plains coreopsis
Burclover	Plantain
Buttercup	Prickly lettuce*
Canada thistle*‡	Prostrate knotweed*‡
Carolina geranium	Purple scabious
Coast fiddleneck (tarweed)	Russian thistle*
Common chickweed	Scotch thistle*
Common mullein	Shepherd's-purse
Common purslane	Smallseed falseflax
Common snowberry*	Smartweed (green, ladysthumb, pale)
Common yarrow	Snow speedwell
Conical catchfly	Tansymustard*
Corn gromwell*‡	Treacle mustard
Cowcockle	(Bushy Wallflower)
Curly dock	Tumble/Jim Hill mustard
Cutleaf evening primrose*‡	Volunteer sunflower*
Dandelion	Waterpod
Dogfennel	Western snowberry*
False chamomile	Wild buckwheat*‡
Field pennycress (fanweed)	Wild carrot
Filaree	Wild garlic*
Flixweed*	Wild mustard
Groundsel (common)	Wild sunflower*‡
Henbit	Woolly croton*
Horsemint (beebalm)	
Kochia*	
Lambsquarters	

CIMARRON® X-tra at 1.0 oz/acre

All weeds listed above plus:	Pensacola bahiagrass*
Annual sowthistle	Pigweed (prostrate)
Aster	Pineapple-weed
Bittercress	Plumeless thistle*
Bull thistle*	Redstem filaree
Chicory	Rough fleabane
Clover	Sand sagebrush*
Cocklebur	Seaside arrowgrass
Corn cockle	Sericea lespedeza*
Crown vetch	Silky crazyweed (locoweed)
Flodman thistle*	Sweet clover
Fringed sagebrush*	Wavyleaf thistle*
Goldenrod	Wild lettuce
Hempnettle	Wood sorrel
London rocket	Yankee weed
Maximillion sunflower	Yellowspine thistle*
Pennsylvania smartweed	

CIMARRON® X-tra at 2.0 oz/acre

All weeds listed above plus:	wild roses
Black henbane	Perennial Pepperweed
Blackberry	Poison hemlock
Bouncingbet	Purple loosestrife
Broom snakeweed*	Rosering gaillardia
Buckhorn plantain	Rush skeletonweed*‡
Bull thistle	Salsify
Bur beakchervil	Scouringrush
Common crupina	Snowberry
Common speedwell‡	Spotted knapweed*
Common tansy	St. Johnswort
Dewberry	Teasel
Dyer's woad	Turkey mullein‡
Field bindweed‡	Western salsify
Gorse	Whitetop (hoary cress)
Gunweed	Wild caraway
Halogeton	Wild parsnip
Honeysuckle	Yucca*‡
Houndstongue	
Multiflora rose and other	

* See the **Specific Weed Problems** section.

‡ **Weed suppression** is a reduction in weed competition (reduced population and/or vigor) as visually compared to an untreated area. The degree of suppression varies with the rate used, the size of the weeds, and the environmental conditions following treatment.

SPECIFIC WEED PROBLEMS

Note: For best results, thorough spray coverage of all weed species listed below is very important.

Annual sunflower: Apply CIMARRON® X-tra at 0.5 oz/acre in the spring or early summer, prior to emerged plants reaching 6 inches of new growth.

Blue/Purple Mustard, Flixweed, and Tansymustard: For best results, apply CIMARRON® X-tra at 0.5 oz/acre in tank mixtures with 2,4-D or MCPA postemergence to mustards, but before bloom.

Broom Snakeweed: For best results, apply CIMARRON® X-tra at 2.0 oz/acre in the fall. Applications of CIMARRON® X-tra in the spring will provide suppression only.

Bull Thistle: For control of bull thistle apply CIMARRON® X-tra at 0.5 oz/acre in the spring or early summer prior to flowering, or in the fall after newly emerged plants have reached the rosette stage of growth. Fall applications should be made before the soil freezes.

Canada Thistle: For suppression of emerged Canada thistle foliage, apply DuPont™ CIMARRON® X-tra at 0.5 oz/acre after the majority of thistles have emerged and are actively growing. Applications can be made through flowering and prior to completion of seed development. The application will inhibit the ability of emerged thistles to compete with grass. For suppression of shoot regrowth from underground roots during the season of application, CIMARRON® X-tra should be applied just prior to flower bud formation, through completion of seed development. CIMARRON® X-tra applied early in the spring shortly after thistle emergence will not suppress shoot regrowth from underground roots. Use of CIMARRON® X-tra at 1.0 oz/acre may provide some additional suppression of shoot regrowth from underground roots.

CIMARRON® X-tra may cause a reduction in shoot regrowth from underground roots the following season, however this reduction is inconsistent due to the natural variability in growth from perennial root systems.

Corn Gromwell, Cutleaf Evening Primrose and Prostrate Knotweed: Apply CIMARRON® X-tra at 0.5 oz/acre when weeds are actively growing, are no larger than 2" tall, and when crop canopy will allow thorough coverage. Tank mixing 2,4-D or MCPA with CIMARRON® X-tra can improve results.

Kochia, Russian thistle, Prickly lettuce: Naturally occurring resistant biotypes of these weeds are known to occur. For best results, use CIMARRON® X-tra at 0.5 oz/acre in a tank mix with dicamba (such as "Banvel" or "Clarity") and 2,4-D. CIMARRON® X-tra should be applied in the spring when kochia, Russian thistle, and prickly lettuce are less than 2" tall or 2" across and are actively growing.

Multiflora Rose: For control with broadcast applications, apply CIMARRON® X-tra at 2.0 oz/acre as a broadcast application when multiflora rose is less than 3' tall. Application should be made in the spring, soon after multiflora rose is fully leafed.

Musk Thistle: Apply CIMARRON® X-tra at 0.5 oz/acre in the spring or early summer prior to flowering or in the fall after newly emerged plants have reached the rosette stage of growth. Certain biotypes of musk thistles are less sensitive to CIMARRON® X-tra and may not be controlled. Use of CIMARRON® X-tra at 1.0 oz/acre may provide some additional control of these less sensitive biotypes, but may not achieve acceptable control. Consult with your local DuPont representative, dealer or applicator for specific use rate and tank mix recommendations for your area. Fall applications should be made before the soil freezes.

Pensacola bahiagrass control in established Bermudagrass pasture: Apply CIMARRON® X-tra at 1.0 oz/acre after greenup in the spring but before bahiagrass seedhead formation. Application should be made when moisture is sufficient to enhance grass growth.

CIMARRON® X-tra is very effective for removal of bahiagrass from bermudagrass pastures. In highly infested pastures, the use of CIMARRON® X-tra can clear the areas of useful forage until the bermudagrass has time to cover the area. Therefore, CIMARRON® X-tra treatments should be spread out over a period of years. Do not apply to an entire farm or ranch in one

year. Fertilization (particularly with nitrogen and potassium) and/or replanting may accelerate the process of reestablishment of bermudagrass. Under heavy bahiagrass pressure, grazing pressure, or adverse weather conditions (heat and drought), bahiagrass regrowth may occur.

CIMARRON® X-tra should not be applied in liquid fertilizer solutions for Pensacola bahiagrass control, as poor control and/or regrowth may occur.

CIMARRON® X-tra should not be used for the control of common or Argentine bahiagrass.

Plumeless Thistle: For control of plumeless thistle apply CIMARRON® X-tra at 1.0 oz/acre in the spring or early summer prior to flowering or in the fall after newly emerged plants have reached the rosette stage of growth. Fall applications should be made before the soil freezes.

Rush skeletonweed: For best results, apply CIMARRON® X-tra at 2.0 oz/acre with 8 fluid ounces of dicamba (such as "Banvel" or "Clarity") and 16 fluid ounces of 2,4-D per acre.

Sagebrush (Sand, Fringed): For control of sagebrush, apply CIMARRON® X-tra at 1.0 oz/acre after the plants are actively growing. Applications can be made in the spring or early summer after 6 inches of new growth has occurred. Tank mixtures with 2,4-D ester improve control (refer to the Tank Mixtures section of this label for additional information).

Scotch Thistle: For suppression of scotch thistle apply CIMARRON® X-tra at 0.5 oz/acre in the spring or early summer prior to flowering.

Sericea lespedeza: For best results, apply CIMARRON® X-tra at 1.0 oz/acre beginning at flower bud initiation through the full bloom stage of growth. Do not make applications if drought conditions exist at intended time of application.

Snowberry (Western, Common, Mountain): For control of snowberry, apply CIMARRON® X-tra at 0.5 oz/acre after the plants are actively growing. Applications can be made throughout the growing season but before fall defoliation. Tank mixtures with 2,4-D ester improve control (refer to the Tank Mixtures section of this label for additional information).

Spotted Knapweed: For best results, apply CIMARRON® X-tra at 2.0 oz/acre with 8 fluid ounces of dicamba (such as "Banvel" or "Clarity") and 16 fluid ounces of 2,4-D per acre.

Sunflower (wild or volunteer): Apply CIMARRON® X-tra at 0.5 oz/acre plus 2,4-D or MCPA after the majority of sunflowers have emerged, are 2" to 4" tall and are actively growing.

Wild Buckwheat: For best results, apply CIMARRON® X-tra at 0.5 oz/acre plus 2,4-D or MCPA when plants have no more than 3 true leaves (not counting the cotyledons). If plants are not actively growing, delay treatment until environmental conditions favor active weed growth.

Wild Garlic: For best results, apply CIMARRON® X-tra at 0.5 oz/acre in the early spring when wild garlic is less than 12" tall with 2" to 4" of new growth.

Woolly Croton: For best results, apply CIMARRON® X-tra at 0.5 oz/acre in the late spring or early summer from cotyledon through 2 true leaf stage.

Yellowspine Thistle, Wavyleaf Thistle, Flodman Thistle: For control of yellowspine thistle, wavyleaf thistle, or flodman thistle apply CIMARRON® X-tra at 1.0 oz/acre in the spring or early summer prior to flowering.

Yucca: For best results, apply DuPont™ CIMARRON® X-tra at 2.0 oz/acre plus 2,4-D, dicamba, dicamba plus 2,4-D, or "Remedy" from two weeks before blooming to two weeks after blooming.

Spray Adjuvants

Unless otherwise recommended, applications of CIMARRON® X-tra must include either a crop oil concentrate or a nonionic surfactant. Consult local DuPont fact sheets, technical bulletins, and service policies prior to using other adjuvant systems. If another herbicide is tank mixed with CIMARRON® X-tra, select adjuvants authorized for use with both products. Products must contain only EPA-exempt ingredients (40 CFR 1001).

Petroleum Crop Oil Concentrate (COC) or Modified Seed Oil (MSO)

- Apply at 1% v/v (1 gal per 100 gal spray solution) or 2% under arid conditions.
- Oil adjuvants must contain at least 80% high quality, petroleum (mineral) or modified vegetable seed oil with at least 15% surfactant emulsifiers.

Nonionic Surfactant (NIS)

- Apply at 0.25% v/v (1 qt per 100 gal spray solution) or 0.5% under arid conditions.
- Surfactant products must contain at least 60% nonionic surfactant with a hydrophilic/lipophilic balance (HLB) greater than 12.

Special Adjuvant Types

- Combination adjuvant products may be used at doses that provide the required amount of NIS, COC, MSO. Consult product literature for use rates and restrictions.
- In addition to the adjuvants specified above, other adjuvant types may be used if they provide the same functionality and have been evaluated and approved by DuPont. Consult separate DuPont technical bulletins for detailed information before using adjuvant types not specified on this label.

Exceptions: On Fescue pastures use no more than 1/2 to 1 pint non-ionic surfactant per 100 gals.

Antifoaming agents may be used if needed.

Ammonium Nitrogen Fertilizer

- Use up to 2% v/v of a high-quality urea ammonium nitrate (UAN), such as 28% N or 32% N, or up to 17 lb/acre of a spray grade ammonium sulfate (AMS).

Do not use low rates of liquid fertilizer as a substitute for spray adjuvants.

Ground Application

To obtain optimum spray distribution and thorough coverage, use flat-fan or low-volume flood nozzles. For flat-fan nozzles, use at least 10 GPA for broadcast applications to pasture, rangeland or CRP.

For flood nozzles on 30" spacings, use at least 10 gallons per acre (GPA), flood nozzles no larger than TK10 (or equivalent), and a pressure of at least 30 pounds per square inch (psi). For 40" nozzle spacings, use at least 13 GPA; for 60" spacings, use

at least 20 GPA. It is essential to overlap the nozzles 100% for all spacings.

With "Raindrop RA" nozzles, use at least 30 GPA and ensure that nozzle spray patterns overlap 100%.

Use 50-mesh screens or larger.

Aerial Application

Use nozzle types and arrangements that provide optimum spray distribution and maximum coverage.

Use a minimum of 3 GPA.

When applying CIMARRON® X-tra by air in areas adjacent to sensitive crops, use solid stream nozzles oriented straight back. Adjust the swath to avoid spray drift damage to sensitive crops downwind and/or use ground equipment to treat the border edge of fields. See the **Spray Drift Management** section of this label.

TANK MIXTURES

When tank mixing CIMARRON® X-tra, use the most restrictive label limitations for each product used in the mix.

With Insecticides and Fungicides

CIMARRON® X-tra may be tank mixed or used sequentially with insecticides and fungicides registered for use on pastures, rangeland or CRP. However, under certain conditions (drought stress or cold weather), tank mixes or sequential applications of CIMARRON® X-tra with organophosphate insecticides (such as parathion) may produce temporary grass yellowing or, in severe cases, grass injury. The potential for grass injury is greatest when wide fluctuations in day/night temperatures occur just prior to or soon after application. Test these mixtures in a small area before treating large areas. Do not use CIMARRON® X-tra plus Malathion, since grass injury will result.

Herbicide Tank Mixtures for Pastures or Rangeland:

CIMARRON® X-tra may be tank mixed with other suitable registered herbicides to control weeds listed as **Weeds Suppressed**, weeds resistant to CIMARRON® X-tra, or weeds not listed under **Weeds Controlled**. Read and follow all manufacturer's label recommendations for the companion herbicide. If those recommendations conflict with this label, do not tank mix the herbicide with CIMARRON® X-tra.

CIMARRON® X-tra can be applied in a tank mix with one of the following products. Refer to companion herbicide labels to confirm that the product is labeled for control of the weeds listed above and is registered for use in your state.

Product	Rate (oz product/A)
"Grazon" P+D	8 to 32
"Tordon" 22K	4 to 16
"Weedmaster"	8 to 32
"Remedy"	8
Product	Rate (oz A.I./A)
2,4-D	8 to 16
Dicamba (such as "Banvel" or "Clarity")	2 to 16
2,4-D + Dicamba	3 + 1 to 12 + 4

Herbicide Tank Mixtures for CRP:

Preplant

DuPont™ CIMARRON® X-tra may be tank mixed with glyphosate (such as DuPont™ Glyphosate or "Roundup UltraMax") as a pre-plant (prior to the planting of CRP grasses) treatment to control broadleaf and grassy weeds. When using a glyphosate tank mix, allow at least 7 days after application before planting grasses. Refer to glyphosate containing product labels and fact sheets for all use instructions, label rates, weed control claims, warnings and precautions..

Postemergence

For best weed control performance in CRP, use CIMARRON® X-tra in a tank mix with 2,4-D (ester formulations perform best) or dicamba (such as "Banvel" or "Clarity").

CIMARRON® X-tra can be tank mixed with 2,4-D at 1/4 lb a.i./A for all labeled grasses larger than the 5-leaf stage. For fully tillered stands, up to 1/2 lb a.i./A of 2,4-D may be used. A spray adjuvant may be added. However, the addition of spray adjuvant may increase the chance of grass injury.

CIMARRON® X-tra can also be tank mixed with dicamba (such as "Banvel" or "Clarity"). Use not more than 1/8 to 1/4 lb a.i./A of dicamba plus CIMARRON® X-tra after majority of grasses are in the 3-leaf stage. In established grasses (2nd year stands), use not more than 1/4 to 1/2 lb a.i./A dicamba plus CIMARRON® X-tra. A spray adjuvant may be added. However, the addition of spray adjuvant may increase the chance of grass injury.

With Liquid Nitrogen Solution Fertilizer

Liquid nitrogen fertilizer solutions may be used as a carrier in place of water. Run a tank mix compatibility test before mixing CIMARRON® X-tra in fertilizer solution.

CIMARRON® X-tra must first be slurried with clean water and then added to liquid nitrogen solutions (e.g., 28-0-0, 32-0-0). Ensure that the agitator is running while the CIMARRON® X-tra is added.

Use of this mixture may result in temporary grass yellowing and stunting.

If using low rates of liquid nitrogen fertilizer (between 5 and 50% of the spray solution volume) in the spray solution, the addition of a non-ionic surfactant is necessary. Add surfactant at 1/4 pt per 100 gal of spray solution (0.03%). Do not use a spray adjuvant other than non-ionic surfactant.

When using high rates of liquid nitrogen fertilizer (equal to or greater than 50% of the spray solution volume) in the spray solution, adding a spray adjuvant increases the risk of grass injury. Consult your agricultural dealer, consultant, fieldman, or DuPont representative for a specific recommendation before adding an adjuvant to these tank mixtures.

If 2,4-D or MCPA is included with CIMARRON® X-tra and fertilizer mixture, ester formulations tend to be more compatible (See manufacturer's label). Do not add a spray adjuvant when using CIMARRON® X-tra in tank mix with 2,4-D ester and liquid nitrogen fertilizer solutions greater than 5% of the spray solution volume.

The use of liquid nitrogen fertilizer solutions greater than 5% of the spray solution volume with CIMARRON® X-tra at rates greater than 0.5 oz/acre may cause grass injury.

Do not use low rates of liquid fertilizer as a substitute for a spray adjuvant.

Do not tank mix CIMARRON® X-tra with liquid fertilizer solutions with a pH less than 3.0.

Minimum Rotational Intervals

Minimum rotation intervals are determined by the rate of breakdown of CIMARRON® X-tra applied. CIMARRON® X-tra breakdown in the soil is affected by soil pH, presence of soil microorganisms, soil temperature, and soil moisture. Low soil pH, high soil temperature, and high soil moisture increase CIMARRON® X-tra breakdown in soil, while high soil pH, low soil temperature, and low soil moisture slow CIMARRON® X-tra breakdown. Of these 3 factors, only soil pH remains relatively constant. Soil temperature, and to a greater extent, soil moisture, can vary significantly from year to year and from area to area. For this reason, soil temperatures and soil moisture should be monitored regularly when considering crop rotations.

Before using CIMARRON® X-tra, carefully consider your crop rotation plans and options. For rotational flexibility, do not treat all of your pasture, rangeland or CRP acres at the same time.

Soil pH Limitations

CIMARRON® X-tra should not be used on soils having a pH above 7.9, because soil residual activity could extend crop rotation intervals beyond normal. Under certain conditions, CIMARRON® X-tra could remain active in the soil for 34 months or more, injuring subsequent crops.

Checking Soil pH

Before using CIMARRON® X-tra, determine the soil pH of the areas of intended use. To obtain a representative pH value for the test area, take several 0" to 4" samples from different areas of the field and analyze them separately. Consult local extension publications for additional information on recommended soil sampling procedures.

BIOASSAY

A field bioassay must be completed before rotating to any crop or grass species/variety not listed in the above Rotation Intervals Table, or if the soil pH is not in the specified range, or if the use rate applied is not specified in the table. To conduct a field bioassay, grow test strips of the crop(s) or grass(es) you plan to grow in fields previously treated with CIMARRON® X-tra. Crop or grass response to the bioassay will indicate whether or not to rotate to the crop(s) or grass(es) grown in the test strips. If a field bioassay is planned, check with your local Agricultural dealer or DuPont representative for information detailing the field bioassay procedure.

GRAZING/HAYING

There are no grazing or hay harvest restrictions for DuPont™ CIMARRON® X-TRA when applied to range, pasture, CRP, and/or undesirable vegetation in uncultivated areas (fence rows, farmyards, and rights-of-way) which are adjacent to, or pass through or transect, treated pastures, rangeland, or CRP.

Coveralls and shoes plus socks must be worn if cutting within 4 hours of treatment.

CROP ROTATION

Rotation Intervals in Pasture, Rangeland or CRP for Overseeding and Renovation

Location	Crop or Grass Species	DuPont™ CIMARRON® X-tra Rate	Minimum Rotation Interval (months) *
ALL AREAS WITH SOIL PH OF 7.5 OR LESS	Russian wildrye	0.5 oz/acre 1.0 oz/acre	2
	Green needlegrass, switchgrass, sheep fescue, meadow brome, smooth brome, red fescue, Russian wildrye	0.5 oz/acre 1.0 oz/acre 2.0 oz/acre	2
	Switchgrass	0.5 oz/acre 1.0 oz/acre 2.0 oz/acre	3
	Timothy	0.5 oz/acre 1.0 oz/acre	2
		2.0 oz/acre	4
	Meadow foxtail	0.5 oz/acre 1.0 oz/acre	3
		2.0 oz/acre	4
	Alta fescue	0.5 oz/acre 1.0 oz/acre	2
		2.0 oz/acre	3
	Orchardgrass	0.5 oz/acre 1.0 oz/acre	2
2.0 oz/acre		3	
ALL AREAS WITH SOIL PH OF 7.9 OR LESS	Mountain brome, blue grama thickspike wheatgrass, western wheatgrass	0.5 oz/acre 1.0 oz/acre 2.0 oz/acre	2
	Alkali sacaton	0.5 oz/acre 1.0 oz/acre	2
		2.0 oz/acre	3
	Sideoats grama, switchgrass	0.5 oz/acre 1.0 oz/acre 2.0 oz/acre	4
	Big bluestem	0.5 oz/acre 1.0 oz/acre	3
	Wheat (except durum)	0.5 oz/acre	4
	Durum, wheat or oat	0.5 oz/acre	16
Barley	0.5 oz/acre	24	

* The minimum rotation interval represents the period of time from the last application to the anticipated date of the next planting.

MIXING INSTRUCTIONS

1. Fill the tank 1/4 to 1/3 full of water (If using liquid nitrogen fertilizer solution in place of water, see Tank Mixtures sections for additional details).
2. While agitating, add the required amount of CIMARRON® X-tra.
3. Continue agitation until the CIMARRON® X-tra is fully dispersed, at least 5 minutes.
4. Once the CIMARRON® X-tra is fully dispersed, maintain agitation and continue filling tank with water. CIMARRON® X-tra should be thoroughly mixed with water before adding any other material.
5. As the tank is filling, add tank mix partners (if desired) then add the necessary volume of spray adjuvants. Always add spray adjuvants last.
6. If the mixture is not continuously agitated, settling will occur. If settling occurs, thoroughly re-agitate before using.

7. Apply CIMARRON® X-tra spray mixture within 24 hours of mixing to avoid product degradation.
8. If CIMARRON® X-tra and a tank mix partner are to be applied in multiple loads, pre-slurry the CIMARRON® X-tra in clean water prior to adding to the tank. This will prevent the tank mix partner from interfering with the dissolution of the CIMARRON® X-tra.

Do not use CIMARRON® X-tra with spray additives that reduce the pH of the spray solution to below 3.0.

SPRAY EQUIPMENT

For specific application equipment, refer to the manufacturer's recommendations for additional information on GPA, pressure, speed, nozzle types and arrangements, nozzle heights above the target canopy, etc.

Be sure to properly calibrate air or ground equipment before application. Select a spray volume and delivery system that will ensure thorough coverage and a uniform spray pattern with minimum drift. Use higher spray volumes to obtain better coverage when the crop canopy is dense. Avoid swath

overlapping, and shut off spray booms while starting, turning, slowing, or stopping to avoid crop injury.

Do not make applications using equipment and/or spray volumes or under weather conditions that might cause spray to drift onto nontarget sites. For additional information on spray drift, refer to the **Spray Drift Management** section of the label.

Continuous agitation is required to keep DuPont™ CIMARRON® X-tra in suspension.

SPRAYER CLEANUP

Spray equipment must be clean before CIMARRON® X-tra is sprayed. Follow the cleanup procedures specified on the labels of previously applied products. If no directions are provided, follow the six steps outlined in **After Spraying CIMARRON® X-tra** section of this label.

At the End of the Day

When multiple loads of CIMARRON® X-tra herbicide are applied, it is recommended that at the end of each day of spraying, the interior of the tank be rinsed with fresh water and then partially filled, and the boom and hoses flushed. This will prevent the buildup of dried pesticide deposits that can accumulate in the application equipment.

After Spraying CIMARRON® X-tra and Before Spraying Crops Other Than Pasture, Rangeland or CRP

To avoid subsequent injury to desirable crops, thoroughly clean all mixing and spray equipment immediately following applications of CIMARRON® X-tra as follows:

1. Drain tank; thoroughly rinse spray tanks, boom, and hoses with clean water. Loosen and physically remove any visible deposits.
 2. Fill the tank with clean water and 1 gal of household ammonia* (contains 3% active) for every 100 gal of water. Flush the hoses, boom, and nozzles with the cleaning solution. Then add more water to completely fill the tank. Circulate the cleaning solution through the tank and hoses for at least 15 min. Flush the hoses, boom, and nozzles again with the cleaning solution, and then drain the tank.
 3. Remove the nozzles and screens and clean separately in a bucket containing cleaning agent and water.
 4. Repeat step 2.
 5. Rinse the tank, boom, and hoses with clean water.
 6. If only Ammonia is used as a cleaner, the rinsate solution may be applied back to the crop(s) recommended on this label. Do not exceed the maximum-labeled use rate. If other cleaners are used, consult the cleaner label for rinsate disposal instructions. If no instructions are given, dispose of the rinsate on site or at an approved waste disposal facility.
- * Equivalent amounts of an alternate-strength ammonia solution or a DuPont-approved cleaner can be used in the cleanout procedure. Carefully read and follow the individual cleaner instructions. Consult your agricultural dealer, applicator, or DuPont representative for a listing of approved cleaners.

Notes:

1. Attention: Do not use chlorine bleach with ammonia, as dangerous gases will form. Do not clean equipment in an enclosed area.
2. Steam-cleaning aerial spray tanks is recommended prior to performing the above cleanout procedure to facilitate the removal of any caked deposits.
3. When CIMARRON® X-tra is tank mixed with other pesticides, all required cleanout procedures should be examined and the most rigorous procedure should be followed.
4. In addition to this cleanout procedure, all precleanout guidelines on subsequently applied products should be followed as per the individual labels.
5. Where routine spraying practices include shared equipment frequently being switched between applications of CIMARRON® X-tra and applications of other pesticides to CIMARRON® X-tra sensitive crops during the same spray season, it is recommended that a sprayer be dedicated to CIMARRON® X-tra to further reduce the chance of crop injury.

SPRAY DRIFT MANAGEMENT

The interaction of many equipment and weather-related factors determines the potential for spray drift. The user is responsible for considering all these factors when making application decisions. Follow the additional precautions below to minimize the potential for spray drift.

All aerial and ground application equipment must be properly maintained and calibrated using appropriate carriers.

AVOIDING SPRAY DRIFT IS THE RESPONSIBILITY OF THE APPLICATOR.

Drift Control Adjuvants

A drift control adjuvant may be used to reduce the potential for drift. However, because it is the combined physical-chemical properties of all the ingredients in the spray mix that can determine drift potential, the applicator must confirm that the drift control adjuvant used is having the desired effect with the tank mix that is being applied. If a drift control adjuvant is used, follow the use directions and precautions on the manufacturer's label. Do not use an adjuvant which increases viscosity with application systems that cannot accommodate viscous sprays.

Ground Application: With ground equipment, spray drift can be lessened by keeping the spray boom as low as possible (i.e., a release height of 4 feet or less above the application target); by applying 10 gallons or more of spray per acre; by keeping the operating spray pressures at the manufacturer's recommended minimum pressures for the specific nozzle type used; and by spraying when the wind velocity is low (follow all applicable state regulations).

Do not make ground applications within a surface temperature inversion when applying near an area requiring protection to avoid an unreasonable adverse effect. Applicators may determine presence of an inversion by noting the presence of ground fog, light variable wind, or layering of smoke and dust. Be particularly alert to the potential for a surface temperature inversion when winds are calm.

Direct the sprays no higher than the tops of target vegetation, and maintain spray pressures at levels which provide coarse to very coarse spray droplets to minimize drift.

Aerial Application: The following drift management requirements must be followed to avoid off-target drift movement from aerial applications:

1. The distance between the outer most operating nozzles on the boom must not exceed 75% of the wingspan. For helicopters, use a boom length and position that prevents droplets from entering the rotor vortices.
2. Nozzles should always point backward parallel with the air stream.

Where states have more stringent regulations, they must be observed. The applicator should be familiar with and take into account the information presented below.

IMPORTANCE OF DROPLET SIZE

Since the most effective way to reduce drift potential is to apply large droplets, equipment producing a coarse to very coarse droplet spectrum must be used when applying this product. The best drift management strategy is to apply the coarsest drop size spectrum that provides sufficient coverage and control. The presence of sensitive species nearby, the environmental conditions, and pest pressure may affect how an applicator balances drift control and coverage. **APPLYING LARGER DROPLETS REDUCES DRIFT POTENTIAL, BUT WILL NOT PREVENT DRIFT IF APPLICATIONS ARE MADE IMPROPERLY OR UNDER UNFAVORABLE ENVIRONMENTAL CONDITIONS!**

See "WIND", "TEMPERATURE AND HUMIDITY", and "SURFACE TEMPERATURE INVERSIONS" sections of this label.

Controlling Droplet Size – Ground Application

- **Volume** - Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets.
- **Pressure** - Do not exceed the manufacturer's recommended pressures. Use the lower spray pressures recommended for the nozzle. Higher pressure generally reduces droplet size and does not improve canopy penetration. **WHEN HIGHER FLOW RATES ARE NEEDED, USE A HIGHER-CAPACITY NOZZLE INSTEAD OF INCREASING PRESSURE.**
- **Nozzle Type** - Use a nozzle type according to manufacturer's specifications which is designed for the intended application, and that produces a coarse to very coarse droplet size spectrum. With most nozzle types, narrower spray angles produce larger droplets. To further reduce drift, low-drift or drift reducing nozzles should be used.

Controlling Droplet Size - Aircraft

- **Number of Nozzles** - Use the minimum number of nozzles that provide uniform coverage.
- **Nozzle Orientation** - For some nozzle types, such as solid streams, orienting nozzles so that the spray is emitted backwards, parallel to the air stream minimizes the effects of air shear and will produce a coarser droplet spectrum than other orientations. For applications of this product, nozzles

must be oriented in a manner that results in the application of a coarse to very coarse droplet size spectrum.

- **Nozzle Type** - Use a nozzle type according to manufacturer's specifications which is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Solid stream and other drift reducing nozzles should be used.

BOOM LENGTH AND HEIGHT

- **Boom Height (ground)** - Setting the boom at the lowest referenced height (if specified) which provides uniform coverage reduces the exposure of droplets to evaporation and wind. The boom should remain level with the crop and have minimal bounce. Apply at a height no greater than 4 feet above the top of the largest plants.
- **Application Height (aircraft)** - Apply at a height no greater than 10 feet above the top of the largest plants, unless a greater height is required for aircraft safety. Making applications at the lowest height that is safe reduces exposure of droplets to evaporation and wind.
- **Boom Length (aircraft)** - The distance between the outer-most operating nozzles on the boom must not exceed 3/4 (75%) of the wingspan - longer booms increase drift potential. For helicopters, use a boom length and position that prevents droplets from entering the rotor vortices.

SWATH ADJUSTMENT

When applications are made with a crosswind, the swath will be displaced downwind. Therefore, on the up and downwind edges of the field, the applicator must compensate for this displacement by adjusting the path of the application equipment upwind. Swath adjustment distance should increase with increasing drift potential (higher wind, smaller drops, etc.).

WIND (GROUND AND AERIAL APPLICATION)

Drift potential is lowest with a sustained wind of 2-10 mph. However, many factors, including droplet size and equipment type, determine drift potential at any given wind speed. Application should be avoided during gusty conditions, and when winds are below 2 mph due to variable wind direction and high potential for a temperature inversion. Avoid applying during calm conditions which may be conducive to air inversions.

Note: Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they affect spray drift.

TEMPERATURE AND HUMIDITY (GROUND AND AERIAL APPLICATIONS)

When making applications in low relative humidity, set up equipment to produce larger droplets to compensate for evaporation. Droplet evaporation is most severe when conditions are both hot and dry.

SURFACE TEMPERATURE INVERSIONS (GROUND AND AERIAL APPLICATIONS)

Applications must not occur during a local, surface temperature inversion because drift potential is high. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain in a concentrated cloud. This cloud can move in unpredictable directions due to the light variable winds which are common during inversions. Temperature inversions are characterized by increasing temperatures with height and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of the smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

SHIELDED SPRAYERS (GROUND APPLICATION)

Shielding the boom or individual nozzles can reduce the effects of wind. However, it is the responsibility of the applicator to verify that the shields are preventing drift and not interfering with uniform deposition of the product.

Do not apply this product in a way that will contact workers or other people, either directly or through drift. Only protected handlers may be in the area during application.

SENSITIVE AREAS

This product should be applied only when the potential for drift to adjacent sensitive areas (e.g. residential areas, bodies of water, known habitat for threatened or endangered species, non-target crops) is minimal (e.g. when wind is blowing away from the sensitive areas). Small quantities of spray may seriously injure susceptible crops either during active growth periods or dormancy.

WEED RESISTANCE

Biotypes of certain weeds listed on this label are resistant to DuPont™ CIMARRON® X-tra and other herbicides with the same mode of action*, even at exaggerated application rates. Biotypes are naturally occurring individuals of a species that are identical in appearance but have slightly different genetic compositions; the mode of action of a herbicide is the chemical interaction that interrupts a biological process necessary for plant growth and development. If weed control is unsatisfactory, it may be necessary to retreat problem areas using a product with a different mode of action, such as postemergence broadleaf and/or grass herbicides. If resistant weed biotypes such as kochia, prickly lettuce, and Russian thistle are suspected or known to be present use a tankmix partner with CIMARRON® X-tra to help control these biotypes, or use a planned herbicide rotation program where other residual broadleaf herbicides having different modes of action are used.

* Naturally occurring weed biotypes that are resistant to ALS inhibitor herbicides (such as "Amber" herbicide) may also be resistant to CIMARRON® X-tra.

PRECAUTIONS

- CIMARRON® X-tra may cause injury to desirable trees and plants when contacting their roots, stems or foliage. These plants are most sensitive to CIMARRON® X-tra during their development or growing stage. FOLLOW THE PRECAUTIONS IN THIS LABEL WHEN USING CIMARRON® X-tra.
- Injury to or loss of desirable trees or vegetation may result from failure to observe the following:
 - Do not apply CIMARRON® X-tra, or drain or flush equipment on or near desirable trees or other plants, or on areas where their roots may extend, or in locations where the chemical may be washed or moved into contact with their roots.
 - Do not use on lawns, walks, driveways, tennis courts, golf courses, athletic fields, commercial sod operations, or other high-maintenance, fine turfgrass areas, or similar areas.
 - Do not use on grasses grown for seed.
- Do not contaminate irrigation ditches or water used for domestic purposes.
- Do not apply to irrigated land where the tailwater will be used to irrigate crops.
- Do not apply to frozen ground as surface runoff may occur.
- Do not apply to snow-covered ground.
- Grass species or varieties may differ in their response to various herbicides. DuPont recommends that you first consult your state experiment station, university, or extension agent as to sensitivity to any herbicide. If no information is available, limit the initial use of CIMARRON® X-tra to a small area. Components in a grass seed mixture will vary in tolerance to CIMARRON® X-tra so the final stand may not reflect the seed ratio.
- Under certain conditions such as heavy rainfall, high pH, prolonged cold weather, or wide fluctuations in day/night temperatures prior to or soon after CIMARRON® X-tra application, temporary discoloration and/or grass injury may occur. DuPont™ CIMARRON® X-tra should not be applied to grass that is stressed by severe weather conditions, drought, low fertility, water-saturated soil, disease, or insect damage, as grass injury may result. Severe winter stress, drought, disease, or insect damage before or following application also may result in grass injury.
- Applications of CIMARRON® X-tra to pastures, rangeland or CRP undersown with legume crops may cause severe injury to the legumes. Legumes in a seeding mixture may be severely injured or killed following an application of CIMARRON® X-tra.
- Treatment of powdery, dry soil or light, sandy soil when there is little likelihood of rainfall soon after treatment may result in off target movement and possible damage to susceptible crops when soil particles are moved by wind or water. Injury to crops may result if treated soil is washed, blown, or moved onto land used to produce crops. Injury may be more severe when the crops are irrigated. Do not apply CIMARRON® X-tra when these

conditions are identified and powdery, dry soil or light or sandy soil are known to be prevalent in the area to be treated.

- Applications made where runoff water flows onto agricultural land may injure crops. Applications made during periods of intense rainfall, to soils saturated with water, or soils through which rainfall will not readily penetrate may result in runoff and movement of DuPont™ CIMARRON® X-tra. Treated soil should be left undisturbed to reduce the potential for CIMARRON® X-tra movement by soil erosion due to wind or water.
- For ground applications applied to weeds when dry, dusty field conditions exist, control of weeds in wheel track areas may be reduced.
- Do not apply more than the equivalent of 1 ounce of chlorsulfuron per acre per year.
- Do not apply more than the equivalent of 1 ounce of metsulfuron methyl per acre per year.
- Avoid disturbing (e.g. cultivating or mowing) treated areas for at least 7 days following application.

STORAGE AND DISPOSAL

Pesticide Storage: Store product in original container only. Do not contaminate water, other pesticides, fertilizer, food or feed in storage.

Pesticide Disposal: Do not contaminate water, food, or feed by disposal. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: For Plastic Containers: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke. **For Fiber Sacks:** Completely empty fiber sack by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into manufacturing or application equipment. Then dispose of sack in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

Container Refilling and Disposal (For Containers up to 250 gal): This is a refillable container. If the container is to be refilled, do not rinse with any material or introduce any pesticide other than CIMARRON® X-tra. Reseal and return the container to any authorized DuPont refilling facility. If the container is not to be refilled, triple rinse (or equivalent) and offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or by open burning, if allowed by state and local authorities. If burned, keep out of smoke.

For minor spills, leaks, etc., follow all precautions indicated on this label and clean up immediately. Take special care to avoid contamination of equipment and facilities during cleanup procedures and disposal of wastes. In the event of a major spill, fire or other emergency, call 1-800-441-3637 day or night.

Container Disposal for Bulk Containers: When this container is empty, replace the cap and seal all openings that have been opened during use, and return the container to the point of purchase or to a designated location named at time of purchase of this product. The container must only be refilled with this pesticide product. **DO NOT REUSE THE CONTAINER FOR ANY OTHER PURPOSE.** Prior to refilling, inspect carefully for damage such as cracks, punctures, abrasions, worn-out threads and closure devices. Check for leaks after refilling and before transporting. Do not transport if this container is damaged or leaking. If the container is damaged, leaking or obsolete, contact DuPont at 1-800-441-3637. If not returned to the point of purchase or to a designated location, triple rinse emptied container and offer for recycling. Disposal of this container must be in compliance with state and local regulations.

For minor spills, leaks, etc., follow all precautions indicated on this label and clean up immediately. Take special care to avoid contamination of equipment and facilities during cleanup procedures and disposal of wastes. In the event of a major spill, fire or other emergency, call 1-800-441-3637 day or night.

NOTICE TO BUYER: Purchase of this material does not confer any rights under patents of countries outside of the United States.

The DuPont Oval Logo, DuPont™ and CIMARRON® are trademarks or registered trademarks of E. I. duPont de Nemours and Company.

"Raindrop RA" is a trademark of Delavan Corp.

"Amber" is a trademark of Syngenta Crop Protection Inc.

"Weedmaster" and "Clarity" are trademarks of BASF Corp.

"Banvel" is a trademark of Micro Flo Company LLC

"Grazon", "Remedy", "Tordon" are trademarks of Dow AgroSciences

"Roundup UltraMax" is a trademark of Monsanto

D - 1274 103107

LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read This Limitation of Warranty and Liability Before Buying or Using This Product. If the Terms Are Not Acceptable, Return the Product at Once, Unopened, and the Purchase Price Will Be Refunded.

It is impossible to eliminate all risks associated with the use of this product. Such risks arise from weather conditions, soil factors, off target movement, unconventional farming techniques, presence of other materials, the manner of use or application, or other unknown factors, all of which are beyond the control of DuPont. These risks can cause: ineffectiveness of the product; crop injury, or; injury to non-target crops or plants.

DuPont does not agree to be an insurer of these risks. **TO THE FULLEST EXTENT PERMITTED BY LAW, WHEN YOU BUY OR USE THIS PRODUCT, YOU AGREE TO ACCEPT THESE RISKS.**

DuPont warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for the purpose stated in the Directions for Use, subject to the inherent risks described above, when used in accordance with the Directions for Use under normal conditions.

DUPONT MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR OF MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

TO THE FULLEST EXTENT PERMITTED BY LAW, IN NO EVENT SHALL DUPONT OR SELLER BE LIABLE FOR ANY INCIDENTAL, CONSEQUENTIAL OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. BUYER'S OR USER'S BARGAINED-FOR EXPECTATION IS CROP PROTECTION. THE EXCLUSIVE REMEDY OF THE USER OR BUYER AND THE EXCLUSIVE LIABILITY OF DUPONT OR SELLER, FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY OR CONTRACT, NEGLIGENCE, TORT OR STRICT LIABILITY), WHETHER FROM FAILURE TO PERFORM OR INJURY TO CROPS OR OTHER PLANTS, AND RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT, OR AT THE ELECTION OF DUPONT OR SELLER, THE REPLACEMENT OF THE PRODUCT.

DuPont or its Ag Retailer must have prompt notice of any claim so that an immediate inspection of buyer's or user's growing crops can be made. Buyer and all users shall promptly notify DuPont or a DuPont Ag Retailer of any claims, whether based on contract, negligence, strict liability, other tort or otherwise or be barred from any remedy.

This Limitation of Warranty and Liability may not be amended by any oral or written agreement.

For product information call: 1-888-6-DUPONT

Internet address: www.cropprotection.dupont.com

© 2005-2007 E. I. du Pont de Nemours and Company, 1007 Market Street, Wilmington, DE 19898

All rights reserved.