


**DuPont Crop
Protection**



SUPPLEMENTAL LABELING

**DUPONT™ LOROX® DF
HERBICIDE**

**FOR USE ON PARSLEY GROWN
FOR SEED IN OREGON AND
WASHINGTON**

DUPONT™ LOROX® DF HERBICIDE

EPA Reg. No. 352-686

**FOR USE ON PARSLEY GROWN FOR SEED
IN THE STATES OF OREGON AND WASHINGTON**

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

This bulletin contains new or supplemental instructions for use of this product which does not appear on the package label.

Follow the instructions carefully.

This labeling must be in the possession of the user at time of pesticide application.

**IMPORTANT
BEFORE USING LOROX® DF HERBICIDE,
READ AND FOLLOW ALL APPLICABLE
DIRECTIONS, RESTRICTIONS, AND
PRECAUTIONS ON THE EPA-REGISTERED
LABEL.**

GENERAL INFORMATION

DuPont™ LOROX® DF herbicide is recommended for control of weeds in parsley grown for seed.

For best results apply preemergence to weeds or early postemergence while weeds are small. Consult the LOROX® DF label for list of weeds controlled. If weeds are up at time of application, add a non-ionic surfactant at 1 quart per 100 gallons to improve weed control performance. Before using other kinds of adjuvants, test on a small area to be sure the treatment is safe to the crop.

Apply LOROX® DF at 1.0 to 2.0 pounds per acre after parsley has a minimum of 3 true leaves or crop injury may result. More than one application can be made providing the user has prior experience to indicate adequate crop safety. Wait at least 21 days between applications. Do not treat plants that are under stress. Avoid late fall applications prior to cold weather as crop injury may occur. Do not make more than two applications in the first growing season and do not make more than two applications in the second growing season. When using on a new variety for the first time, treat a small area to determine crop safety prior to treating a larger area.

User must accept all crop injury liability if an adjuvant other than a non-ionic surfactant at 1 quart per 100 gallons is used or if more than one application per crop is made.

Note: All parsley seed screenings shall be disposed of in such a way that they cannot be distributed or used for human food or animal feed. The seed conditioner shall keep records of screening disposal for three years from the date of disposal and shall furnish the records to the state department of agriculture immediately upon request. Disposal records shall consist of documentation from a controlled dumpsite, incinerator, or other equivalent disposal site and shall show the lot numbers, amount of material disposed of, its grower(s), and date of disposal.

No portion of the parsley seed plant, including but not limited to green chop, hay, pellets, meal, whole seed, cracked seed, roots, bulbs, leaves and seed screenings may be used or distributed for food or feed purposes.

Parsley seed shall bear a tag or container label which forbids use of the seed for human consumption or animal feed.

Parsley seed may not be distributed for human consumption or animal feed.

IMPORTANT

BEFORE USING THIS PRODUCT HERBICIDE, READ AND CAREFULLY NOTE THE CAUTIONARY STATEMENTS AND OTHER PROCEDURAL INFORMATION APPEARING ON THE EPA REGISTERED LABEL OR ON OTHER SUPPLEMENTAL LABELS.

This bulletin contains new or supplemental instructions for use of these products in combination which does not appear on the package label. Follow the instructions carefully.

This labeling must be in the possession of the user at the time of pesticide application.

Read the Limitation of Warranty and Liability on the Section 3 Federal product label before buying or using THIS product. If terms are not acceptable, return the unopened package at once to Seller for full refund of purchase price paid. Otherwise, use by Buyer or any other User constitutes acceptance of the terms of the Limitation of Warranty and Liability on the Section 3 Federal product label.

DR-680 012907-2