

## **PLEASE NOTE**

**This image contains more than one label approved for this product on this date.**

352-722

**ACCEPTED**  
AUG 30 2007  
Under the Federal Insecticide,  
Fungicide, and Rodenticide Act,  
as amended for the pesticides  
registered under  
EPA Reg. No. 352-722

08/30/2007

1/2

**SUPPLEMENTAL  
LABELING**

**DuPont Crop  
Protection**

**DUPONT™ BREAKFREE™  
HERBICIDE ROTATIONAL CROP  
RESTRICTIONS AFTER  
FINAL APPLICATION**

**RESTRICTED USE PESTICIDE**  
**Due to Oncogenicity**  
For retail sale to and use only by Certified Applicators, or persons under their direct supervision, and only for those uses covered by the Certified Applicator's certification.

**DUPONT™ BREAKFREE™ HERBICIDE**

EPA Reg. No. 352-722

**ROTATIONAL CROP RESTRICTIONS AFTER FINAL APPLICATION**

**ATTENTION**

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This label must be in the possession of the user at the time of pesticide application.
- Read the label affixed to the container for BREAKFREE™ herbicide before applying. Follow all applicable use directions and precautions on the EPA-registered label.
- Use of BREAKFREE™ according to this supplemental labeling is subject to all restrictions and limitations imposed by the label affixed to the product container.

**DIRECTIONS FOR USE**

Refer to product label for General Use Precautions, Mixing and Application instructions.

**Rotational Crop Restrictions**


When tank mixing with other herbicides, follow the most restrictive crop rotation guidelines on the label of each product used. The following rotational crops may be planted as indicated:

| Crop   | Interval                             |
|--|--------------------------------------|
| Corn (1)   | Anytime - 0 Months After Application |
| Wheat  | 4 Months After Application           |
| Alfalfa, Barley, Buckwheat, Clover, Dry beans (2), Guar, Kudzu, Lentil, Lespedeza, Lupin (4), Millet, pearl or proso, Oats, Pea (5), Potatoes, Rye, Sorghum, Soybeans, Sugar beets, Sunflower, Trefoil, Tobacco, Triticale, Vetch, Wild rice | Spring Following Application (3)     |

Numbers within parentheses (-) in the table refer to Specific Rotational Crop Requirements below.

- (1) If crop treated with BREAKFREE™ is lost, corn may be replanted immediately. Do not make a second application of BREAKFREE™.
- (2) Dry beans includes: adzuki, kidney, lima, navy, pinto
- (3) Approved rotation crops list does not include any species of succulent beans and peas
- (4) Lupin includes: grain, white, white sweet
- (5) Pea includes: blackeyed, chick, cow, Crowder, field, pigeon, Southern

**Note:** These rotational crop restrictions supercede prior restrictions of 4 months for wheat and the spring following application for sorghum, soybeans, and tobacco.



For product information call 1-888-6-DUPONT


© 2007 E. I. du Pont de Nemours and Company, Crop Protection, Wilmington, Delaware 19898



**NEXT**

**LABEL**

352-722



08/30/2007

# SUPPLEMENTAL LABELING

1/2

## DuPont Crop Protection

## DUPONT™ BREAKFREE™ HERBICIDE WEED CONTROL IN SWEET CORN

### RESTRICTED USE PESTICIDE

#### Due to Oncogenicity

For retail sale to and use only by Certified Applicators, or persons under their direct supervision, and only for those uses covered by the Certified Applicator's certification.

## DUPONT™ BREAKFREE™ HERBICIDE

EPA Reg. No. 352-722

## WEED CONTROL IN SWEET CORN

### ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This label must be in the possession of the user at the time of pesticide application.
- Read the label affixed to the container for BREAKFREE™ herbicide before applying. Follow all applicable use directions and precautions on the EPA-registered label.
- Use of BREAKFREE™ according to this supplemental labeling is subject to all restrictions and limitations imposed by the label affixed to the product container.

### DIRECTIONS FOR USE

#### DuPont™ BREAKFREE™ Applied Alone

Used alone, BREAKFREE™ herbicide may be applied to sweet corn using rates recommended in the product label for BREAKFREE™ and application methods and timing listed below.

**Carriers and Spray Volume:** Either water or liquid fertilizers such as solutions, slurries or suspensions may be used as liquid carriers. Refer to product label for BREAKFREE™ for details.

**Application Timing and Methods:** For the optimum period of effective weed control during the time most critical to corn production, preplant applications of BREAKFREE™ should occur as close as possible to planting. Preemergence applications should occur as close as possible to planting, but prior to weed emergence. Refer to product label for BREAKFREE™ for details.

**Note:** Do not apply BREAKFREE™ to sweet corn as an early postemergence application.

- **Early Preplant:** On medium and fine textured soils (see Table 1 in main label), BREAKFREE™ may be applied up to 30 days prior to planting.
- **Preplant Incorporation:** BREAKFREE™ and certain tank mixes may be mechanically incorporated in the top 2 inches of the soil with field cultivators, discs, or spring tooth harrows at any time within 14 days prior to planting.
- **Preemergence Surface:** BREAKFREE™ and certain tank mixes may be applied to the soil surface as a broadcast or banded application.
- **Postplant-Preemergence:** BREAKFREE™ may be applied immediately after planting but prior to corn emergence.
- **Banding Preemergence:** BREAKFREE™ may be applied in a 10 to 14 inch band after corn planting but prior to corn emergence.

**Sprinkler Irrigation:** Do not apply BREAKFREE™ by sprinkler irrigation. Use a sprinkler system only to incorporate BREAKFREE™ after application. Refer to product label for BREAKFREE™ for details.

For product information call 1-888-6-DUPONT

© 2007 E. I. du Pont de Nemours and Company, Crop Protection, Wilmington, Delaware 19898

2/2

**Cultivation:** Cultivation should be delayed as long as possible. If weeds develop, a shallow cultivation or rotary hoeing will generally result in improved weed control. Refer to product label for BREAKFREE™ for details.

**Soil Texture and Organic Matter:** The use rate of BREAKFREE™ is determined by a combination of two factors, soil texture and organic matter, which must be determined prior to application. Refer to Table 1 in main product label for soil texture classifications.

**Use Rates for Conventional Tillage Systems:** Refer to table 2 in main label for application rates and detailed information for conventional systems.

**Use Rates for Reduced Tillage Systems:** Refer to table 3 in main label for application rates and detailed information for reduced tillage systems.

**Weeds Controlled:** Refer to table 4 in main label for details.

**BREAKFREE™ Applied in Tank Mix Combination**

Additional weeds may be controlled with tank mixes of BREAKFREE™ and other products labeled for use on sweet corn. Tank mix combinations may be used in conventional, reduced, or no-till systems and may be applied using application methods and timing listed above unless otherwise specified in the tank mix product label. Refer to the product label for BREAKFREE™ for details on carriers, mixing instructions and testing for compatibility of tank mixtures.

**IMPORTANT  
BEFORE USING BREAKFREE™, READ AND  
FOLLOW ALL APPLICABLE DIRECTIONS,  
RESTRICTIONS AND PRECAUTIONS ON THE  
EPA-REGISTERED LABEL.**

This bulletin contains new or supplemental instructions for use of this product which do not appear on the EPA-registered package label. Follow the instructions carefully.

This labeling must be in the possession of the user at the time of pesticide application.

Read the Limitation of Warranty and Liability on the Section 3 Federal product label before buying or using THIS product. If terms are not acceptable, return the unopened package at once to Seller for full refund of purchase price paid. Otherwise, use by Buyer or any other User constitutes acceptance of the terms of the Limitation of Warranty and Liability on the Section 3 Federal product label.

DR-753 072307-2

