



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY  
WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY  
AND POLLUTION PREVENTION

December 2, 2015

Tim McPherson  
Senior Product Registration Manager  
DuPont Crop Protection  
Stine-Haskell Research Center  
P.O. Box 30  
Newark, DE 19714

Subject: PRIA Label Amendment – The addition of California rice and increase single application rates (no increase to amount per year)  
Product Name: DuPont Coragen Insect Control  
EPA Registration Number: 352-729  
Application Date: 2/6/2015  
Decision Number: 499886

Dear Mr. McPherson:

The amended label referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act, as amended, is acceptable. This approval does not affect any conditions that were previously imposed on this registration. You continue to be subject to existing conditions on your registration and any deadlines connected with them.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. You must submit one copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 18 months from the date of this letter. After 18 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

Page 2 of 2  
EPA Reg. No. 352-729  
Decision No. 499886

Your release for shipment of the product constitutes acceptance of these conditions. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6. If you have any questions, please contact Jacquelyn Marchese by phone at 703-347-0559, or via email at [marchese.jacquelyn@epa.gov](mailto:marchese.jacquelyn@epa.gov).

Sincerely,

A handwritten signature in black ink, appearing to read 'VE', followed by a horizontal line extending to the right.

Venus Eagle, Product Manager 01  
Invertebrate and Vertebrate Branch 3  
Registration Division (7505P)

Enclosure



# DuPont™ Coragen®

insect control  
with the active ingredient RynaXypyr®

GROUP	28	INSECTICIDE
-------	----	-------------

CORAGEN® is a suspension concentrate.

Contains 1.67 lb. active ingredient per gallon.

<i>Active Ingredient</i>	<i>By Weight</i>
Chlorantraniliprole	
3-Bromo-N-[4-chloro-2-methyl-6-[(methylamino)carbonyl]phenyl]-1-(3-chloro-2-pyridinyl)-1H-pyrazole-5-carboxamide	18.4%
<i>Other Ingredients</i>	81.6%
TOTAL	100.0%

EPA Reg. No. 352-729

**Nonrefillable Container**


Net: \_\_\_\_\_

OR

**Refillable Container**

Net: \_\_\_\_\_

E. I. du Pont de Nemours and Company  
1007 Market Street  
Wilmington, Delaware 19898  
Phone: 1-800-441-7515 (Toll Free)



EPA Est. No. \_\_\_\_\_

Not for sale, sale into, distribution and/or use in Nassau, Suffolk, Kings, and Queens counties of New York State.

## PRECAUTIONARY STATEMENTS

### KEEP OUT OF REACH OF CHILDREN

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

### FIRST AID

For questions regarding emergency medical treatment, you may contact 1-800-441-3637 for information.

### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

When used as directed this product does not present a hazard to humans or domestic animals.

### PERSONAL PROTECTIVE EQUIPMENT

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants.
- Shoes plus socks.

After the product has been diluted in accordance with label directions for use, shirt, pants, socks, and shoes are sufficient Personal Protective Equipment. Follow manufacturer's instructions for cleaning/maintaining personal protective equipment (PPE). If no such instructions for washables are available, use detergent and hot water. Keep and wash PPE separately from other laundry.

### USER SAFETY RECOMMENDATIONS

**USERS SHOULD:** Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

## ENVIRONMENTAL HAZARDS

This pesticide is toxic to aquatic invertebrates, oysters, and shrimp. Do not apply directly to water. Drift and runoff may be hazardous to aquatic organisms in water adjacent to use sites.

### Surface Water Advisory-

This product may impact surface water quality due to runoff of rain water. This is especially true for poorly draining soils and soils with shallow ground water. This product is classified as having high potential for reaching surface water via runoff for several months or more after application. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of chlorantraniliprole from runoff water and sediment. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 48 hours.

### Ground Water Advisory-

This chemical has properties and characteristics associated with chemicals detected in ground water. This chemical may leach into ground water if used in areas where soils are permeable, particularly where the water table is shallow.

## DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

DuPont™ CORAGEN® insect control must be used only in accordance with the directions on this label, in separate EPA-approved labeling or exemptions under FIFRA (Supplemental Labels, Special Local Need Registrations, FIFRA Section 18 exemptions, FIFRA 2(ee) Bulletins), or as otherwise permitted by FIFRA. Always read the entire label, including the Limitation of Warranty and Liability.

## RESTRICTIONS

- Do not treat plants grown for transplanting. Not for use in nurseries, plant propagation houses, or greenhouses by commercial transplant producers on plants being grown for transplanting.
- This product is only for commercial use.
- Not for use on ornamental plants or plants being grown for ornamental purposes.
- May be used on crops on this label grown for seed production.
- Not for residential use.
- Do not apply CORAGEN® through any irrigation system unless specified in the crop section of this label or in EPA approved supplemental labeling.

### For New York State Only:

The following restrictions are required to permit use of CORAGEN® insect control in the State of New York:

- This product may not be applied within 100 feet of a water body (lake, pond, river, stream, wetland, or drainage ditch).
- Aerial application of this product is prohibited.
- Not for sale, sale into, distribution and/or use in Nassau, Suffolk, Kings, and Queens counties of New York State.

## AGRICULTURAL USE REQUIREMENTS

CORAGEN® insect control must be used only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on the label about personal protective equipment, restricted-entry interval, and notification to workers (as applicable).

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

For any requirements specific to your State or Tribe, consult the State or Tribal agency responsible for pesticide regulation.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 4 hours.

For early entry into treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, wear:

- Long-sleeved shirt and long pants
- Shoes plus socks

CORAGEN® is a suspension concentrate that can be applied as: an in-furrow spray at planting, transplant water treatment, hill drench at planting, surface band at planting, soil shank injection at planting, drip chemigation, or foliar spray (including overhead sprinkler chemigation on certain crops as specified on this label) to control listed insects. Not all application methods

are allowed on all crops; see specific crop sections of this label for which application methods may be used. DuPont™ CORAGEN® is mixed with water for application. CORAGEN® may be used on crops on this label grown for seed production.

CORAGEN® is a member of the anthranilic diamide class of insecticides with a novel mode of action acting on insect ryanodine receptors. Although CORAGEN® has contact activity, it is most effective through ingestion of treated plant material. After exposure to CORAGEN®, affected insects will rapidly stop feeding, become paralyzed, and typically die within 1 - 3 days. Time applications to the most susceptible insect pest stage, typically at egg lay, egg hatch, and/or at egg hatch and/or newly hatched larvae, before populations reach damaging levels. When pest populations are high, use the highest listed application rate for that pest.

### **INTEGRATED PEST MANAGEMENT**

DuPont supports the use of Integrated Pest Management (IPM) programs to control pests. This product may be used as part of an IPM program, which can include biological, cultural, and genetic practices, aimed at preventing economic pest damage. IPM principles and practices include field scouting or other detection methods, correct target pest identification, population monitoring, rotation of insecticides with different modes-of-action, and treating when target pest populations reach locally determined action thresholds. Consult your state cooperative extension service, professional consultants or other qualified authorities to determine appropriate action treatment threshold levels for treating specific pest/crop or site systems in your area.

### **SCOUTING**

Monitor insect populations to determine whether or not there is a need for application of CORAGEN® based on locally determined economic thresholds and pest management guidelines.. More than one treatment of CORAGEN® may be required to control a population of pests.

### **INSECT RESISTANCE MANAGEMENT**

For resistance management, CORAGEN® is a Group 28 Insecticide. Repeated and exclusive use of CORAGEN® (chlorantraniliprole, belonging to the anthranilic diamide class of chemistry), or other Group 28 Insecticide may lead to the buildup of resistant strains of insects in some crops.

Some insects are known to develop resistance to products used repeatedly for control. Because the development of resistance cannot be predicted, this product may be used as part of resistance management strategies established for the use area. These strategies may include incorporation of cultural and biological control practices, alternation of mode-of-action classes of insecticides on succeeding generations and targeting the most susceptible life stage. Consult your local or state agricultural authorities for details.

Unless directed otherwise in the specific crop/pest sections of this label, the best practices are to follow these instructions to delay the development of insecticide resistance:

- Avoid using the same mode of action (same IRAC group number) on consecutive generations of insect pests.
- Apply CORAGEN® or other Group 28 insecticides using a “treatment window” approach to avoid exposure of successive insect pest generations to the same mode of action.
- A “treatment window” is defined as the period of residual activity provided by single or sequential applications of products with the same mode of action. This “treatment window” should not exceed approximately the length of one generation of the target pest.
- Within the “Group 28 treatment window”, make no more than 3 applications of CORAGEN® or other Group 28 insecticides within a single generation of the target pest on a crop.
- Following a “Group 28 treatment window”, rotate to a treatment window of effective products with a different mode of action. This “Non-Group 28 Window” should approximate the duration of one generation of the target pest.
- Avoid using less than the labeled rates of CORAGEN® when applied alone or in tank mixtures.
- Target the most susceptible insect life stages, whenever possible.
- Monitor insect populations for product effectiveness.

If resistance to CORAGEN® develops in your area, CORAGEN® or other products with a similar mode of action, may not provide adequate control. If poor performance cannot be attributed to improper application or extreme weather conditions, a resistant strain of insect may be present. If you experience difficulty with control and resistance is a reasonable cause, immediately consult your local company representative or agricultural advisor for the best alternate method of control for your area. For additional information on insect resistance monitoring, visit the Insecticide Resistance Action Committee (IRAC) on the web at <http://www.irac-online.org>.

### **APPLICATION**

Apply at the specified rates when insect populations reach locally determined economic action thresholds. Consult the cooperative extension service, professional consultants or other qualified authorities to determine appropriate threshold levels for treatment in your area.

Apply follow-up treatments of DuPont™ CORAGEN®, as specified, to keep pest populations within threshold limits. Refer to the Resistance Management section of this label for further guidance on follow-up treatments. See individual crop sections of this label for specific minimum spray intervals.

Use sufficient water to obtain thorough, uniform coverage.

CORAGEN® may be applied by: ground (including an in-furrow spray at planting, transplant water treatment, hill drench at planting, surface band at planting, soil shank injection at planting, drip chemigation, or foliar), or aerial application equipment. Not all application methods are allowed on all crops; see specific crop sections of this label for which application methods may be used. CORAGEN® may be applied via overhead sprinkler chemigation systems on some crops; see specific crop sections of this label for crops where overhead sprinkler chemigation may be used. For aerial application use the following directions unless otherwise specified in specific crop/pest sections of this label or EPA-approved supplemental labeling, use a minimum of 5 gallons per acre (gpa) of water. The highest labeled rate for a specified pest may be necessary when aerial applications are made. For all other application methods use the following directions, unless otherwise specified in specific crop/pest sections of this label or EPA-approved supplemental labeling, use a minimum of 10 gal per acre (GPA) of water for all crops.

**Use of Adjuvants** - In some situations where coverage is difficult to achieve such as closed canopy, dense foliage, plants with waxy leaf surfaces, or less than optimum application equipment, an adjuvant may improve performance. Use only adjuvant products that are labeled for agricultural use and follow the directions on the manufacturer's label. Always conduct a premix test for compatibility. Use an adjuvant that does not affect foliage and/or fruit finish. Refer to specific crop sections of this label for additional adjuvant guidance.

## **SPRAY PREPARATION**

Spray equipment must be clean and free of previous pesticide deposits before applying CORAGEN®. Fill spray tank 1/4 to 1/2 full of water. Add CORAGEN® directly to spray tank. Mix thoroughly to fully disperse the insecticide, once dispersed continued agitation is required. Use mechanical or hydraulic means; do not use air agitation. Do not store spray mix solutions overnight in spray tank. Observe the most restrictive of the labeling limitations and precautions of all products used in mixtures.

## **TANK MIXTURES**

This product can be mixed with pesticide products that are labeled for use on the same crops as CORAGEN®. Do not exceed labeled dosage rates. This product cannot be mixed with any product containing a label prohibition against such mixing.

CORAGEN® may be mixed with certain liquid fertilizers for at-plant soil applications. Do not mix CORAGEN® directly with pure liquid fertilizers.

It is the pesticide user's responsibility to ensure that all products in a tank mix are registered for the intended use. Users must follow the most restrictive directions and precautionary language of the products of the mixture (for example, first aid from one product, spray drift management from another).

Before using a tank mix for the first time, always determine the compatibility of CORAGEN® with the tank mixtures by using a jar test.

**Compatibility** -Since formulations may be changed and new ones introduced, premix a small quantity of a desired tank mix and observe for possible adverse changes (settling out, flocculation, etc.).

Steps to conduct a jar test to determine physical tank mix compatibility of CORAGEN® with other products:

- Add clean water to jar proportional to the planned water volume that will be used in the spray tank (a jar size of 8-16 oz is acceptable).
- Using the most restrictive PPE of the products to be tested, mix proper proportions of CORAGEN® and desired tank mix partner(s) as will be present in the spray tank, add one product at a time following the sequence of addition according to formulation type provided in this label.
- Seal and shake mixture after each product is added.
- Allow to stand for 1 hour.
- View jar to determine if settling, flocculation, crystallization or any other undesirable changes have happened.
- If none of the above is observed or the solution can be easily remixed after shaking, the mixture is compatible with CORAGEN®.
- If the tank mix is not compatible, a higher water volume, reduced rate of the tank mix partner(s), reduced number of tank mix partners or a compatibility agent may be needed.

**Tank Mixtures and Crop Safety** - Crop varieties can differ in their responsiveness to tank mixtures, and environmental conditions can have an influence on product performance and crop response. It is not possible to test CORAGEN® alone or with all possible tank mix combinations on all varieties under all environmental conditions. When considering the use of a tank mixture on a labeled crop without prior experience, or which is not specifically described on CORAGEN® product labeling or in other DuPont product use instruction, it is important to check crop safety first. To test for crop safety prepare

a small volume of the intended tank mixture, apply it to an area of the target crop as directed by both this and the tank mix partner product labels, and observe the treated crop to ensure that a phytotoxic response does not occur.

Use of CORAGEN® in any tank mixture applications that is not specifically described on CORAGEN® product labeling or in other DuPont product use instructions, could potentially result in crop injury. Follow the precautions on this label and on the label for any other product to be used in tank mixtures before making such applications to your crops. DuPont will not be responsible for any crop injury arising from the use of a tank mixture that is not specifically described on DuPont™ CORAGEN® product labeling or in other DuPont product use instruction.

**Tank Mixing Sequence** -Fill spray tank 1/4 to 1/2 full of water. While agitating, add the different formulation types in the sequence indicated below\*. Allow time for complete mixing and dispersion after addition of each product before adding the next product.

1. Water soluble bag (WSB)
2. Water soluble granules (SG)
3. Water dispersible granules (WG, XP, DF)
4. Wettable powders (WP)
5. CORAGEN® and other water based suspension concentrates (SC)
6. Water soluble concentrates (SL)
7. Suspoemulsions (SE)
8. Oil based suspension concentrates (OD)
9. Emulsifiable concentrates (EC)
10. Surfactants, oils adjuvants
11. Soluble fertilizers
12. Drift retardants

\* Unless otherwise specified by manufacturer directions for use or by local experience.

## **SPRAY TANK CLEANOUT**

Prior to application, start with clean, well maintained application equipment. Immediately following application, thoroughly clean all spray equipment to reduce the risk of forming hardened deposits which might become difficult to remove.

Drain spray equipment. Thoroughly rinse sprayer and flush hoses, boom and nozzles with clean water. Clean all other associated application equipment. Take all necessary safety precautions when cleaning equipment. Do not clean near wells, water sources or desirable vegetation. Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

## **SPRAY DRIFT MANAGEMENT**

The interaction of many equipment and weather-related factors determines the potential for spray drift. The applicator is responsible for considering all these factors when making application decisions.

**AVOIDING SPRAY DRIFT IS THE RESPONSIBILITY OF THE APPLICATOR.**

### **IMPORTANCE OF DROPLET SIZE**

The most effective drift management strategy is to apply the largest droplets which are consistent with pest control objectives. The presence of sensitive species nearby, the environmental conditions, and pest pressure may affect how an applicator balances drift control and coverage. Applying larger droplets reduces drift potential, but will not prevent drift if applications are made improperly or under unfavorable environmental conditions. A droplet size classification system describes the range of droplet sizes produced by spray nozzles. The American Society of Agricultural and Biological Engineers (ASABE) provide a Standard that describes droplet size spectrum categories defined by a number of reference nozzles (fine, coarse, etc.). Droplet spectra resulting from the use of a specific nozzle may also be described in terms of volume mean diameter (VMD). Coarser droplet size spectra have larger VMD's and lower drift potential.

### **CONTROLLING DROPLET SIZE - GROUND APPLICATION**

**Nozzle Type** - Select a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. The use of low-drift nozzles will reduce drift potential.

**Pressure** - The lowest spray pressures recommended for the nozzle produce the largest droplets. Higher pressure reduces droplet size and does not improve canopy penetration. When higher flow rates are needed, using a higher-capacity nozzle instead of increasing pressure results in the coarsest droplet spectrum.

**Flow Rate/Orifice Size** - Using the highest flow rate nozzles (largest orifice) that are consistent with pest control objectives reduces the potential for spray drift. Nozzles with higher rated flows produce coarser droplet spectra.

## **CONTROLLING DROPLET SIZE - AIRCRAFT**

**Number of Nozzles** -Using the minimum number of nozzles with the highest flow rate that provide uniform coverage will produce a coarser droplet spectrum.

**Nozzle Orientation** -Orienting nozzles in a manner that minimizes the effects of air shear will produce the coarsest droplet spectra. For some nozzles such as solid stream, pointing the nozzles straight back parallel to the airstream will produce a coarser droplet spectrum than other orientations.

**Nozzle Type** -Solid stream, or other low drift nozzles produce the coarsest droplet spectra.

Do not apply as a ULV application.

## **BOOM LENGTH AND HEIGHT**

**Boom Length (aircraft)** -The boom length must not exceed 3/4 of the wing length; using shorter booms decreases drift potential. For helicopters use a boom length and position that prevents droplets from entering the rotor vortices.

**Boom Height (aircraft)** -Application more than 10 ft above the canopy increases the potential for spray drift. Applications made at the lowest height consistent with pest control objectives, and the safe operation of the aircraft will reduce the potential for spray drift.

**Boom Height (ground)** -Applications made at the lowest height consistent with pest control objectives, and that allow the applicator to keep the boom level with the application site and minimize bounce, will reduce the exposure of spray droplets to evaporation and wind and reduce spray drift potential.

## **WIND**

Drift potential increases at wind speeds of less than 3 mph (due to variable direction and inversion potential) or more than 10 mph. However, many factors, including droplet size and equipment type determine drift potential at any given wind speed. AVOID APPLICATIONS DURING GUSTY OR WINDLESS CONDITIONS. Do not make applications when wind speeds are greater than 15 mph.

Note: Local terrain can influence wind patterns. Every applicator must be familiar with local wind patterns and how they affect spray drift.

## **TEMPERATURE AND HUMIDITY**

When making applications in hot and dry conditions, set up equipment to produce larger droplets to reduce effects of evaporation.

## **SURFACE TEMPERATURE INVERSIONS**

Do not make applications into temperature inversions. Drift potential is high during a surface temperature inversion. Surface inversions restrict vertical air mixing, which causes small-suspended droplets to remain close to the ground and move laterally in a concentrated cloud. Surface inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates a surface inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

## **SHIELDED SPRAYERS**

Shielding the boom or individual nozzles can reduce the effects of wind. However, it is the responsibility of the applicator to verify that the shields are preventing drift and not interfering with uniform deposition of the product.

## **AIR ASSISTED (AIRBLAST) FIELD CROP SPRAYERS**

Air assisted field crop sprayers carry droplets to the target via a downward directed air stream. Some may reduce the potential for drift, but if a sprayer is unsuitable for the application and/or set up improperly, high drift potential can result. It is the responsibility of the applicator to determine that a sprayer is suitable for the intended application, is configured properly, and that drift is not occurring.

Note: Air assisted field sprayers can affect product performance by affecting spray coverage and canopy penetration. Consult the application equipment section of this label to determine if use of an air assisted sprayer is recommended.

## **CHEMIGATION**

The following types of irrigation equipment may be used for chemigation applications: drip (trickle), or strip tubing irrigation systems. DuPont™ CORAGEN® may be applied through overhead sprinkler irrigation systems (see CHEMIGATION USING OVERHEAD SPRINKLER SYSTEMS - CEREAL GRAINS, CORN (FIELD, POP, SEED), COTTON, GRASS FORAGE, FODDER, and HAY, LEGUMES, MINT (PEPPERMINT AND SPEARMINT), NON-GRASS ANIMAL FEEDS, OILSEED GROUP, PEANUT, POTATO, SOYBEAN AND SUGARCANE section of this label).

Apply CORAGEN® in sufficient water and of sufficient duration to ensure the recommended rate is applied evenly to the entire treated area. Do not allow irrigation water to collect or runoff during chemigation; do not allow pooling of irrigation water. Inject CORAGEN® downstream from any water filtration system.



CORAGEN® must not be applied at the same time that a drip/irrigation line clean out product is being used as performance may be reduced. Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water.

If you have questions about calibration, you should contact state extension service specialists, equipment manufacturers, or other experts. A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

Wear personal protective equipment as defined in the PPE section of the label for applicators and other handlers when making adjustments or repairs on the chemigation system when CORAGEN® is in the irrigation water. When the application is finished, allow the entire irrigation and injector system to be thoroughly flushed clean before stopping the system. A pesticide supply tank is recommended for the application of DuPont™ CORAGEN® in chemigation systems.

Do not connect any irrigation system used for pesticide applications to a public water system unless the pesticide label-prescribed safety devices are in place. See **“Required System Safety Devices for All Chemigation Systems” at the end of the Chemigation section.** Public water system means a system for the provision to the public of piped water for human consumption, if such system has at least 15 service connections or regularly serves an average of at least 25 individuals at least 60 days out of the year.

## **APPLICATION INSTRUCTIONS**

### ***DRIP (TRICKLE) CHEMIGATION***

CORAGEN® must be applied in a manner that ensures the product is in the root zone. CORAGEN® must be in the root zone to provide effective control of target pests. CORAGEN® is most effective when it is applied so that the roots are at or near the site of application; manage irrigation so that significant quantities of CORAGEN® remain in the root zone where it is most effective. Unless directed otherwise in the specific crop sections of this label, a total of two applications can be made per crop season. Any subsequent CORAGEN® treatments must be foliar applications.

1. Do not begin applications until after crop emergence in direct seeded crops.
2. Do not make applications if soil moisture is below the level required for active plant growth.
3. This product must be applied uniformly in the root zone or poor performance will result. Drip tape or emitters must be located within or directly adjacent to the root zone.
4. The drip system must be properly designed, free of leaks, and operated in manner that provides uniform application of water throughout the field.
5. In most situations, this product should be applied during the first 1/3 of the irrigation cycle, starting just after the system has come up to pressure.
6. The minimum injection period is the time that it takes water to move from the injection point to the furthest emitter in the irrigation zone (propagation time). If this time is not known, it can be calculated by measuring the time for a soluble dye to move from the injection point to the furthest emitter. A longer injection improves uniformity throughout the zone, but needs to allow for at least an equal period of water to flush the system and move the product through the soil.

### ***CHEMIGATION USING OVERHEAD SPRINKLER SYSTEMS - CEREAL GRAINS, CORN (FIELD, POP, SEED) COTTON GRASS FORAGE, FODDER, AND HAY, LEGUMES, MINT (PEPPERMINT AND SPEARMINT), NON-GRASS ANIMAL FEEDS, OILSEED GROUP, PEANUT, POTATO, SOYBEAN, SUGARCANE***

Types of Chemigation Systems: CORAGEN® may be applied to CEREAL GRAINS, CORN (FIELD, POP, SEED) COTTON GRASS FORAGE, FODDER, and HAY, LEGUMES, MINT (PEPPERMINT AND SPEARMINT), NON-GRASS ANIMAL FEEDS, OILSEED GROUP, PEANUT, POTATO, SOYBEAN, SUGARCANE through overhead sprinkler irrigation systems, including the following; center pivot, end tow, hand move, lateral move, side roll, solid set and wheel line. The irrigation system used must provide uniform water distribution.

#### **Directions for Chemigation:**

##### **Preparation**

A pesticide tank is recommended for the application of CORAGEN® in chemigation systems.

Thoroughly clean the injection system and tank of any fertilizer or chemical residues using a standard clean-out procedure. Dispose of any residues in accordance with State and Federal laws. With the mix tank 1/4 to 1/2 full with water and the agitator running, measure the required amount of CORAGEN® and add it to the tank. Then add additional water to bring your total pesticide mixture up to the desired volume for your application. Note: Always add the CORAGEN® to water, never put CORAGEN® into a dry tank or other mixing equipment without first adding water. See "Tank Mixing Sequence" section of the container label for tank mixing sequence. Continue to agitate the mixture throughout the application process. Use mechanical or hydraulic agitation, do not use air agitation.

##### **Injection Into Chemigation Systems**

Inject the proper amount of CORAGEN® into the irrigation water flow using a positive displacement injection pump or a Venturi injector. Injection should occur at a point in the main irrigation water flow to ensure thorough mixing with the irrigation water. For continuously moving systems, inject the solution containing CORAGEN® into the irrigation water line continually and uniformly throughout the irrigation cycle.

Apply in no more than 0.2 inches of water per acre. For overhead sprinkler systems that are stationary, add the solution containing CORAGEN® to the irrigation water line and apply no more than 0.2 inches of water per acre.

### **Uniform Water Distribution**

The irrigation system used for application of CORAGEN® must provide for uniform distribution of CORAGEN® treated water. Non-uniform distribution can result in crop injury, lack of effectiveness or illegal pesticide residues in or on the crop being treated. Ensure the irrigation system is calibrated to uniformly distribute the chemigation application to the crop. Contact the equipment manufacturer, the local University Extension agent or other experts if you have questions about achieving uniform distribution of the application.

### **Equipment Calibration**

Calibrate the irrigation system and injector before applying DuPont™ CORAGEN®. Calibrate the injection pump while the system is running using the expected irrigation rate. If you have questions about calibration, you should contact your state extension service specialists, equipment manufacturer or other experts.

### **Monitoring of Chemigation Applications**

A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of a responsible person, shall shut the system down and make necessary adjustments should the need arise. Wear the personal protective equipment as defined in the PPE section of the label for applicators and other handlers when making adjustments or repairs on the chemigation system when CORAGEN® is in the irrigation water.

### **Operation**

Start the water pump and sprinkler, and let the system achieve the desired pressure and speed before starting the injector. Start the injector and calibrate the injection system according to the directions above. This procedure is necessary to deliver the desired rate per acre in a uniform manner. When the application is finished, allow the entire irrigation and injector system to be thoroughly flushed clean before stopping the system.

- End guns must be turned off during the application, if they irrigate nontarget areas or if they do not provide uniform application and coverage.
- It is recommended that nozzles in the immediate area of wells, control panels, chemical supply tanks and system safety devices be plugged to prevent contamination of these areas.
- Do not apply when wind speed favors drift beyond the area intended for treatment.
- Do not apply when system connections or fittings leak or when nozzles do not provide uniform distribution.
- Do not allow irrigation water to collect or run-off during chemigation.

### **Cleaning the System**

Thoroughly clean the injection system and tank of any fertilizer or chemical residues using a standard clean-out procedure. Dispose of any residues in accordance with State and Federal laws. Consult your owner's manual or your local equipment dealer for cleanout procedures for your injection system.

### **REQUIRED SYSTEM SAFETY DEVICES FOR ALL CHEMIGATION SYSTEMS**

1. The system must contain a functional check valve, vacuum relief valve and low-pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.
2. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
3. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
4. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
5. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
6. Systems must use a metering device, such as a positive displacement pump or a Venturi injector, that provides uniform injection of the product, is effectively designed and constructed of materials compatible with the product, and is capable of being fitted with a system interlock.
7. Chemigation systems connected to public water systems must contain a functional, reduced- pressure zone, backflow preventer (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.

### **SOIL APPLICATIONS**

CORAGEN® must be applied in a manner that ensures the product is in the root zone. CORAGEN® must be in the root zone to provide effective control of target pests. CORAGEN® is most effective when it is applied so that the roots are at or

near the site of application; manage irrigation so that significant quantities of CORAGEN® remain in the root zone where it is most effective. Maintaining soil moisture to field capacity or to meet crop needs and environmental conditions aids in product availability to the roots and can improve efficacy. Applications of CORAGEN® to the root zone allow the active ingredient to be transported from the roots through the xylem providing upward systemicity. CORAGEN® is translocated to the canopy beginning immediately after the application, reaching an effective concentration in 1 to 3 days for seedlings and up to 7 days for larger plants. As the plant grows, the roots continue to absorb the available DuPont™ CORAGEN® from the reservoir in the soil providing extended protection of the plant canopy including new growth.

The length of control provided following soil applications will depend on the rate used, the pest being controlled and the environmental conditions; such as soil type, soil moisture, soil pH, etc. Use the higher specified rate within the rate range when pests are expected to occur later in the crop growth cycle or when pests are expected to be present continuously. CORAGEN® will primarily have activity in the foliage of treated plants and will not provide protection within the blooms and fruit. Foliar applications of other products may be needed to protect these parts of the plant. Unless directed otherwise in the specific crop sections of this label, only one soil application of CORAGEN® may be made per crop season, except for drip chemigation where a total of two applications can be made per season. If two drip applications are made then the application rate must not exceed 5 fl oz product (0.066 lb ai/acre) per application.

If CORAGEN® is applied as an at plant soil application, only one subsequent drip chemigation application can be made.

#### **In-Furrow Spray at Planting**

Apply as a narrow band spray into the furrow at the seeding depth.

#### **Transplant water treatment or Hill Drench**

Transplants should be adequately watered before transplanting. Apply at transplanting in a minimum of 2 fluid ounces of treatment solution per transplant. Ensure water volume is sufficient to thoroughly wet the root zone.

#### **Surface Band at Planting**

Apply as a narrow (2 inches or less) surface band spray above the seed line at planting. Incorporate surface band application within 24 hours of application using sufficient irrigation (usually 0.5 – 1.0 inches of water) to reach the seeding depth.

#### **Soil Shank Injection**

Use soil shank injection at planting. Applications must be incorporated using sufficient irrigation (usually 0.5 – 1.0 inches of water) to reach the root zone. Shank injection should be placed in the seed row or just below the seed line, within 1 - 2 inches of the seed line.

For insecticide resistance management it is important to avoid consecutive applications of insecticides with the same mode of action on successive generations of the same pest. See crops on label for recommended treatment rates and additional use information.

### **CROP ROTATION**

Crops on this label and the following crops or crop groups may be planted immediately following harvest: Artichoke, globe; Asparagus; Banana/Plantain; Brassica (Cole) Leafy Vegetables (Crop Group 5); Bulb Vegetables (Crop Group 3-07); Bushberry subgroup (Crop subgroup 13-07B); Cacao; Caneberry subgroup (Berry and Small Fruit Crop Group subgroup 13-07A); Cereal Grains (Crop Group 15); Forage, Fodder, and Straw of Cereal Grains (Crop Group 16); Citrus (Crop Group 10-10); Coffee; Corn (field, pop, seed, and sweet); Cotton; Cucurbit Vegetables (Crop Group 9); Figs; Fruiting Vegetables (Crop Group 8-10); Grass Forage, Fodder, and Hay Group (Crop Group 17); Herbs subgroup (Crop Group subgroup 19A); Grape; Hops; Large Shrub/Tree Berry subgroup (Crop subgroup 13-07C); Leafy Vegetables (nonbrassica, Crop Group 4); Legume Vegetables (Crop Group 6); Foliage of Legume Vegetables (Crop Group 7); Low Growing Berry subgroup (Crop subgroup 13-07G); Nongrass Animal Feeds (Forage, Fodder, Straw, and Hay Crop Group 18); Okra; Oilseed Group (Crop Group 20); Olives; Peanut; Persimmons; Pome Fruits (Crop Group 11-10); Pineapple; Pomegranates; Prickly Pear Cactus; Rice; Root and Tuber Vegetables (Crop Group 1); Leaves of Root and Tuber Vegetables (Crop Group 2); Small Fruit Vine Climbing subgroup, except fuzzy kiwifruit (Berry and Small Fruit Crop Group subgroup 13-07F); Soybean; Spice subgroup (Crop Group subgroup 19B); Spearmint and Peppermint; Stone Fruits (Crop Group 12-12); Sugarcane; Tea; Tree Nuts and Pistachio (Crop Group 14); Tobacco; and Tropical Fruits (acerola, atemoya, avocado, biriba, black sapote, canistel, cherimoya, custard apple, ilama, feijoa, guava, jaboticaba, longan, lychee, mamey sapote, mango, papaya, passionfruit, pulasan, rambutan, sapodilla, soursop, Spanish lime, star apple, starfruit, sugar apple, wax jambu, and White sapote (Casimiroa), and and/or hybrids of these).

All other crops cannot be planted until 12 months after the last application of CORAGEN®.

**DUPONT™ CORAGEN® RATE**

Crop	Application Method	Target Pest	Lb ai per acre	fluid ounces product per acre	Last Application (Days to Harvest)	REI (Hours)
Artichoke, globe	FOLIAR	Artichoke plume moth	0.045 - 0.098	3.5 - 7.5	3	4

**USE RESTRICTIONS**

Make no more than 4 applications per acre per crop.  
 Do not apply more often than each 14 days. Apply no more than 15.4 fl oz CORAGEN® or 0.2 lb a.i. of chlorantraniliprole containing products per acre per calendar year. Make applications between bud formation and harvest of an individual fruit.  
 Apply in a minimum of 10 gallons water per acre by air and 50 - 200 gallons of water per acre by ground (use sufficient water to obtain thorough coverage without excessive runoff).

Crop	Application Method	Target Pest	Lb ai per acre	fluid ounces product per acre	Last Application (Days to Harvest)	REI (Hours)
Asparagus	FOLIAR	Beet armyworm Western yellowstriped armyworm	0.045 - 0.098	3.5 - 7.5	1	4

**USE RESTRICTIONS**

Make no more than 4 applications per acre per crop season.  
 Minimum interval between treatments is 3 days.  
 Do not apply more than 15.4 fl oz CORAGEN® or 0.2 lb a.i. of chlorantraniliprole containing products per acre per calendar year.

Crop	Application Method	Target Pest	Lb ai per acre	fluid ounces product per acre	Last Application (Days to Harvest)	REI (Hours)
Onion bulbs, and onion green subgroups: (EPA Crop Group 3-07A and 3-07B) including Chive, fresh leaves; chive, Chinese, fesh leaves; Daylily, bulb; elegans hosta; Fritillaria, bulb; fritillaria, leaves; Garlic, bulb; Garlic, great-headed, bulb; Garlic, serpent, bulb; kurrat; lady's leek; leek; leek, wild; Lily, bulb; onion, Beltsville bunching; Onion, bulb; Onion, chinese, bulb; onion, fresh; onion, green; onion, macrostem; Onion, pearl; Onion, potato, bulb; onion, tree, tops; onion, Welsh, tops; Shallot, bulb; shallot, fresh leaves; cultivars, varieties, and/or hybrids of these	FOLIAR	Beet armyworm Western yellowstriped armyworm	0.045 - 0.098	3.5 - 7.5	1	4

**USE RESTRICTIONS**

Make no more than 4 applications per acre per crop.  
 Minimum interval between treatments is 7 days.  
 Do not apply more than 15.4 fl oz CORAGEN® or 0.2 lb a.i. of chlorantraniliprole containing products per acre per crop.  
 Do not apply more than 46.2 fl oz CORAGEN® or 0.6 lb a.i. of chlorantraniliprole containing products per acre per year; in NY do not apply more than 15.4 fl oz CORAGEN® or 0.2 lb a.i. of chlorantraniliprole containing products per acre per calendar year.

**DUPONT™ CORAGEN® RATE**

<b>Crop</b>	<b>Application Method</b>	<b>Target Pest</b>	<b>Lb ai per acre</b>	<b>fluid ounces product per acre</b>	<b>Last Application (Days to Harvest)</b>	<b>REI (Hours)</b>
Head and Stem Brassica and Leafy Brassica Greens (EPA Crop Subgroups 5 A and 5 B) including: Broccoli, Broccoli chinese (gai lon), Broccoli raab (rapini), Brussels sprouts, Cabbage, Chinese cabbage (bok choy), Chinese cabbage (napa), Cabbage, Chinese mustard (gai choy), Cauliflower, Caval broccolo, Collards, Kale, Kohlrabi, Mizuna, Mustard greens, Mustard spinach, Rape greens	SOIL AT PLANTING†(an in-furrow spray, transplant water treatment, hill drench, surface band, soil shank injection)	Beet armyworm Diamondback moth* Cabbage looper Cabbage maggot** Corn earworm Cross-striped cabbageworm Hawaiian beet webworm Imported cabbageworm Western Yellowstriped Armyworm	0.045 - 0.098	3.5 - 7.5 See rate conversion chart for rate per 1000 linear feet.	3	4
	DRIP CHEMIGATION†	Beet armyworm Diamondback moth* Cabbage looper Corn earworm Cross-striped cabbageworm Hawaiian beet webworm Imported cabbageworm Western Yellowstriped Armyworm	0.045 - 0.098	3.5 - 7.5		
	FOLIAR††	Silverleaf whiteflies (nymphs)*** Beet armyworm Cabbage looper Corn earworm Cross-striped cabbageworm Diamondback moth* Hawaiian beet webworm Imported cabbageworm Western Yellowstriped Armyworm	0.065 - 0.098	5.0 - 7.5		
			Beet armyworm Cabbage looper Corn earworm Cross-striped cabbageworm Diamondback moth* Hawaiian beet webworm Imported cabbageworm Western Yellowstriped Armyworm	0.045 - 0.098	3.5 - 7.5	

**USE RESTRICTIONS**

Make no more than 4 applications per acre per crop.

Minimum interval between treatments is 3 days for foliar applications and 10 days for drip chemigation applications.

Application via drip chemigation: drip tape must be placed directly underneath a single row to ensure CORAGEN® is applied in the root zone.

Do not apply more than 15.4 fl oz CORAGEN® or 0.2 lb a.i. of chlorantraniliprole containing products per acre per crop.

Do not apply more than 61.6 fl oz CORAGEN® or 0.8 lb a.i. of chlorantraniliprole containing products per acre per calendar year; in NY do not apply more than 15.4 fl oz CORAGEN® or 0.2 lb a.i. of chlorantraniliprole containing products per acre per calendar year.

†**SOIL APPLICATIONS** (an in-furrow spray at planting, transplant water treatment, hill drench at planting, surface band at planting, soil shank injection at planting, or drip chemigation): CORAGEN® must be applied uniformly in the root zone or poor performance will result. Surface band application requires sufficient overhead watering following application to ensure the treatment is moved into the root zone.

Do not apply more than 7.5 fl oz (0.098 lb ai per acre) of CORAGEN® to the soil at planting.

Do not apply more than 10 fl oz (0.132 lb ai per acre) of CORAGEN® per crop by any combination of at plant soil application and drip chemigation. **For drip chemigation applications made in the second half of the crop growing cycle: translocation of CORAGEN® into aerial portions of the plant may take up to 7 - 10 days.**

Do not make more than 2 drip chemigation applications of CORAGEN® per crop.

Do not make more than one drip chemigation application per crop if an at plant application of CORAGEN® was made. Refer to the SOIL APPLICATION section of this label for additional guidance; also see the rate conversion chart for application rate per 1000 linear feet.

†† FOLIAR. For best performance use an effective adjuvant. See the "Use of Adjuvants" section of the label.

\* **Diamondback moth resistance management:** Do not apply CORAGEN® more than twice to any generation of diamondback moth or within any 30 day period. After the second application of CORAGEN® for diamondback moth, rotate to another effective insecticide with a different mode of action (i.e. a product with a different IRAC group number). Application(s) to the next generation of diamondback moth must be with an effective product with a different mode of action.

Do not apply less than 3.5 oz. of CORAGEN® per application per acre for diamondback moth control. Do not make more than 6 total applications per calendar year for control of diamondback moth at the same farm location.

\*\* Suppression only. Transplant water treatment only.

\*\*\*Suppression only. Use in conjunction with an effective adult whitefly control program.

**DUPONT™ CORAGEN® RATE**

Crop	Application Method	Target Pest	Lb ai per acre	fluid ounces product per acre	Last Application (Days to Harvest)	REI (Hours)
Cereal Grains (EPA Crop Group 15) except Corn and Rice including: Barley, Buckwheat, Pearl Millet, Proso Millet, Oats, Rye, Sorghum (milo), Sorghum spp. [grain sorghum, sudangrass (seed crop), and hybrids of these grown for its seed], Teosinte, Triticale, Wheat, Wild Rice	FOLIAR	Corn earworm Beet armyworm European corn borer Fall armyworm Sorghum webworm Southwestern corn borer Sugarcane borer True armyworm	0.045 - 0.098	3.5 - 7.5	1	4
		Grasshoppers	0.026 - 0.065	2.0 - 5.0		

**USE RESTRICTIONS**  
 Make no more than 4 applications per acre per crop.  
 Minimum interval between treatments is 7 days.  
 Do not apply more than 15.4 fl oz CORAGEN® or 0.2 lb a.i. of chlorantraniliprole containing products per acre per calendar year.  
 CORAGEN® may be applied by overhead sprinkler chemigation systems. See "CHEMIGATION USING OVERHEAD SPRINKLER SYSTEMS - CEREAL GRAINS, CORN (FIELD, POP, SEED) COTTON GRASS FORAGE, FODDER, and HAY, LEGUMES, MINT (PEPPERMINT AND SPEARMINT), NON-GRASS ANIMAL FEEDS, OILSEED GROUP, PEANUT, POTATO, SOYBEAN, AND SUGARCANE" section for instructions on overhead sprinkler chemigation.  
**Grasshopper** - Apply foliarly when grasshopper populations reach local established thresholds to prevent crop damage. Correct timing of spray applications to nymphal stages and thorough coverage is critical to achieve optimum control. For best results, add Methylated Seed Oil (MSO) adjuvant at 1 gallon per 100 gallons of spray volume (1% v/v) when eggs have hatched and the majority of the grasshopper population is 2nd – 3rd instar nymphs. Once grasshoppers contact and/or ingest CORAGEN® there will be rapid feeding cessation; full expression of mortality may take a week or more. Do not make more than two sequential applications of CORAGEN® before rotating to another registered insecticide having a different mode-of-action.

Crop	Application Method	Target Pest	Lb ai per acre	fluid ounces product per acre	Last Application (Days to Harvest)	REI (Hours)
Forage, fodder, and Straw of Cereal Grains, (EPA Crop Group 16) except Corn and Rice. Including Forage, fodder, and straw of all commodities included in the cereal grains group, except corn and rice. Includes Sorghum spp. [sorghum, forage; sorghum, stover; sudangrass, and hybrids of these grown for forage and/or stover].	FOLIAR	Corn earworm Beet armyworm European corn borer Fall armyworm Sorghum webworm Southwestern corn borer Sugarcane borer True armyworm	0.045 - 0.098	3.5 - 7.5	1	4
		Grasshoppers	0.026 - 0.065	2.0 - 5.0		

**USE RESTRICTIONS**  
 Make no more than 4 applications per acre per crop.  
 Minimum interval between treatments is 7 days.  
 Do not apply more than 15.4 fl oz CORAGEN® or 0.2 lb a.i. of chlorantraniliprole containing products per acre per calendar year.  
 CORAGEN® may be applied by overhead sprinkler chemigation systems. See "CHEMIGATION USING OVERHEAD SPRINKLER SYSTEMS - CEREAL GRAINS, CORN (FIELD, POP, SEED) COTTON GRASS FORAGE, FODDER, and HAY, LEGUMES, MINT (PEPPERMINT AND SPEARMINT), NON-GRASS ANIMAL FEEDS, OILSEED GROUP, PEANUT, POTATO, SOYBEAN, AND SUGARCANE" section for instructions on overhead sprinkler chemigation.  
**Grasshopper** - Apply foliarly when grasshopper populations reach local established thresholds to prevent crop damage. Correct timing of spray applications to nymphal stages and thorough coverage is critical to achieve optimum control. For best results, add Methylated Seed Oil (MSO) adjuvant at 1 gallon per 100 gallons of spray volume (1% v/v) when eggs have hatched and the majority of the grasshopper population is 2nd – 3rd instar nymphs. Once grasshoppers contact and/or ingest CORAGEN® there will be rapid feeding cessation; full expression of mortality may take a week or more. Do not make more than two sequential applications of CORAGEN® before rotating to another registered insecticide having a different mode-of-action.

**DUPONT™ CORAGEN® RATE**

Crop	Application Method	Target Pest	Lb ai per acre	fluid ounces product per acre	Last Application (Days to Harvest)	REI (Hours)
Corn (field) Corn (pop)	SOIL AT PLANTING† In-furrow spray	European corn borer	0.065 - 0.098	5.0 - 7.5	14	4
		Fall armyworm Southern armyworm		See rate conversion chart for rate per 1000 linear ft.		
	FOLIAR	Corn earworm Beet armyworm European corn borer Fall armyworm Southern armyworm Southwestern corn borer True armyworm Western bean cutworm	0.045 - 0.098	3.5 - 7.5		
		Grasshoppers	0.026 - 0.065	2.0 - 5.0		

**USE RESTRICTIONS**

Make no more than 4 applications per acre per crop.  
 Minimum interval between treatments is 7 days.  
 Do not apply more than 15.4 fl oz CORAGEN® or 0.2 lb a.i. of chlorantraniliprole containing products, whether applications are made to the soil, foliarly or as a seed treatment per acre per calendar year.  
 CORAGEN® may be applied by overhead sprinkler chemigation systems. See "CHEMIGATION USING OVERHEAD SPRINKLER SYSTEMS - CEREAL GRAINS, CORN (FIELD, POP, SEED) COTTON GRASS FORAGE, FODDER, and HAY, LEGUMES, MINT (PEPPERMINT AND SPEARMINT), NON-GRASS ANIMAL FEEDS, OILSEED GROUP, PEANUT, POTATO, SOYBEAN, AND SUGARCANE" section for instructions on overhead sprinkler chemigation.  
 †**SOIL APPLICATIONS:**  
 In-Furrow Spray at Planting  
 Apply as a narrow band spray into the furrow at the seeding depth.  
 CORAGEN® must be applied in a manner that ensures the product is in the root zone. CORAGEN® is most effective when it is applied so that the roots are at or near the site of application; manage irrigation so that significant quantities of CORAGEN® remain in the root zone where it is most effective. Unless directed otherwise in the specific crop sections of this label, only one soil application of CORAGEN® may be made per crop.  
**Grasshopper** - Apply foliarly when grasshopper populations reach local established thresholds to prevent crop damage. Correct timing of spray applications to nymphal stages and thorough coverage is critical to achieve optimum control. For best results, add Methylated Seed Oil (MSO) adjuvant at 1 gallon per 100 gallons of spray volume (1% v/v) when eggs have hatched and the majority of the grasshopper population is 2nd – 3rd instar nymphs. Once grasshoppers contact and/or ingest CORAGEN® there will be rapid feeding cessation; full expression of mortality may take a week or more. Do not make more than two sequential applications of CORAGEN® before rotating to another registered insecticide having a different mode-of-action.

Crop	Application Method	Target Pest	Lb ai per acre	fluid ounces product per acre	Last Application (Days to Harvest)	REI (Hours)
Corn (sweet) Corn (grown for seed)	SOIL AT PLANTING† In-furrow spray	European corn borer	0.065 - 0.098	5.0 - 7.5	1	4
		Fall armyworm Southern armyworm		See rate conversion chart for rate per 1000 linear ft.		
	FOLIAR	Corn earworm Beet armyworm European corn borer Fall armyworm Southern armyworm Southwestern corn borer True armyworm Western bean cutworm	0.045 - 0.098	3.5 - 7.5		
		Grasshoppers	0.026 - 0.065	2.0 - 5.0		

**USE RESTRICTIONS**

Make no more than 4 applications per acre per crop.  
 Minimum interval between treatments is 1 day.  
 Do not apply more than 15.4 fl oz CORAGEN® or 0.2 lb a.i. of chlorantraniliprole containing products whether applications are made to the soil, foliarly or as a seed treatment per acre per calendar year.  
 CORAGEN® may be applied by overhead sprinkler chemigation systems. See "CHEMIGATION USING OVERHEAD SPRINKLER SYSTEMS - CEREAL GRAINS, CORN (FIELD, POP, SEED) COTTON GRASS FORAGE, FODDER, and HAY, LEGUMES, MINT (PEPPERMINT AND SPEARMINT), NON-GRASS ANIMAL FEEDS, OILSEED GROUP, PEANUT, POTATO, SOYBEAN, AND SUGARCANE" section for instructions on overhead sprinkler chemigation.  
 †**SOIL APPLICATIONS:**  
 In-Furrow Spray at Planting  
 Apply as a narrow band spray into the furrow at the seeding depth.  
 CORAGEN® must be applied in a manner that ensures the product is in the root zone. CORAGEN® is most effective when it is applied so that the roots are at or near the site of application; manage irrigation so that significant quantities of CORAGEN® remain in the root zone where it is most effective. Unless directed otherwise in the specific crop sections of this label, only one soil application of CORAGEN® may be made per crop.  
**Grasshopper** - Apply foliarly when grasshopper populations reach local established thresholds to prevent crop damage. Correct timing of spray applications to nymphal stages and thorough coverage is critical to achieve optimum control. For best results, add Methylated Seed Oil (MSO) adjuvant at 1 gallon per 100 gallons of spray volume (1% v/v) when eggs have hatched and the majority of the grasshopper population is 2nd – 3rd instar nymphs. Once grasshoppers contact and/or ingest CORAGEN® there will be rapid feeding cessation; full expression of mortality may take a week or more. Do not make more than two sequential applications of CORAGEN® before rotating to another registered insecticide having a different mode-of-action.

**DUPONT™ CORAGEN® RATE**

Crop	Application Method	Target Pest	Lb ai per acre	fluid ounces product per acre	Last Application (Days to Harvest)	REI (Hours)
Cotton	FOLIAR	Beet armyworm Cotton bollworm** Fall armyworm Saltmarsh caterpillar Southern armyworm Tobacco budworm** Western Yellowstriped Armyworm	0.045 - 0.098	3.5 - 7.5	21	4
		Cabbage looper Soybean looper*	0.065 - 0.098	5.0 - 7.5		
		Grasshoppers	0.026 - 0.065	2.0 - 5.0		

**USE RESTRICTIONS**

Make no more than 4 applications per acre per crop.

Do not apply more than 15.4 fl oz CORAGEN® or 0.2 lb a.i. of chlorantraniliprole containing products per acre per calendar year.

The minimum interval between treatments is 5 days.

CORAGEN® may be applied by overhead sprinkler chemigation systems. See "CHEMIGATION USING OVERHEAD SPRINKLER SYSTEMS - CEREAL GRAINS, CORN (FIELD, POP, SEED) COTTON GRASS FORAGE, FODDER, and HAY, LEGUMES, MINT (PEPPERMINT AND SPEARMINT), NON-GRASS ANIMAL FEEDS, OILSEED GROUP, PEANUT, POTATO, SOYBEAN, AND SUGARCANE" section for instructions on overhead sprinkler chemigation.

\*Suppression only.

\*\*For Heliothine control (cotton bollworm and/or tobacco budworm in conventional non-transgenic/non-Bt cotton) make the first application at rates of 0.065 - 0.09 lb. ai per acre (5.0 - 7.0 oz product). Subsequent applications can be at rates of 0.045 - 0.09 lb. ai acre (3.5 - 7.0 oz product) depending on pest pressure.

For control of cotton bollworm (*Helicoverpa zea*) in Bt transgenic cotton varieties, the initial application, and subsequent applications, of CORAGEN® can be applied at 3.5 to 5 fluid ounces per acre as a foliar spray. Apply when cotton bollworm populations reach local established treatment thresholds to prevent crop damage.

**Grasshopper** - Apply foliarly when grasshopper populations reach local established thresholds to prevent crop damage. Correct timing of spray applications to nymphal stages and thorough coverage is critical to achieve optimum control. For best results, add Methylated Seed Oil (MSO) adjuvant at 1 gallon per 100 gallons of spray volume (1% v/v) when eggs have hatched and the majority of the grasshopper population is 2nd – 3rd instar nymphs. Once grasshoppers contact and/or ingest CORAGEN® there will be rapid feeding cessation; full expression of mortality may take a week or more. Do not make more than two sequential applications of CORAGEN® before rotating to another registered insecticide having a different mode-of-action.



**DUPONT™ CORAGEN® RATE**

<b>Crop</b>	<b>Application Method</b>	<b>Target Pest</b>	<b>Lb ai per acre</b>	<b>fluid ounces product per acre</b>	<b>Last Application (Days to Harvest)</b>	<b>REI (Hours)</b>
Cucurbit Vegetables, (EPA Crop Group 9) Including: Chayote (fruit), Chinese wax-gourd (Chinese preserving melon), Citron melon, Cucumber, Gherkin, Edible gourd (includes hyotan, cucuzza, hechima, Chinese okra), Momordica spp. (includes balsam apple, balsam pear, bitter melon, Chinese cucumber), Muskmelon (includes true cantaloupe, cantaloupe, casaba, crenshaw melon, golden pershaw melon, honeydew melon, honey balls, mango melon, Persian melon, pineapple melon, Santa Claus melon, and snake melon), Pumpkin, Summer squash (includes crookneck squash, scallop squash, straightneck squash, vegetable marrow, zucchini), Winter squash (includes butternut squash, calabaza, hubbard squash, acorn squash, spaghetti squash), Watermelon	SOIL AT PLANTING† (an in-furrow spray, transplant water treatment, hill drench, surface band, soil shank injection)	Beet armyworm	0.045 - 0.098	3.5 - 7.5	1	4
		Cabbage looper		See rate conversion chart for rate per 1000 linear ft.		
	DRIP	Leafminers (larvae)*	0.065 - 0.098	5.0 - 7.5		
		Silverleaf whiteflies (nymphs)**				
	CHEMIGATION† (Make application(s) within the first half of the crop growing cycle, typically up to peak bloom crop stage.	Melon worm	0.026 - 0.045	2.0 - 3.5		
		Beet armyworm	0.045 - 0.098	3.5 - 7.5		
	FOLIAR	Cabbage looper				
		Pickle worm				
	FOLIAR	Leafminers (larvae)*	0.065 - 0.098	5.0 - 7.5		
		Silverleaf whiteflies (nymphs)**				
FOLIAR	Melon worm	0.026 - 0.045	2.0 - 3.5			
	Beet armyworm	0.045 - 0.098	3.5 - 7.5			
FOLIAR	Cabbage looper					
	Hawaiian beet webworm					
FOLIAR	Pickle worm					
	Western yellowstriped armyworm					
FOLIAR	Leafminers (larvae)*	0.065 - 0.098	5.0 - 7.5			
	Silverleaf whiteflies (nymphs)**					

**USE RESTRICTIONS**  
 Make no more than 4 applications per acre per crop.  
 Minimum interval between treatments is 5 days for foliar applications and 10 days for drip chemigation applications.  
 Do not apply more than 15.4 fl oz CORAGEN® or 0.2 lb a.i. of chlorantraniliprole containing products per acre per crop.  
 Do not apply more than 46.2 fl oz CORAGEN® or 0.6 lb a.i. of chlorantraniliprole containing products per acre per calendar year; in NY do not apply more than 15.4 fl oz CORAGEN® or 0.2 lb a.i. of chlorantraniliprole containing products per acre per calendar year.  
 †**SOIL APPLICATIONS** (an in-furrow spray at planting, transplant water treatment, hill drench at planting, surface band at planting, soil shank injection at planting, or drip chemigation): CORAGEN® must be applied uniformly in the root zone or poor performance will result. Surface band application requires sufficient overhead watering following application in to ensure the treatment is moved into the root zone. Do not apply more than 7.5 fl oz (0.098 lb ai per acre) of CORAGEN® to the soil at planting.  
 Do not apply more than 10 fl oz (0.132 lb ai per acre) of CORAGEN® per crop by any combination of at plant soil application and drip chemigation.  
 Do not make more than 2 drip chemigation applications of CORAGEN® per crop.  
 Do not make more than one drip chemigation application per crop if an at plant application of CORAGEN® was made. Refer to the SOIL APPLICATION section of this label for additional guidance; also see the rate conversion chart for application rate per 1000 linear feet.  
 \*Control of Liriomyza species except suppression only for *L. huidabrensis* and *L. langei*.  
 \*\*Suppression only. Use in conjunction with an effective adult whitefly control program.

**DUPONT™ CORAGEN® RATE**

<b>Crop</b>	<b>Application Method</b>	<b>Target Pest</b>	<b>Lb ai per acre</b>	<b>fluid ounces product per acre</b>	<b>Last Application (Days to Harvest)</b>	<b>REI (Hours)</b>
Fruiting Vegetables Including: Eggplant, Groundcherry (Physalis spp.), okra, Pepino, Pepper, (including bell pepper, chili pepper, cooking pepper, pimento, sweet pepper), Tomatillo, Tomato	SOIL AT PLANTING† (an in-furrow spray, transplant water treatment, hill drench, surface band, soil shank injection)	Beet armyworm Fall armyworm Loopers Southern armyworm Tomato fruitworm Tomato pinworm Western yellow striped armyworm	0.045 - 0.098	3.5 - 7.5 See rate conversion chart for rate per 1000 linear ft.	1	4
		Leafminers (larvae)* Silverleaf whiteflies (nymphs)**	0.065 - 0.098	5.0 - 7.5		
	DRIP CHEMIGATION†	Beet armyworm Colorado potato beetle European corn borer Fall armyworm Garden webworm Hornworms Loopers Southern armyworm Tomato fruitworm Tomato pinworm Western yellow striped armyworm	0.045 - 0.098	3.5 - 7.5		
		Leafminers (larvae)* Silverleaf whiteflies (nymphs)**	0.065 - 0.098	5.0 - 7.5		
	FOLIAR	Hornworms	0.026 - 0.065	2.0 - 5.0		
		Beet armyworm Colorado potato beetle European corn borer Fall armyworm Garden webworm Loopers Southern armyworm Tomato fruitworm Tomato pinworm Western yellow striped armyworm	0.045 - 0.098	3.5 - 7.5		
		Leafminers (larvae)* Silverleaf whiteflies (nymphs)**	0.065 - 0.098	5.0 - 7.5		

**USE RESTRICTIONS**  
 Make no more than 4 applications per acre per crop.  
 Minimum interval between treatments is 5 days for foliar applications and 10 days for drip chemigation applications.  
 Do not apply more than 15.4 fl oz CORAGEN® or 0.2 lb a.i. of chlorantraniliprole containing products per acre per crop.  
 Do not apply more than 46.2 fl oz CORAGEN® or 0.6 lb a.i. of chlorantraniliprole containing products per acre per calendar year; in NY do not apply more than 15.4 fl oz CORAGEN® or 0.2 lb a.i. of chlorantraniliprole containing products per acre per calendar year.  
 \* Control of Liriomyza species except suppression only for *L. huidabrensis* and *L. langei*.  
 \*\* Suppression only. Use in conjunction with an effective adult whitefly control program.  
 †**SOIL APPLICATIONS** (an in-furrow spray at planting, transplant water treatment, hill drench at planting, surface band at planting, soil shank injection at planting, or drip chemigation): CORAGEN® must be applied uniformly in the root zone or poor performance will result. Surface band application requires sufficient overhead watering following application to ensure the treatment is moved into the root zone. Do not apply more than 7.5 fl oz (0.098 lb ai per acre) of CORAGEN® to the soil at planting. Do not apply more than 10 fl oz (0.132 lb ai per acre) of CORAGEN® per crop by any combination of at plant soil application and drip chemigation. **For drip chemigation applications made in the second half of the crop growing cycle: translocation of CORAGEN® into aerial portions of the plant may take up to 7 - 10 days.**  
 Do not make more than 2 drip chemigation applications of CORAGEN® per crop .  
 Do not make more than one drip chemigation application per crop if an at plant application of CORAGEN® was made. Refer to the SOIL APPLICATION section of this label for additional guidance; also see the rate conversion chart for application rate per 1000 linear feet.

**DUPONT™ CORAGEN® RATE**

Crop	Application Method	Target Pest	Lb ai per acre	fluid ounces product per acre	Last Application (Days to Harvest)	REI (Hours)
Grass Forage, Fodder and Hay: (EPA Crop Group 17) Any grass, Gramineae family (either green or cured) except sugarcane and those included in the cereal grains group, that will be fed to or grazed by livestock, all pasture and range grasses and grasses grown for hay or silage	FOLIAR	Beet armyworm	0.045 - 0.098	3.5 - 7.5	0	4
		Corn earworm				
		Fall armyworm	0.026 - 0.065	2.0 - 5.0		
		Sod webworm	0.065 - 0.098	5.0 - 7.5		
		Southern armyworm				
		True armyworm				
		Grasshoppers				
		Billbug (grubs)*				
		Cutworms				
		European crane fly (larvae)*				

**USE RESTRICTIONS**  
 Make no more than 4 applications per acre per crop.  
 Minimum interval between treatments is 7 days.  
 Do not apply more than 15.4 fl oz CORAGEN® or 0.2 lb a.i. of chlorantraniliprole containing products per acre per calendar year.  
 CORAGEN® may be applied by overhead sprinkler chemigation systems. See "CHEMIGATION USING OVERHEAD SPRINKLER SYSTEMS - CEREAL GRAINS, CORN (FIELD, POP, SEED) COTTON GRASS FORAGE, FODDER, and HAY, LEGUMES, MINT (PEPPERMINT AND SPEARMINT), NON-GRASS ANIMAL FEEDS, OILSEED GROUP, PEANUT, POTATO, SOYBEAN, AND SUGARCANE" section for instructions on overhead sprinkler chemigation.  
 \* Suppression only. Grass grown for seed only.  
**Grasshopper** - Apply foliarly when grasshopper populations reach local established thresholds to prevent crop damage. Correct timing of spray applications to nymphal stages and thorough coverage is critical to achieve optimum control. For best results, add Methylated Seed Oil (MSO) adjuvant at 1 gallon per 100 gallons of spray volume (1% v/v) when eggs have hatched and the majority of the grasshopper population is 2nd – 3rd instar nymphs. Once grasshoppers contact and/or ingest CORAGEN® there will be rapid feeding cessation; full expression of mortality may take a week or more. Do not make more than two sequential applications of CORAGEN® before rotating to another registered insecticide having a different mode-of-action.

Crop	Application Method	Target Pest	Lb ai per acre	fluid ounces product per acre	Last Application (Days to Harvest)	REI (Hours)
Herb subgroup (EPA Crop Subgroup 19A) Including Angelica; balm; basil; borage; burnet; camomile; catnip; chervil (dried); chive, Chinese; clary; coriander (leaf); costmary; culantro (leaf); curry (leaf); dillweed; horehound; hyssop; lavender; lemongrass; lovage (leaf); marigold; marjoram; nasturtium; parsley (dried); pennyroyal; rosemary; rue; sage; savory, summer and winter; sweet bay; tansy; tarragon; thyme; wintergreen; woodruff; and wormwood	FOLIAR	Beet armyworm Cabbage looper Corn earworm Fall armyworm Southern armyworm	0.045 - 0.065	3.5 - 5.0	1	4

**USE RESTRICTIONS**  
 Make no more than 4 applications per acre per crop.  
 Minimum interval between treatments is 3 days.  
 Do not apply more than 15.4 fl oz CORAGEN® or 0.2 lb a.i. of chlorantraniliprole containing products per acre per crop.  
 Do not apply more than 61.6 fl oz CORAGEN® or 0.8 lb a.i. of chlorantraniliprole containing products per acre per calendar year; in NY do not apply more than 15.4 fl oz CORAGEN® or 0.2 lb a.i. of chlorantraniliprole containing products per acre per calendar year.  
**PLANT TOLERANCE PHYTOTOXICITY**  
 CORAGEN® has been tested on numerous crops and cultivars with no observable phytotoxicity at label rates. However, neither the manufacturer nor the seller has determined whether or not CORAGEN® can be used safely on all herbs and spices for which it is registered for use.  
 Since all herbs and spices and their varieties and cultivars have not been tested for phytotoxicity it is recommended that a small number of plants be sprayed initially to determine if there is any phytotoxicity prior to large scale applications to herbs and spices. The user assumes all risks arising from application of CORAGEN® in a manner that is inconsistent with its labeling.

**DUPONT™ CORAGEN® RATE**

Crop	Application Method	Target Pest	Lb ai per acre	fluid ounces product per acre	Last Application (Days to Harvest)	REI (Hours)
Hops (except California)	FOLIAR	Western yellowstriped armyworm	0.045 - 0.098	3.5 - 7.5	0	4

**USE RESTRICTIONS**

Make no more than 4 applications per acre per crop.  
 Minimum interval between treatments is 7 days.  
 Do not apply more than 15.4 fl oz CORAGEN® or 0.2 lb a.i. of chlorantraniliprole containing products per acre per calendar year.

Crop	Application Method	Target Pest	Lb ai per acre	fluid ounces product per acre	Last Application (Days to Harvest)	REI (Hours)
Leafy Vegetables except brassica (EPA Crop Group 4) Including: Amaranth leafy, Arugula (roquette), Cardoon, Celery, Celery (chinese), Celtuce, Chevril, Chinese spinach, Chrysanthemum (edible leaved), Chrysanthemum, garland, Corn salad, Cress (garden), Cress (upland), Dandelion, leaves, Dock (sorrel), Endive (escarole), Florence fennel, Lettuce (head & leaf), Orach, Parsley, Purslane (garden), (winter), Radicchio (red chicory), Rhubarb, Spinach, Spinach (vine), Spinach (New Zealand), Swiss chard, Tampala	SOIL AT PLANTING† (an in-furrow spray, transplant water treatment, hill drench, surface band, soil shank injection)	Beet armyworm Corn earworm Cabbage looper Tobacco budworm	0.045 - 0.098	3.5 - 7.5 See rate conversion chart for rate per 1000 linear ft.	1	4
		Leafminers (larvae)** Silverleaf whiteflies (nymphs)***	0.065 - 0.098	5.0 - 7.5		
	DRIP CHEMIGATION†	Diamondback moth* Beet armyworm Corn earworm Cabbage looper Hawaiian beet webworm Tobacco budworm	0.045 - 0.098	3.5 - 7.5		
		Leafminers (larvae)** Silverleaf whiteflies (nymphs)***	0.065 - 0.098	5.0 - 7.5		
	FOLIAR	Corn earworm Diamondback moth* Beet armyworm Cabbage looper Hawaiian beet webworm Tobacco budworm Western yellowstriped armyworm	0.045 - 0.098	3.5 - 7.5		
		Leafminers (larvae)** Silverleaf whiteflies (nymphs)***	0.065 - 0.098	5.0 - 7.5		

**USE RESTRICTIONS**

Make no more than 4 applications per acre per crop.  
 Minimum interval between treatments is 3 days for foliar applications and 10 days for drip chemigation applications.  
 Do not apply more than 15.4 fl oz CORAGEN® or 0.2 lb a.i. of chlorantraniliprole containing products per acre per crop.  
 Do not apply more than 61.6 fl oz CORAGEN® or 0.8 lb a.i. of chlorantraniliprole containing products per acre per calendar year; in NY do not apply more than 15.4 fl oz CORAGEN® or 0.2 lb a.i. of chlorantraniliprole containing products per acre per calendar year.  
 \* Diamondback moth resistance management: Do not apply CORAGEN® more than twice to any generation of diamondback moth or within any 30 day period. After the second application of CORAGEN® for diamondback moth, rotate to another effective insecticide with a different mode of action (i.e. a product with a different IRAC group number). Application(s) to the next generation of diamondback moth must be with an effective product with a different mode of action. Do not apply less than 3.5 oz. of CORAGEN® per application per acre for diamondback moth control. Do not make more than 6 total applications per calendar year for control of diamondback moth at the same farm location.  
 \*\* Control of *Liriomyza* species except suppression only for *L. huidabrensis* and *L. langei*.  
 \*\*\* Suppression only. Use in conjunction with an effective adult whitefly control program.  
 †**SOIL APPLICATIONS** (an in-furrow spray at planting, transplant water treatment, hill drench at planting, surface band at planting, soil shank injection at planting, or drip chemigation): CORAGEN® must be applied uniformly in the root zone or poor performance will result. Surface band application requires sufficient watering in to ensure the treatment is moved into the root zone. Do not apply more than 7.5 fl oz (0.098 lb ai per acre) of CORAGEN® to the soil at planting. Do not apply more than 10 fl oz (0.132 lb ai per acre) of CORAGEN® per crop by any combination of at plant soil application and drip chemigation. Do not make more than 2 drip chemigation applications of CORAGEN® per crop. **For drip chemigation applications made in the second half of the crop growing cycle: translocation of CORAGEN® into aerial portions of the plant may take up to 7 - 10 days.**  
 Do not make more than one drip chemigation application per crop if an at plant application of CORAGEN® was made. Refer to the SOIL APPLICATION section of this label for additional guidance; also see the rate conversion chart for application rate per 1000 linear feet.

**DUPONT™ CORAGEN® RATE**

<b>Crop</b>	<b>Application Method</b>	<b>Target Pest</b>	<b>Lb ai per acre</b>	<b>fluid ounces product per acre</b>	<b>Last Application (Days to Harvest)</b>	<b>REI (Hours)</b>
Legume vegetables (EPA Crop Group 6) (For soybean see separate soybean crop section below.) (Succulent or Dried, Including Bean (Lupinus) (includes grain lupin, sweet lupin, white lupin, and white sweet lupin); bean (Phaseolus) (includes field bean, kidney bean, lima bean, navy bean, pinto bean, runner bean, snap bean, tepary bean, wax bean); bean (Vigna) (includes adzuki bean, asparagus bean, blackeyed pea, catjang, Chinese longbean, cowpea, crowder pea, moth bean, mung bean, rice bean, southern pea, urd bean, yardlong bean); broad bean (fava); chickpea (garbanzo); guar; jackbean; lablab bean; lentil; pea (Pisum) (includes dwarf pea, edible-podded pea, English pea, field pea, garden pea, green pea, snowpea, sugar snap pea); pigeon pea; sword bean	SOIL AT PLANTING† In-furrow spray	Corn earworm Beet armyworm European corn borer Fall armyworm	0.065 - 0.098	5.0 - 7.5 See rate conversion chart for rate per 1000 linear ft.	1	4
	FOLIAR	Corn earworm Beet armyworm European corn borer Fall armyworm Cabbage looper Soybean looper Western bean cutworm	0.045 - 0.098	3.5 - 7.5		
		Leafminers (larvae)* Silverleaf whiteflies (nymphs)**	0.098	7.5		
		Grasshoppers	0.026 - 0.065	2.0 - 5.0		

**USE RESTRICTIONS**  
 Make no more than 4 applications per acre per crop.  
 Minimum interval between treatments is 3 days.  
 Do not apply more than 15.4 fl oz CORAGEN® or 0.2 lb a.i. of chlorantraniliprole containing products per acre per crop.  
 Do not apply more than 46.2 fl oz CORAGEN® or 0.6 lb a.i. of chlorantraniliprole containing products per acre per calendar year; in NY do not apply more than 15.4 fl oz CORAGEN® or 0.2 lb a.i. of chlorantraniliprole containing products per acre per calendar year.  
 CORAGEN® may be applied by overhead sprinkler chemigation systems. See "CHEMIGATION USING OVERHEAD SPRINKLER SYSTEMS - CEREAL GRAINS, CORN (FIELD, POP, SEED, SWEET[?]) COTTON GRASS FORAGE, FODDER, and HAY, LEGUMES, MINT (PEPPERMINT AND SPEARMINT), NON-GRASS ANIMAL FEEDS, OILSEED GROUP, PEANUT, POTATO, SOYBEAN, AND SUGARCANE" section for instructions on overhead sprinkler chemigation.  
 \*Control of *Liriomyza* species except suppression only for *L. huidabrensis* and *L. langei*.  
 \*\*Suppression only. Use in conjunction with an effective adult whitefly control program.  
 †SOIL APPLICATIONS:  
 In-Furrow Spray at Planting  
 Apply as a narrow band spray into the furrow at the seeding depth. CORAGEN® must be applied in a manner that ensures the product is in the root zone. CORAGEN® must be in the root zone to provide effective control of target pests. CORAGEN® is most effective when it is applied so that the roots are at or near the site of application; manage irrigation so that significant quantities of CORAGEN® remain in the root zone where it is most effective. Unless directed otherwise in the specific crop sections of this label, only one soil application of CORAGEN® may be made per crop.  
**Grasshopper** - Apply foliarly when grasshopper populations reach local established thresholds to prevent crop damage. Correct timing of spray applications to nymphal stages and thorough coverage is critical to achieve optimum control. For best results, add Methylated Seed Oil (MSO) adjuvant at 1 gallon per 100 gallons of spray volume (1% v/v) when eggs have hatched and the majority of the grasshopper population is 2nd – 3rd instar nymphs. Once grasshoppers contact and/or ingest CORAGEN® there will be rapid feeding cessation; full expression of mortality may take a week or more. Do not make more than two sequential applications of CORAGEN® before rotating to another registered insecticide having a different mode-of-action.

**DUPONT™ CORAGEN® RATE**

Crop	Application Method	Target Pest	Lb ai per acre	fluid ounces product per acre	Last Application (Days to Harvest)	REI (Hours)
Foliage of Legume Vegetables (EPA Crop Group 7) except soybean including: of any legume vegetable included in the legume vegetables that will be used as animal feed.	SOIL AT PLANTING† In-furrow spray	Corn earworm Beet armyworm European corn borer Fall armyworm	0.065 - 0.098	5.0 - 7.5 See rate conversion chart for rate per 1000 linear ft.	1	4
	FOLIAR	Corn earworm Beet armyworm European corn borer Fall armyworm Cabbage looper Soybean looper Western bean cutworm	0.045 - 0.098	3.5 - 7.5		
		Leafminers (larvae)* Silverleaf whiteflies (nymphs)**	0.098	7.5		
		Grasshoppers	0.026 - 0.065	2.0 - 5.0		

**USE RESTRICTIONS**

Make no more than 4 applications per acre per crop.  
 Minimum interval between treatments is 3 days.  
 Do not apply more than 15.4 fl oz CORAGEN® or 0.2 lb a.i. of chlorantraniliprole containing products per acre per crop.  
 Do not apply more than 46.2 fl oz CORAGEN or 0.6 lb a.i. of chlorantraniliprole containing products per acre per calendar year; in NY do not apply more than 15.4 fl oz CORAGEN® or 0.2 lb a.i. of chlorantraniliprole containing products per acre per calendar year.  
 CORAGEN® may be applied by overhead sprinkler chemigation systems. See "CHEMIGATION USING OVERHEAD SPRINKLER SYSTEMS - CEREAL GRAINS, CORN (FIELD, POP, SEED) COTTON GRASS FORAGE, FODDER, and HAY, LEGUMES, MINT (PEPPERMINT AND SPEARMINT), NON-GRASS ANIMAL FEEDS, OILSEED GROUP, PEANUT, POTATO, SOYBEAN, AND SUGARCANE" section for instructions on overhead sprinkler chemigation.  
 \*Control of *Liriomyza* species except suppression only for *L. huidabrensis* and *L. langei*.  
 \*\*Suppression only. Use in conjunction with an effective adult whitefly control program.  
 †**SOIL APPLICATIONS:**  
 In-Furrow Spray at Planting  
 Apply as a narrow band spray into the furrow at the seeding depth. CORAGEN® must be applied in a manner that ensures the product is in the root zone. CORAGEN® must be in the root zone to provide effective control of target pests. CORAGEN® is most effective when it is applied so that the roots are at or near the site of application; manage irrigation so that significant quantities of CORAGEN® remain in the root zone where it is most effective. Unless directed otherwise in the specific crop sections of this label, only one soil application of CORAGEN® may be made per crop.  
**Grasshopper** - Apply foliarly when grasshopper populations reach local established thresholds to prevent crop damage. Correct timing of spray applications to nymphal stages and thorough coverage is critical to achieve optimum control. For best results, add Methylated Seed Oil (MSO) adjuvant at 1 gallon per 100 gallons of spray volume (1% v/v) when eggs have hatched and the majority of the grasshopper population is 2nd – 3rd instar nymphs. Once grasshoppers contact and/or ingest CORAGEN® there will be rapid feeding cessation; full expression of mortality may take a week or more. Do not make more than two sequential applications of CORAGEN® before rotating to another registered insecticide having a different mode-of-action.

Crop	Application Method	Target Pest	Lb ai per acre	fluid ounces product per acre	Last Application (Days to Harvest)	REI (Hours)
Mint: Peppermint and Spearmint	FOLIAR	Armyworms Cutworms Loopers Mint root borer	0.045 - 0.098	3.5 - 7.5	3	4

**USE RESTRICTIONS**

Make no more than 4 applications per acre per crop.  
 Minimum interval between treatments is 14 days.  
 Do not apply more than 15.4 fl oz CORAGEN® or 0.2 lb a.i. of chlorantraniliprole containing products per acre per calendar year.  
 CORAGEN® may be applied by overhead sprinkler chemigation systems. See "CHEMIGATION USING OVERHEAD SPRINKLER SYSTEMS - CEREAL GRAINS, CORN (FIELD, POP, SEED) COTTON GRASS FORAGE, FODDER, and HAY, LEGUMES, MINT (PEPPERMINT AND SPEARMINT), NON-GRASS ANIMAL FEEDS, OILSEED GROUP, PEANUT, POTATO, SOYBEAN, AND SUGARCANE" section for instructions on overhead sprinkler chemigation.  
 In mint growing areas where the mint root borer degree day model is being used and mint is being grown under sprinkler irrigation: apply CORAGEN® at 5.0 fl oz/acre (0.065 lb a.i. per acre) as a foliar spray or via overhead sprinkler chemigation. Application should be timed between 900 and 1250 growing degree days. Foliar sprays must be followed by sprinkler irrigation before swathing. When making a foliar spray, be sure to include an adjuvant to help obtain thorough coverage. Use only adjuvant products that are labeled for agricultural use and follow the directions on the manufacturer's label. Always conduct a premix test for compatibility. Use an adjuvant that does not affect foliage.  
**Mint Root Borer** - For applications after the last cutting of mint, apply CORAGEN(r) soon after the last cutting of mint, but before the Mint Root Borer forms an overwintering hibernaculum. If CORAGEN® is applied as a broadcast spray, follow application with at least 2 inches water per acre of overhead irrigation. For furrow irrigated mint, apply CORAGEN® as a broadcast spray soon after harvest. Follow application with two furrow irrigations in order to move CORAGEN® into the mint root zone before the mint root borer forms a hibernaculum. If CORAGEN® is applied via overhead chemigation, use a minimum of 2 acre inches of water to move the CORAGEN® into the mint root zone.

**DUPONT™ CORAGEN® RATE**

Crop	Application Method	Target Pest	Lb ai per acre	fluid ounces product per acre	Last Application (Days to Harvest)	REI (Hours)
Non-grass animal feeds (EPA Crop Group 18) Including: Alfalfa;bean, velvet; clover (Trifolium, Melilotus); kudzu; lespedeza; lupin; sainfoin; trefoil; vetch; vetch, crown; vetch, milk	FOLIAR	Alfalfa caterpillar Alfalfa looper Beet armyworm Western yellowstriped armyworm	0.045 - 0.098	3.5 - 7.5	0	4
		Grasshoppers	0.026 - 0.065	2.0 - 5.0		
<p><b>USE RESTRICTIONS</b>            Make no more than 4 applications per acre per crop.            Make one application per cutting.            Do not apply more than 15.4 fl oz CORAGEN® or 0.2 lb a.i. of chlorantraniliprole containing products per acre per calendar year.            CORAGEN® may be applied by overhead sprinkler chemigation systems. See "CHEMIGATION USING OVERHEAD SPRINKLER SYSTEMS - CEREAL GRAINS, CORN (FIELD, POP, SEED) COTTON GRASS FORAGE, FODDER, and HAY, LEGUMES, MINT (PEPPERMINT AND SPEARMINT), NON-GRASS ANIMAL FEEDS, OILSEED GROUP, PEANUT, POTATO, SOYBEAN, AND SUGARCANE" section for instructions on overhead sprinkler chemigation.  <b>Grasshopper</b> - Apply foliarly when grasshopper populations reach local established thresholds to prevent crop damage. Correct timing of spray applications to nymphal stages and thorough coverage is critical to achieve optimum control. For best results, add Methylated Seed Oil (MSO) adjuvant at 1 gallon per 100 gallons of spray volume (1% v/v) when eggs have hatched and the majority of the grasshopper population is 2nd – 3rd instar nymphs. Once grasshoppers contact and/or ingest CORAGEN® there will be rapid feeding cessation; full expression of mortality may take a week or more. Do not make more than two sequential applications of CORAGEN® before rotating to another registered insecticide having a different mode-of-action.</p>						

Crop	Application Method	Target Pest	Lb ai per acre	fluid ounces product per acre	Last Application (Days to Harvest)	REI (Hours)
Peanut	FOLIAR	Corn earworm Beet armyworm Fall armyworm Green cloverworm Southern armyworm Tobacco budworm Velvetbean caterpillar	0.045 - 0.098	3.5 - 7.5	1	4
		Cabbage looper Granulate cutworm Soybean looper	0.065 - 0.098	5.0 - 7.5		
		Grasshoppers	0.026 - 0.065	2.0 - 5.0		
<p><b>USE RESTRICTIONS</b>            Make no more than 4 applications per acre per crop.            Minimum interval between treatments is 3 days.            Do not apply more than 15.4 fl oz CORAGEN® or 0.2 lb a.i. of chlorantraniliprole containing products per acre per calendar year.            CORAGEN® may be applied by overhead sprinkler chemigation systems. See "CHEMIGATION USING OVERHEAD SPRINKLER SYSTEMS - CEREAL GRAINS, CORN (FIELD, POP, SEED) COTTON GRASS FORAGE, FODDER, and HAY, MINT (PEPPERMINT AND SPEARMINT), LEGUMES, NON-GRASS ANIMAL FEEDS, OILSEED GROUP, PEANUT, POTATO, SOYBEAN, AND SUGARCANE" section for instructions on overhead sprinkler chemigation.  <b>Grasshopper</b> - Apply foliarly when grasshopper populations reach local established thresholds to prevent crop damage. Correct timing of spray applications to nymphal stages and thorough coverage is critical to achieve optimum control. For best results, add Methylated Seed Oil (MSO) adjuvant at 1 gallon per 100 gallons of spray volume (1% v/v) when eggs have hatched and the majority of the grasshopper population is 2nd – 3rd instar nymphs. Once grasshoppers contact and/or ingest CORAGEN® there will be rapid feeding cessation; full expression of mortality may take a week or more. Do not make more than two sequential applications of CORAGEN® before rotating to another registered insecticide having a different mode-of-action.</p>						

**DUPONT™ CORAGEN® RATE**

Crop	Application Method	Target Pest	Lb ai per acre	fluid ounces product per acre	Last Application (Days to Harvest)	REI (Hours)
Potato	FOLIAR	Beet and Yellowstriped Armyworms Cabbage looper Colorado potato beetle European corn borer Potato tuberworm	0.045 - 0.098	3.5 - 7.5	14	4
		Grasshoppers	0.026 - 0.065	2.0 - 5.0		

**USE RESTRICTIONS**  
 Make no more than 4 applications per acre per crop.  
 Do not apply more than 15.4 fl oz CORAGEN® or 0.2 lb a.i. of chlorantraniliprole containing products per acre per calendar year. The minimum interval between treatments is 5 days.

**Colorado potato beetle resistance management:** Do not apply CORAGEN® more than twice to a generation of Colorado potato beetle or within any 30 day period. Application(s) to the next generation of Colorado potato beetle must be with an effective product with a different mode of action.

**Potato tuberworm:** CORAGEN® may be applied at rates of 3.2 - 5.0 fl oz per acre to control potato tuberworm. Begin application when field scouting indicates the presence of tuberworm adults and/or larvae. Potato tuberworm often have overlapping generations so repeat applications of CORAGEN® may be needed based on field scouting. Avoid treating successive generations with the same mode of action. It is important to protect the crop just prior to harvest when foliage starts to senesce. Use the high rate of CORAGEN® where potato tuberworm pressure is high. Failure to adequately control potato tuberworm larvae prior to crop senescence or vine kill increases the risk of tuber damage. Foliar sprays alone, by air or ground, may not provide adequate control of larvae in the mid to lower crop canopy. For best results, apply via overhead chemigation or integrate chemigation applications into the foliar spray program. For best results with foliar sprays, add Methylated Seed Oil (MSO) adjuvant at 1 gallon per 100 gallons of spray volume (1% v/v). For chemigation applications, apply in 0.1 to 0.2 acre inches of water and add MSO at 12 to 16 fl oz/acre. CORAGEN® may be applied via overhead sprinkler chemigation systems.

Do not apply CORAGEN® more than once to Colorado potato beetle via overhead chemigation. CORAGEN® may only be applied to potatoes as a direct foliar spray or via chemigation through overhead sprinkler irrigation systems. See "CHEMIGATION USING OVERHEAD SPRINKLER SYSTEMS - CEREAL GRAINS, CORN (FIELD, POP, SEED) COTTON GRASS FORAGE, FODDER, and HAY, LEGUMES, MINT (PEPPERMINT AND SPEARMINT), NON-GRASS ANIMAL FEEDS, OILSEED GROUP, PEANUT, POTATO, SOYBEAN, AND SUGARCANE" section for instructions on overhead sprinkler chemigation.

**Cabbage looper:** West of the Rocky Mountains - CORAGEN® may be applied at 2.0 - 3.4 fl oz per acre ( 0.026 - 0.044 Lb ai/acre) to control early stage instars (1st - 3rd instar).

**Colorado potato beetle:** West of the Rocky Mountains - CORAGEN® may be applied at 2.0 - 3.4 fl oz per acre ( 0.026 - 0.044 LB ai/acre) to control local populations of Colorado Potato Beetle believed to be sensitive to most commonly used insecticides. Apply just prior to or just after egg hatch while larvae are small. In some areas, where local populations of Colorado Potato Beetle have elevated levels of resistance to insecticides, CORAGEN® should be used at the 5.0 fluid ounce per acre application rate. With resistant populations of Colorado Potato Beetle, back-to-back applications on 5 to 7 day intervals may be required to achieve maximum control.

**Grasshopper** - Apply foliarly when grasshopper populations reach local established thresholds to prevent crop damage. Correct timing of spray applications to nymphal stages and thorough coverage is critical to achieve optimum control. For best results, add Methylated Seed Oil (MSO) adjuvant at 1 gallon per 100 gallons of spray volume (1% v/v) when eggs have hatched and the majority of the grasshopper population is 2nd – 3rd instar nymphs. Once grasshoppers contact and/or ingest CORAGEN® there will be rapid feeding cessation; full expression of mortality may take a week or more. Do not make more than two sequential applications of CORAGEN® before rotating to another registered insecticide having a different mode-of-action.



**DUPONT™ CORAGEN® RATE**

<b>Crop</b>	<b>Application Method</b>	<b>Target Pest</b>	<b>Lb ai per acre</b>	<b>fluid ounces product per acre</b>	<b>Last Application (Days to Harvest)</b>	<b>REI (Hours)</b>
Oilseed Group: (EPA Crop Group 20) except milkweed including: Borage; calendula; canola; castor oil plant; Chinese tallowtree; cottonseed; crambe; cuphea; euphorbia; evening primrose; flax seed; gold of pleasure; hare's ear mustard; jojoba; lesquerella; lunaria; meadowfoam; mustard seed; niger seed; oil radish; poppy seed; rapeseed; rose hip; safflower; sesame; stokes aster; sunflower; sweet rocket; tallowwood; tea oil plant; vernonia; cultivars, varieties, and/or hybrids of these	FOLIAR	Diamondback moth Banded sunflower moth Sunflower moth	0.045 - 0.098	3.5 - 7.5	1	4
		Grasshoppers	0.026 - 0.065	2.0 - 5.0		

**USE RESTRICTIONS**  
 Make no more than 4 applications per acre per crop.  
 Minimum interval between treatments is 5 days.  
 Do not apply more than 15.4 fl oz CORAGEN® or 0.2 lb a.i. of chlorantraniliprole containing products per acre per calendar year.  
 CORAGEN® may be applied by overhead sprinkler chemigation systems. See "CHEMIGATION USING OVERHEAD SPRINKLER SYSTEMS - CEREAL GRAINS, CORN (FIELD, POP, SEED) COTTON GRASS FORAGE, FODDER, and HAY, LEGUMES, MINT (PEPPERMINT AND SPEARMINT), NON-GRASS ANIMAL FEEDS, OILSEED GROUP, PEANUT, POTATO, SOYBEAN, AND SUGARCANE" section for instructions on overhead sprinkler chemigation.  
**Banded sunflower moth and sunflower moth** - For best results apply when moth populations reach local established treatment thresholds and as blooms begin to open (sunflower growth stage R-5.0 to R-5.1) to prevent crop damage. Applications may be required at 5-7 day intervals when moth pressure is heavy.  
**Grasshopper** - Apply foliarly when grasshopper populations reach local established thresholds to prevent crop damage. Correct timing of spray applications to nymphal stages and thorough coverage is critical to achieve optimum control. For best results, add Methylated Seed Oil (MSO) adjuvant at 1 gallon per 100 gallons of spray volume (1% v/v) when eggs have hatched and the majority of the grasshopper population is 2nd – 3rd instar nymphs. Once grasshoppers contact and/or ingest CORAGEN® there will be rapid feeding cessation; full expression of mortality may take a week or more. Do not make more than two sequential applications of CORAGEN® before rotating to another registered insecticide having a different mode-of-action.

**DUPONT™ CORAGEN® RATE**

Crop	Application Method	Target Pest	Lb ai per acre	fluid ounces product per acre	Last Application (Days to Harvest)	REI (Hours)
Root and Tuber Vegetables (EPA Crop Group 1), except potato: including Arracacha; arrowroot; artichoke, Chinese; artichoke, Jerusalem; beet, garden; beet, sugar; burdock, edible; canna, edible; carrot; cassava, bitter and sweet; celeriac; chayote (root); chervil, turnip-rooted; chicory; chufa; dasheen (taro); ginger; ginseng; horseradish; leren; parsley, turnip-rooted; parsnip; radish; radish, oriental; rutabaga; salsify; salsify, black; salsify, Spanish; skirret; sweet potato; tanier; turmeric; turnip; yam bean; yam, true.	FOLIAR	Beet armyworm Western yellowstriped armyworm	0.045 - 0.098	3.5 - 7.5	1	4

**USE RESTRICTIONS**  
 Make no more than 4 applications per acre per crop.  
 Minimum interval between treatments is 3 days.  
 Do not apply more than 15.4 fl oz CORAGEN® or 0.2 lb a.i. of chlorantraniliprole containing products per acre per crop.  
 Do not apply more than 61.6 fl oz CORAGEN® or 0.8 lb a.i. of chlorantraniliprole containing products per acre per calendar year; in NY do not apply more than 15.4 fl oz CORAGEN® or 0.2 lb a.i. of chlorantraniliprole containing products per acre per calendar year.

Crop	Application Method	Target Pest	Lb ai per acre	fluid ounces product per acre	Last Application (Days to Harvest)	REI (Hours)
Leaves of Root and Tuber Vegetables (EPA Crop Group 2) (Human Food or Animal Feed) Including: Beet, garden; beet, sugar; burdock, edible; carrot; cassava, bitter and sweet; celeriac; chervil, turnip-rooted; chicory; dasheen (taro); parsnip; radish; radish, oriental (daikon); rutabaga; salsify, black; sweet potato; tanier; turnip; yam, true	FOLIAR	Beet armyworm Western yellowstriped armyworm	0.045 - 0.098	3.5 - 7.5	1	4

**USE RESTRICTIONS**  
 Make no more than 4 applications per acre per crop.  
 Minimum interval between treatments is 3 days.  
 Do not apply more than 15.4 fl oz CORAGEN® or 0.2 lb a.i. of chlorantraniliprole containing products per acre per crop.  
 Do not apply more than 61.6 fl oz CORAGEN® or 0.8 lb a.i. of chlorantraniliprole containing products per acre per calendar year; in NY do not apply more than 15.4 fl oz CORAGEN® or 0.2 lb a.i. of chlorantraniliprole containing products per acre per calendar year.

Crop	Application Method	Target Pest	Lb ai per acre	fluid ounces product per acre	Last Application (Days to Harvest)	REI (Hours)
Rice California only	SOIL APPLICATION† Broadcast spray prior to seeding	Rice water weevil larvae	0.08 - 0.1	6.1 - 7.6		4

**USE RESTRICTIONS**  
 † Broadcast application prior to seeding - apply to the soil prior to seeding and prior to flooding in wet-sown rice culture.  
 Do not apply more than 5 days prior to flooding.  
 Once flood is established, hold the water for a minimum of 14 days before discharging the water.  
 Broadcast application may be made using aerial or ground application equipment.  
 Do not apply more than 7.6 fl oz CORAGEN® or 0.1 lb a.i. of chlorantraniliprole containing products per acre per calendar year.  
 Do not use CORAGEN® treated rice fields for the aquaculture of edible fish or crustacea (including crawfish) during the rice production cycle (planting through harvest).

**DUPONT™ CORAGEN® RATE**

<b>Crop</b>	<b>Application Method</b>	<b>Target Pest</b>	<b>Lb ai per acre</b>	<b>fluid ounces product per acre</b>	<b>Last Application (Days to Harvest)</b>	<b>REI (Hours)</b>
Spice (EPA Crop Subgroup 19B) Including: Allspice; anise (seed); anise, star; annatto (seed); caper (buds); caraway; caraway, black; cardamom; cassia (bark); cassia (buds); celery (seed); cinnamon; clove (buds); coriander (seed); culantro (seed); dill (seed); fennel, common; fennel, Florence (seed); fenugreek; grains of paradise; juniper (berry); lovage (seed); mace; mustard (seed); nutmeg; pepper, black; pepper, white; poppy (seed); saffron; and vanilla	FOLIAR	Beet armyworm Cabbage looper Corn earworm Fall armyworm Southern armyworm	0.045 - 0.065	3.5 - 5.0	1	4

**USE RESTRICTIONS**  
 Make no more than 4 applications per acre per crop.  
 Minimum interval between treatments is 3 days.  
 Do not apply more than 15.4 fl oz CORAGEN® or 0.2 lb a.i. of chlorantraniliprole containing products per acre per crop.  
 Do not apply more than 61.6 fl oz CORAGEN® or 0.8 lb a.i. of chlorantraniliprole containing products per acre per calendar year; in NY do not apply more than 15.4 fl oz CORAGEN® or 0.2 lb a.i. of chlorantraniliprole containing products per acre per calendar year.

**PLANT TOLERANCE PHYTOTOXICITY**  
 CORAGEN® has been tested on numerous crops and cultivars with no observable phytotoxicity at label rates. However, neither the manufacturer nor the seller has determined whether or not CORAGEN® can be used safely on all herbs and spices for which it is registered for use.  
 Since all herbs and spices and their varieties and cultivars have not been tested for phytotoxicity it is recommended that a small number of plants be sprayed initially to determine if there is any phytotoxicity prior to large scale applications to herbs and spices. The user assumes all risks arising from application of CORAGEN® in a manner that is inconsistent with its labeling.

<b>Crop</b>	<b>Application Method</b>	<b>Target Pest</b>	<b>Lb ai per acre</b>	<b>fluid ounces product per acre</b>	<b>Last Application (Days to Harvest)</b>	<b>REI (Hours)</b>
Soybean Including edamame (immature soybean)	FOLIAR	Corn earworm Beet armyworm Fall armyworm Cabbage looper Green cloverworm Southern armyworm Soybean looper Tobacco budworm Velvetbean caterpillar	0.045 - 0.098	3.5 - 7.5	1	4
		Grasshoppers	0.026 - 0.065	2.0 - 5.0		

**USE RESTRICTIONS**  
 Make no more than 4 applications per acre per crop.  
 Minimum interval between treatments is 3 days.  
 Do not apply more than 15.4 fl oz CORAGEN® or 0.2 lb a.i. of chlorantraniliprole containing products per acre per calendar year.  
 CORAGEN® may be applied by overhead sprinkler chemigation systems. See "CHEMIGATION USING OVERHEAD SPRINKLER SYSTEMS - CEREAL GRAINS, CORN (FIELD, POP, SEED) COTTON GRASS FORAGE, FODDER, and HAY, LEGUMES, MINT (PEPPERMINT AND SPEARMINT), NON-GRASS ANIMAL FEEDS, OILSEED GROUP, PEANUT, POTATO, SOYBEAN, AND SUGARCANE" section for instructions on overhead sprinkler chemigation.

**Grasshopper** - Apply foliarly when grasshopper populations reach local established thresholds to prevent crop damage. Correct timing of spray applications to nymphal stages and thorough coverage is critical to achieve optimum control. For best results, add Methylated Seed Oil (MSO) adjuvant at 1 gallon per 100 gallons of spray volume (1% v/v) when eggs have hatched and the majority of the grasshopper population is 2nd – 3rd instar nymphs. Once grasshoppers contact and/or ingest CORAGEN® there will be rapid feeding cessation; full expression of mortality may take a week or more. Do not make more than two sequential applications of CORAGEN® before rotating to another registered insecticide having a different mode-of-action.

**DUPONT™ CORAGEN® RATE**

Crop	Application Method	Target Pest	Lb ai per acre	fluid ounces product per acre	Last Application (Days to Harvest)	REI (Hours)
Strawberry	FOLIAR	Beet armyworm Cabbage looper Corn earworm Japanese beetle (adult) Light brown apple moth	0.045 - 0.098	3.5 - 7.5	1	4

**USE RESTRICTIONS**

Make no more than 4 applications per acre per crop.  
 Minimum interval between treatments is 7 days.  
 Do not apply more than 15.4 fl oz CORAGEN® or 0.2 lb a.i. of chlorantraniliprole containing products per acre per crop.  
 Do not apply more than 30.8 fl oz CORAGEN® or 0.4 lb a.i. of chlorantraniliprole containing products per acre per calendar year; in NY do not apply more than 15.4 fl oz CORAGEN® or 0.2 lb a.i. of chlorantraniliprole containing products per acre per calendar year.

Crop	Application Method	Target Pest	Lb ai per acre	fluid ounces product per acre	Last Application (Days to Harvest)	REI (Hours)
Sugarcane	FOLIAR	Sugarcane borer	0.045 - 0.098	3.5 - 7.5	14	4
		Grasshoppers	0.026 - 0.065	2.0 - 5.0		

**USE RESTRICTIONS**

Make no more than 4 applications per acre per crop.  
 Minimum interval between treatments is 7 days.  
 Do not apply more than 15.4 fl oz CORAGEN® or 0.2 lb a.i. of chlorantraniliprole containing products per acre per calendar year.  
 CORAGEN® may be applied by overhead sprinkler chemigation systems. See "CHEMIGATION USING OVERHEAD SPRINKLER SYSTEMS - CEREAL GRAINS, CORN (FIELD, POP, SEED) COTTON GRASS FORAGE, FODDER, and HAY, LEGUMES, MINT (PEPPERMINT AND SPEARMINT), NON-GRASS ANIMAL FEEDS, OILSEED GROUP, PEANUT, POTATO, SOYBEAN, AND SUGARCANE" section for instructions on overhead sprinkler chemigation.  
**Grasshopper** - Apply foliarly when grasshopper populations reach local established thresholds to prevent crop damage. Correct timing of spray applications to nymphal stages and thorough coverage is critical to achieve optimum control. For best results, add Methylated Seed Oil (MSO) adjuvant at 1 gallon per 100 gallons of spray volume (1% v/v) when eggs have hatched and the majority of the grasshopper population is 2nd - 3rd instar nymphs. Once grasshoppers contact and/or ingest CORAGEN® there will be rapid feeding cessation; full expression of mortality may take a week or more. Do not make more than two sequential applications of CORAGEN® before rotating to another registered insecticide having a different mode-of-action.

Crop	Application Method	Target Pest	Lb ai per acre	fluid ounces product per acre	Last Application (Days to Harvest)	REI (Hours)
Tobacco (except California)	FOLIAR	Split worm (potato tuberworm) Tobacco budworm Tomato hornworm Tobacco hornworm	0.045 - 0.098	3.5 - 7.5	1	4
	SOIL AT PLANTING† (transplant water treatment only)	Tobacco budworm Tomato hornworm Tobacco hornworm	0.065 - 0.098	5.0 - 7.5		

**USE RESTRICTIONS**

Make no more than 4 applications per acre per crop.  
 Minimum interval between treatments is 3 days.  
 Do not apply more than 15.4 fl oz CORAGEN® or 0.2 lb a.i. of chlorantraniliprole containing products per acre per calendar year.  
 †SOIL APPLICATIONS (transplant water treatment at planting): CORAGEN® must be applied uniformly in the root zone or poor performance will result.  
 Do not apply more than 7.5 fl oz (0.098 lb ai per acre) of CORAGEN® to the soil at planting.  
 Refer to the SOIL APPLICATION section of this label for additional guidance.

**CORAGEN® Conversion Chart for Drip (Trickle) Chemigation and At-Plant Soil Application**

Fl oz/acre	Rate in Fluid Ounces Product / 1000 Row-Foot Based on Planted Row Spacing (in inches) of:															
	15	20	25	30	34	36	38	40	44	48	60	66	72	78	80	84
2											0.23	0.25	0.28	0.30	0.30	0.32
3.5				0.20	0.23	0.24	0.25	0.27	0.30	0.32	0.40	0.44	0.48	0.52	0.53	0.56
5		0.19	0.24	0.29	0.33	0.34	0.36	0.38	0.42	0.46	0.57	0.63	0.69	0.75	0.76	0.80
6		0.23	0.29	0.34	0.39	0.41	0.44	0.46	0.50	0.55	0.69	0.76	0.83	0.90	0.91	0.96
7	0.20	0.27	0.33	0.40	0.46	0.48	0.51	0.53	0.59	0.64	0.80	0.88	0.96	1.04	1.07	1.13
7.5	0.22	0.29	0.36	0.43	0.49	0.52	0.55	0.57	0.63	0.69	0.86	0.95	1.03	1.12	1.15	1.21

Level and length of control is affected by rate applied.

Higher labeled rates may be required in heavy texture and/or high organic soils if application is made later in the crop development, or when pest pressure is high.

## **STORAGE AND DISPOSAL**

Do not contaminate water, food, or feed by storage or disposal.

**PESTICIDE STORAGE:** Do not subject to temperatures below 32 degrees F. Store product in original container only in a location inaccessible to children and pets. Do not contaminate water, other pesticides, fertilizer, food or feed in storage. Not for use or storage in or around the home.

**PESTICIDE DISPOSAL:** Do not contaminate water, food or feed by storage or disposal. Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

**CONTAINER HANDLING: Refer to the Net Contents section of this product's labeling for the applicable "Refillable Container" or "Nonrefillable Container" designation.**

**For Small (Capacity Equal to or Less Than 5 Gallons) Nonrefillable Plastic Containers:** Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

**For Large (Capacity Greater Than 5 Gallons) Nonrefillable Plastic Containers:** Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

**For All Refillable Containers:** Refillable container. Refill this container with chlorantraniliprole only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Prior to refilling, inspect carefully for damage such as cracks, punctures, abrasions, worn out threads and closure devices. Check for leaks after refilling and before transporting. Do not transport if this container is damaged or leaking. If the container is damaged, leaking or obsolete, or in the event of a major spill, fire or other emergency, contact DuPont at 1-800-441-3637, day or night.

**NOTICE TO BUYER**— Purchase of this material does not confer any rights under patents of countries outside of the United States.

The DuPont Oval Logo, DuPont™, CORAGEN® and RYNAXYPYR® are trademarks or registered trademarks of E.I. duPont de Nemours and Company.

**D - 1908MSTR 120215-3**

## **LIMITATION OF WARRANTY AND LIABILITY**

Read this Limitation of Warranty and Liability before buying or using this Product. DuPont will not be responsible for losses or damages resulting from use of this product in any manner not specifically directed by DuPont. User assumes all risks associated with such non-directed use. If the terms are not acceptable; return the product at once; unopened and the purchase price will be refunded.

It is impossible to eliminate all risks associated with the use of this product. Such risks arise from weather conditions, soil factors, off target movement, unconventional farming techniques, presence of other materials, the manner of use or application, or other unknown factors, all of which are beyond the control of DuPont. These risks can cause: ineffectiveness of the product, crop injury, or injury to non-target crops or plants. **WHEN YOU BUY OR USE THIS PRODUCT, YOU AGREE TO ACCEPT THESE RISKS.**

DuPont warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for the purpose stated in the Directions for Use, subject to the inherent risks described above, when used in accordance with the Directions for Use under normal conditions.

**TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, DUPONT MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR OF MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL DUPONT OR SELLER BE LIABLE FOR ANY INCIDENTAL, CONSEQUENTIAL OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. BUYER'S OR USER'S BARGAINED-FOR EXPECTATION IS CROP PROTECTION. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER AND THE EXCLUSIVE LIABILITY OF DUPONT OR SELLER, FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY OR CONTRACT, NEGLIGENCE, TORT OR STRICT LIABILITY), WHETHER FROM FAILURE TO PERFORM OR INJURY TO CROPS OR OTHER PLANTS, AND RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT, OR AT THE ELECTION OF DUPONT OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

To the extent consistent with applicable law that allows such requirement, DuPont or its Ag Retailer must have prompt notice of any claim so that an immediate inspection of buyer's or user's growing crops can be made. Buyer and all users shall promptly notify DuPont or a DuPont Ag Retailer of any claims, whether based on contract, negligence, strict liability, other tort or otherwise, or be barred from any remedy.

This Limitation of Warranty and Liability may not be amended by any oral or written agreement.


**For product information call: 1-888-6-DUPONT [1-888-638-7668]**

**© 2008-2015 E. I. du Pont de Nemours and Company, Chestnut Run Plaza, 974 Centre Rd., Wilmington, Delaware 19805.**

**All rights reserved.**



**DuPont Crop  
Protection**



**SUPPLEMENTAL LABELING**

**DUPONT™ CORAGEN®  
INSECT CONTROL WITH ACTIVE  
INGREDIENT RYNAXYPYR®  
FOR USE ON RICE  
IN THE STATE OF CALIFORNIA**

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF CALIFORNIA

**DUPONT™ CORAGEN® INSECT CONTROL**

EPA Reg. No. 352-729

This supplemental label expires on 12/2/18 and must not be used or distributed after this date.

**FOR CONTROL OF RICE WATER WEEVIL IN RICE  
IN THE STATE OF CALIFORNIA**

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Dupont™ CORAGEN® insect control is recommended as a pre-flood, pre-plant broadcast soil application for control of rice water weevil in California rice.

**GENERAL INFORMATION**

CORAGEN™ is a suspension concentration that can be applied as a pre-flood, pre-plant broadcast soil application for control of rice water weevil in California rice. CORAGEN™ is mixed with water for application.

Crop	Application Method	Target Pest	Lb ai per acre	fluid ounces product per acre	Last Application (Days to Harvest)	REI (Hours)
Rice California only	SOIL APPLICATION† Broadcast spray prior to seeding	Rice water weevil larvae	0.08 - 0.1	6.1 - 7.6		4

**USE RESTRICTIONS**

† Broadcast application prior to seeding - apply to the soil prior to seeding and prior to flooding in wet-sown rice culture.  
Do not apply more than 5 days prior to flooding.  
Once flood is established, hold the water for a minimum of 14 days before discharging the water.  
Broadcast application may be made using aerial or ground application equipment.  
Do not apply more than 7.6 fl oz CORAGEN® or 0.1 lb a.i. of chlorantraniliprole containing products per acre per calendar year.  
Do not use CORAGEN® treated rice fields for the aquaculture of edible fish or crustacea (including crawfish) during the rice production cycle (planting through harvest).

**SOIL APPLICATIONS**

CORAGEN® must be applied in a manner that ensures the product is in the root zone. CORAGEN™ must be in the root zone to provide effective control of target pests. CORAGEN® does not readily move in the soil; CORAGEN® will typically move approximately 1 – 3 inches from the site of delivery into the soil.

**IMPORTANT**

Before using CORAGEN®, read and follow all applicable directions, restrictions and precautions on the EPA Registered label. This bulletin contains new or supplemental instructions for use of this product, which do not appear on the EPA-registered package label. Follow the instructions carefully. This labeling must be in the possession of the user at the time of pesticide application.

Read the Limitations of Warranty and Liability on the Section 3 Federal product label before buying or using this product. If terms are not acceptable, return the unopened package at once to the Seller for full refund of purchase price paid. Otherwise, use by the Buyer or any other User constitutes acceptance of the terms of the Limitation of Warranty and Liability on the Section 3 Federal product label.

For product information call 1-888-6-DUPONT

© 2015 E.I. du Pont de Nemours and Company, Chestnut Run Plaza, 974 Centre Road, Wilmington, DE 19805 U.S.A.