



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

June 27, 2019

S. K. Theodorakis
Registration Manager
Corteva Agriscience
Chestnut Run Plaza LR735/4160-8
974 Centre Road
Wilmington, DE 19805

Subject: Registration Review Label Mitigation for Nicosulfuron
Product Name: DuPont Accent SC Herbicide
Application Date: 12/14/2017
EPA Registration Number: 352-747
Decision Number: 541945

Dear Mr. Theodorakis:

The Agency, in accordance with the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA), as amended, has completed reviewing all of the information submitted with your application to support the Registration Review of the above referenced product in connection with the 22 Sulfonylurea (SU) Herbicides Interim Decision, and has concluded that your submission is acceptable. The label referred to above, submitted in connection with registration under FIFRA, as amended, is acceptable.


Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

A copy of your label stamped "Accepted" is enclosed. Products shipped after 12 months from the date of this amendment must bear the new revised label. Your release for shipment of the product bearing the amended label constitutes acceptance of these conditions. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6.

Page 2 of 2
EPA Reg. No. 352-747
Decision No. 541945

If you have any questions about this letter, please contact Erik Kraft by phone at 703-308-9358, or via email at kraft.erik@epa.gov.

Sincerely,

A handwritten signature in black ink, appearing to read "Erik Kraft". The signature is fluid and cursive, with the first name "Erik" being more prominent than the last name "Kraft".

Erik Kraft, Product Manager 24
Fungicide and Herbicide Branch
Registration Division (7505P)
Office of Pesticide Programs

Enclosure

DuPont™ ACCENT® SC**HERBICIDE**

GROUP

2

HERBICIDE

***For use on Field Corn grown for grain, silage, or seed, Popcorn and Sweet Corn.
For use on Grain Sorghum containing INZEN™ Trait.***

Soluble Concentrate

<i>Active Ingredients</i>	By Weight
Nicosulfuron	
2-[[[(4,6-dimethoxypyrimidin-2-yl)aminocarbonyl]aminosulfonyl]-N,N-dimethyl-3-pyridinecarboxamide	4.2%
<i>Other Ingredients</i>	95.8%
TOTAL	100.0%

ACCENT® SC contains 0.33 lbs. of active ingredient per gallon.

®™ Trademarks of Dow AgroSciences, DuPont or Pioneer and their affiliated companies or respective owners

EPA Reg. No. 352-747

EPA Est. No. _____

[Nonrefillable Container]

[Refillable Container]

Net: _____

ACCEPTED**06/27/2019**

Under the Federal Insecticide, Fungicide
and Rodenticide Act as amended, for the
pesticide registered under
EPA Reg. No.

352-747**KEEP OUT OF REACH OF CHILDREN****CAUTION**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (if you do not understand this label, find someone to explain it to you in detail.)

FIRST AID

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-441-3637 for emergency medical treatment information.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Avoid contact with skin or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear:

Long-sleeved shirt and long pants
Chemical resistant gloves made of any water proof material including butyl rubber, natural rubber, neoprene rubber, or nitrile rubber
Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

USERS SHOULD: Users should remove clothing/PPE immediately if pesticide gets inside. Then wash through and put on clean clothing. Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters or rinsate. Do not apply where/when conditions could favor runoff.

Ground Water Advisory

Nicosulfuron is known to leach through soil into groundwater under certain conditions as a result of label use. This chemical may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow.

Surface Water Advisory

This product may impact surface water quality due to runoff of rain water. This is especially true for poorly draining soils and soils with shallow ground water. This product is classified as having high potential for reaching surface water via runoff for several months or more after application. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of this product from runoff water and sediment. Runoff of this product will be greatly reduced by avoiding applications when rainfall or irrigation is expected to occur within 48 hours.

Windblown Soil Particles Advisory

This product has the potential to move off-site due to wind erosion. Soils that are subject to wind erosion usually have a high silt and/or fine to very fine sand fractions and low organic matter content. Other

factors which can affect the movement of windblown soil include the intensity and direction of prevailing winds, vegetative cover, site slope, rainfall, and drainage patterns. Avoid applying this product if prevailing local conditions may be expected to result in off-site movement.

Non-target organism Advisory

This product is toxic to plants and may adversely impact the forage and habitat of non-target organisms, including pollinators, in areas adjacent to the treated area. Protect the forage and habitat of non-target organisms by minimizing spray drift. For further guidance and instructions on how to minimize spray drift, refer to the Spray drift Management section of this label.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 4 hours. PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, including plants, soil, or water, is:

Coveralls

Chemical resistant gloves made of any water proof material such as butyl rubber, natural rubber, neoprene rubber, or nitrile rubber

Shoes plus socks

ACCENT® SC herbicide must be used only in accordance with directions on this label. To the extent consistent with the law, DuPont will not be responsible for losses or damage resulting from use of this product in any manner not specifically directed by DuPont.

PRODUCT INFORMATION

ACCENT® SC herbicide is a suspension concentrate used at a rate 6 - 24 fluid ounces (0.015 - 0.062 lb ai) per acre for selective postemergence grass weed control in field corn grown for seed or grain, popcorn and sweet corn, and 12 - 24 fluid ounces (0.031 - 0.062 lb ai) per acre in grain sorghum containing the INZEN™ trait.

Always shake well before use.

RESTRICTIONS

Aerial application is not permitted in New York and California.

Do not apply ACCENT® SC to corn or grain sorghum containing the INZEN™ trait that exhibits herbicide injury from previous applications made to the current or preceding crop.

Do not use liquid nitrogen fertilizer as the total carrier solution for postemergence applications.

Injury or loss of desirable trees or vegetation may result from failure to observe the following:

- Do not apply ACCENT® SC or drain or flush application equipment on or near desirable trees or other plants, or on areas where their roots may extend, or in locations where the chemical may be washed or moved into contact with their roots.
- Do not use on lawns, walks, driveways, or tennis courts. Prevent drift of spray to desirable plants.
- Do not contaminate any body of water.

Do not apply this product through any type of irrigation system.

PRECAUTIONS

ACCENT® SC may interact with certain insecticides previously applied to the crop. Crop response varies with field corn type, insecticide used, insecticide application methods, and soil type.

ACCENT® SC may be applied to corn previously treated with “Fortress” (AI: chlorethoxyfos Reg. No. 59807-19), “Smartchoice” (AI: bifenthrin + chlorethoxyfos: Reg. No. 5481-561, 5481-579), “Aztec” (AI: cyfluthrin + phostebupirim; Reg. Nos 5481-577, 5481-9028, 5481-9030), or “Force” (AI: tefluthrin; Reg. Nos, 100-1075, 100-1253, 100-1625) insecticides, or other non-organophosphate soil insecticides regardless of soil type.

It is possible that pollen-mediated gene flow from grain sorghum containing the INZEN™ trait to weedy relatives, including johnsongrass or shattercane, may contribute to the development of resistance to ALS herbicides in these biotypes.

Temporary yellowing and reduction in height of grain sorghum hybrids containing the INZEN™ trait may occur following a postemergence application of ACCENT® SC. Crop responses may be more pronounced when conditions exist that result in slowed crop growth, including but not limited to, cloudy, cool, or wet conditions. Normal growth and appearance will resume when normal growing conditions return.

Adherence to the DuPont Stewardship program and Best Management Practices is necessary to reduce the risk of the development of resistance to ALS herbicides in weedy relatives.

Crop injury may occur following an application of ACCENT® SC if there is a prolonged period of cold weather and/or in conjunction with wet soils.

Prevent drift or spray onto desirable plants.

Thoroughly clean application equipment immediately after use (See Sprayer cleanup section of this label).

For all application systems, use 50-mesh or larger strainer screens.

WEED RESISTANCE MANAGEMENT

ACCENT® SC, which contains the active ingredient nicosulfuron is a group 2 herbicide based on the mode of action classification system of the Weed Science Society of America.

Proactively implementing diversified weed control strategies to minimize selection for weed populations resistant to one or more herbicides is a best practice. A diversified weed management program may include the use of multiple herbicides with different sites of action and overlapping weed spectrum with or without tillage operations and/or other cultural practices. Research has demonstrated that using the

labeled rate and directions for use is important to delay the selection for resistance.

The continued effectiveness of this product depends on the successful implementation of a weed resistance management program.

To aid in the prevention of developing weeds resistant to this product, users should:

- Scout fields before application to ensure herbicides and rates will be appropriate for the weed species and weed sizes present.
- Start with a clean field, using either a burndown herbicide application or tillage.
- Control weeds early when they are relatively small (less than 4 inches).
- Apply full rates of this product for the most difficult to control weed in the field at the specified time (correct weed size) to minimize weed escapes.
- Scout fields after application to detect weed escapes or shifts in control of weed species
- Control weed escapes before they reproduce by seed or proliferate vegetatively.
- Report any incidence of non-performance of this product against a particular weed to your DuPont representative, local retailer, or county extension agent.
- Contact your DuPont representative, crop advisor, or extension agent to find out if suspected resistant weeds to this MOA have been found in your region. If resistant biotypes of target weeds have been reported, use the application rates of this product specified for your local conditions. Tank mix products so that there are multiple effective sites of actions for each target weed.
- If resistance is suspected, treat weed escapes with an herbicide having a site of action other than group 2 and/or use nonchemical methods to remove escapes, as practical, with the goal of preventing further seed production.
- Suspected herbicide-resistant weeds may be identified by these indicators:
 - Failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds;
 - A spreading patch of non-controlled plants of a particular weed species; and
 - Surviving plants mixed with controlled individuals of the same species.

Additionally, users should follow as many of the following herbicide resistance management practices as is practical:

- Use a broad spectrum soil-applied herbicide with other sites of action as a foundation in a weed control program.
- Utilize sequential applications of herbicides with alternative sites of action.
- Rotate the use of this product with non-group 2 herbicides.
- Avoid making more than two applications of this product and any other group 2 herbicides within a year unless mixed with an herbicide with a different site of action with an overlapping spectrum for the difficult-to- control weeds.

- Incorporate non-chemical weed control practices, such as mechanical cultivation, crop rotation, cover crops and weed-free crop seeds, as part of an integrated weed control program.
- Use good agronomic principles that enhance crop development and crop competitiveness.
- Thoroughly clean plant residues from equipment before leaving fields suspected to contain resistant weeds.
- Manage weeds in and around fields, during and after harvest to reduce weed seed production.

INTEGRATED PEST MANAGEMENT

This product may be used as part of an Integrated Pest Management (IPM) program that can include biological, cultural, and genetic practices aimed at preventing economic pest damage. IPM principles and practices include field scouting or other detection methods, correct target pest identification, population monitoring, and treating when target pest populations reach locally determined action thresholds. Consult your state cooperative extension service, professional consultant or other qualified authorities to determine appropriate action treatment threshold levels for treating specific pest / crop systems in your area.

MANDATORY SPRAY DRIFT MANAGEMENT

Aerial Applications:

- Do not release spray at a height greater than 10 feet above the vegetative canopy, unless a greater application height is necessary for pilot safety.
- For applications prior to the emergence of crops and target weeds, applicators are required to use a coarse or coarser droplet size (ASABE S572.1)
- For all other applications, applicators are required to use a Medium or coarser droplet size (ASABE S572.1).
- The boom length must not exceed 65% of the wingspan for airplanes or 75% of the rotor blade diameter for helicopters.
- Applicators must use one-half swath displacement upwind at the downwind edge of the field.
- Nozzles must be oriented so the spray is directed toward the back of the aircraft.
- Do not apply when wind speeds exceed 10 miles per hour at the application site.
- Do not apply during temperature inversions.

Ground boom Applications:

- Apply with the nozzle height recommended by the manufacturer, but no more than 3 feet above the ground or crop canopy unless making a turf, pasture, or rangeland application, in which case applicators may apply with a nozzle height no more than 4 feet above the ground.
- For applications prior to the emergence of crops and target weeds, applicators are required to use a coarse or coarser droplet size (ASABE S572.1).
- For all other applications, applicators are required to use a Medium or coarser droplet size (ASABE S572.1).
- Do not apply when wind speeds exceed 10 miles per hour at the application site.
- Do not apply during temperature inversions.

SPRAY DRIFT ADVISORIES

THE APPLICATOR IS RESPONSIBLE FOR AVOIDING OFF-SITE SPRAY DRIFT. BE AWARE OF NEARBY NON-TARGET SITES AND ENVIRONMENTAL CONDITIONS.

IMPORTANCE OF DROPLET SIZE

An effective way to reduce drift potential is to apply large droplets. Use the largest droplets that provide target pest control. While applying larger droplets will reduce spray drift, the potential for drift will be greater if applications are made improperly or under unfavorable environmental conditions.

Controlling Droplet Size – Ground Boom

- **Volume** - Increasing the spray volume so that larger droplets are produced will reduce spray drift. Use

the highest practical spray volume for the application. If a greater spray volume is needed, consider using a nozzle with a higher flow rate.

- **Pressure** - Use the lower spray pressures recommended for the nozzle to produce the target spray volume and droplet size.
- **Spray Nozzle**- Use a spray nozzle that is designed for the intended application. Consider using nozzles designed to reduce drift.

Controlling Droplet Size – Aircraft

- **Adjust Nozzles** - Follow nozzle manufacturers recommendations for setting up nozzles. Generally, to reduce fine droplets, nozzles should be oriented parallel with the airflow in flight.

BOOM HEIGHT – Ground Boom

Use the lowest boom height that is compatible with the spray nozzles that will provide uniform coverage. For ground equipment, the boom should remain level with the crop and have minimal bounce.

RELEASE HEIGHT – Aircraft

Use the lowest boom height that is compatible with the spray nozzles that will provide uniform coverage. For ground equipment, the boom should remain level with the crop and have minimal bounce.

SHIELDED SPRAYERS

Shielding the boom or individual nozzles can reduce spray drift. Verify that the shields are not interfering with the uniform deposition of the spray on the target area.

TEMPERATURE AND HUMIDITY

When making applications in hot and dry conditions, use larger droplets to reduce effects of evaporation.

TEMPERATURE INVERSIONS

Drift potential is high during a temperature inversion. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. The presence of an inversion can be indicated by ground fog or by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing. Avoid applications during temperature inversions.

WIND

Drift potential generally increases with wind speed. **AVOID APPLICATIONS DURING GUSTY WIND CONDITIONS.** Applicators need to be familiar with local wind patterns and terrain that could affect spray drift.

DRIFT CONTROL ADDITIVES

Using product compatible drift control additives can reduce drift potential. When a drift control additive is used, read and carefully observe cautionary statements and all other information on the additive's label. If using an additive that increases viscosity, ensure that the nozzles and other application equipment will function properly with a viscous spray solution. Preferred drift control additives have been certified by the Chemical Producers and Distributors Association (CPDA).

CORN**RESTRICTIONS**

CROPS	Maximum Fl Oz of Product/ Acre/ Single Application	Maximum Lb AI/Acre/Single Application	Maximum Number of Applications per Year	Maximum FL Oz of Product/Acre/Year	Maximum Lb AI/A per Year	Retreat Interval (Days)	Last Treatment Preharvest Interval
Field Corn – Grown for Grain or Silage; Seed, Popcorn, or Sweet Corn	24 fl oz	0.062 lb ai	2	24 fl oz	0.062 lb ai	7	Do not harvest corn grain within 70 days or harvest corn forage or stover within 45 days of an ACCENT® SC application

The combined dosage of sequential applications cannot exceed 24 fluid ounces (0.063 pounds of nicosulfuron active ingredient) per acre of ACCENT® SC in corn.

When tank mixing with other products that contain nicosulfuron, do not apply more than 1.0 ounce (0.062 lb) of nicosulfuron in corn per year.

Do not tank mix ACCENT® SC with products containing bentazon or severe crop injury may occur.

Do not tank mix ACCENT® SC with foliar- applied organophosphate insecticides including chlorpyrifos, malathion, etc., as severe crop injury may occur. To avoid crop injury or antagonism, apply these products at least 7 days before or 3 days after the application of ACCENT® SC.

Do not apply ACCENT® SC within 45 days of crop emergence where the organophosphate insecticide, terbufos (including "Counter" brands) was applied since corn crop injury may occur. Applications made to corn previously treated with chlorpyrifos or other similar organophosphate insecticides may result in unacceptable crop injury. Any crop injury or yield loss resulting from these applications are the responsibility of the grower.

Do not make a late application of ACCENT® SC to field corn grown for grain or silage that is taller than 36 inches or that exhibits 10 or more collars (V10), whichever is more restrictive.

Do not apply ACCENT® SC to any white popcorn inbred, or white popcorn hybrid.

Do not apply ACCENT® SC to sweet corn taller than 18 inches or which exhibits 6 or more leaf collars (V6).

Do not graze, feed forage, grain or fodder (stover) from corn treated areas to livestock within 30 days of an ACCENT® SC application.

FIELD CORN GROWN FOR GRAIN OR SILAGE**Timing to Crop**

ACCENT® SC may be used on field corn, high lysine, waxy, white or other food grade corn hybrids.

ACCENT® SC may be broadcast to corn up to 20" tall (free standing) or that is exhibiting up to and including 6 leaf collars (V6), whichever is more restrictive.

While ACCENT® SC has a wide application window, research has shown best results are obtained when applications are made early postemergence when corn and weeds are small. Target applications to corn

that is less than 12" tall for best overall performance.

Timing to Weeds

Apply ACCENT® SC when grasses are young and actively growing, but before they exceed the sizes indicated in the table Weeds controlled with 12 fluid ounces (0.031 lb ai) ACCENT® SC. Treat heavy infestations of weeds before they become too competitive with the crop, especially where soil moisture and/or fertility are limited. ACCENT® SC provides weed control via foliar absorption. ACCENT® SC only controls those weeds that have emerged. For later-emerging weeds, a second application or a timely cultivation is required. Applications made to weeds larger than the size indicated on this label or to weeds under stress may result in unsatisfactory control. Refer to LATE APPLICATIONS.

LATE APPLICATIONS IN FIELD CORN GROWN FOR GRAIN OR SILAGE

ACCENT® SC may be applied to field corn as a directed postemergence application on corn that is taller than 20" or which has more than 6 collars (whichever occurs first).

- For corn 20" to 36" tall, apply ACCENT® SC with drop nozzles only and avoid spraying into the whorl of cornstalks.
- **Do not** apply to corn that is taller than 36" or that exhibits 10 or more collars (V10), whichever is most restrictive.

Applications made to weeds larger than those listed on this label may vary from complete control to suppression. Level of control will depend on the weed species, stage of growth, and environmental conditions.

Choices must be made between the risks that arise from applications made beyond the proper time for ACCENT® SC use, and the effects of season long grass competition and/or harvest complications. These choices must balance risks from improperly timed ACCENT® SC use that include, but are not limited to:

- Yield loss due to competition: Research indicates competition from foxtail exceeding 4 inches in height may reduce corn yields. Applications to foxtail and other annual grasses that exceed the sizes stated on the label increases the risk of yield losses due to prolonged competition with the crop even though control may be acceptable.
- Incomplete control of grasses beyond labeled size: Applications to grasses that exceed the labeled sizes can result in reduced control. This incomplete control may reduce corn yield.
- Incomplete grass control due to herbicide stress: Grasses under stress from previous herbicide applications may not be actively growing and susceptible to ACCENT® SC.
- Ear malformation: Applications of ACCENT® SC on corn that has 7 to 10 collars (V7 to V10) increases the potential for ear malformation (pinching). This risk may be greatly reduced, but not eliminated, by using drop nozzles properly adjusted so as to not apply ACCENT® SC into the corn whorl.

APPLICATION RATE

Optimum control of the weeds listed can be achieved with 12 fluid ounces (0.031 lb ai) of ACCENT® SC. Weeds that exceed the listed weed sizes by up to 50% may be partially controlled with rates between 12 and 24 fluid ounces (0.031 and 0.062 lb ai) of ACCENT® SC per acre.

ACCENT® SC may be applied at 6 - 12 fluid ounces (0.015 – 0.031 lb ai) for limited control of certain small grass weeds. See the table for reduced rates under ADDITIONAL DIRECTIONS for details.

As weeds mature, their sensitivity to ACCENT® SC decreases. As grassy weeds become mature (more than 3 tillers), they may not reach the size listed below, due to drought or other environmental factors.

Treat grassy weeds that are maturing rapidly before they reach the stages listed below.

When applied as directed, ACCENT® SC will control the following weeds:

Weeds controlled with 12 fluid ounces (0.031 lb ai) ACCENT® SC.

Grasses	Maximum Height or Diameter
Barnyardgrass	4"
Broadleaf signalgrass	2"
Foxtails (bristly, giant, green, yellow)	4"
Itchgrass	6"
Johnsongrass [†]	
seedling	12"
rhizome	18"
Panicum (Texas, browntop)	3"
Fall	4"
Quackgrass*	10"
Ryegrass (Italian, perennial)	6"
Sandbur (field, longspine)*	3"
Timothy	6"
Volunteer cereals (barley, oats, rye, triticale, wheat)	6"***
Wild oats	4"
Wild proso millet	4"
Wirestem muhly*	8"
Witchgrass	6"
Woolly Cupgrass* [†]	4"

* Requires the use of COC plus ammonium nitrogen fertilizer. Cultivation or re-treatment may be required. See "For ADDITIONAL CONTROL OF LATER EMERGING GRASSES"

**10 inches in the states of WA, OR, ID, and MT, where the use of MSO adjuvants are preferred. See SPRAY ADJUVANTS.

† Naturally occurring resistant biotypes are known to occur.

Broadleaves	Maximum Height or Diameter
Burcucumber	3"
Dandelion	6"
Hemp dogbane*	4"
Jimsonweed	3"
Morningglory (ivyleaf, pitted)	3"
tall	2"
Pigweed (redroot, smooth)	4"
Pokeweed*	4"
Smartweeds (ladysthump, PA)	4"
Thistle, Canada*	4"

*Suppression

POPCORN, FIELD CORN GROWN FOR SEED AND SWEET CORN

ACCENT® SC may be broadcast or applied with drop nozzles to popcorn or field corn grown for seed that is less than 20" tall (free-standing) or that exhibits up to and including 5 leaf-collars (V5), whichever is most restrictive. **Do not** apply to corn that is taller than 20" or that exhibits more than 5 leaf-collars (V5),

whichever is more restrictive.

Many seed companies have tested seed corn inbreds or yellow popcorn hybrids for sensitivity to ACCENT® SC and have reported excellent safety. **Do not** apply ACCENT® SC to any white popcorn inbred, or white popcorn hybrid unless specifically approved by the seed company. This includes “White Dynamite” popcorn.

ACCENT® SC may be applied to certain sweet corn hybrids grown for fresh markets or under contract for processing. Applications of ACCENT® SC may be applied broadcast or with drop nozzles (post-directed) on sweet corn up to 12 inches tall or up to and including 5 leaf-collars (V5). For sweet corn 12 - 18 inches tall, apply only with drop nozzles. **Do not** apply to sweet corn taller than 18 inches or those which exhibit 6 or more leaf-collars (V6) and make only one application of ACCENT® SC per year.

Sweet corn hybrid sensitivity to ACCENT® SC is highly variable, and not all hybrids have been tested for crop sensitivity. Contact your DuPont Crop Protection Sales Representative for information on local sweet corn hybrids that have been evaluated with ACCENT® SC.

Not all seed corn inbreds, popcorn or sweet corn hybrids have been tested, nor does DuPont have access to all seed company data. Consequently, to the extent consistent with the law, DuPont is not responsible for any crop injury arising from the use of ACCENT® SC on field corn grown for seed, popcorn or sweet corn. When tank mixing, check the tank mix partner label for sensitivities and instructions for use. It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

SEQUENTIAL APPLICATIONS FOLLOWING PREEMERGENCE HERBICIDES

ACCENT® SC is best used in a planned postemergence weed control program as part of a sequential application herbicide program, following a before-planting application of: BASIS® Blend (AI: rimsulfuron + thifensulfuron methyl; Reg. No. 352-854), BREAKFREE® brands (AI: acetochlor + atrazine; Reg. Nos. 352-893, -894 or -895), CINCH® (AI: S-metolachlor; Reg. No. 352-625), Cinch ATZ (AI: S-metolachlor + atrazine; Reg. No. 624), Cinch ATZ Lite (AI S-metolachlor + atrazine; Reg. No. 352-623), INSTIGATE® (AI: mesotrione + rimsulfuron; Reg. No. 352-873), LEADOFF® (AI: rimsulfuron + thifensulfuron methyl; Reg. No. 352-853), PREQUEL® (AI: isoxaflutole + rimsulfuron; Reg. No. 352-779), RESOLVE® Q (AI: rimsulfuron + thifensulfuron methyl; Reg. No. 352-777) herbicides and/or other pre-emergence applied corn herbicides.

Refer to the label of the respective corn herbicide partner for specific use directions.

Restriction: Do not apply ACCENT® SC to corn that exhibits herbicide injury from previous applications made to the current or preceding crop.

TANK MIX APPLICATIONS

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

For Additional Control of Broadleaf Weeds

ACCENT® SC may be tank mixed with herbicides registered for postemergence application in corn for

additional control of broadleaf weeds. It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

See SPRAY ADJUVANTS for adjuvant rate directions.

ADDITIONAL DIRECTIONS AND/OR DIRECTIONS FOR SPECIFIC WEED PROBLEMS

Reduced Rates of ACCENT® SC

ACCENT® SC may be applied at 6 - 12 fluid ounces (0.015 – 0.031 lb ai) for control of the small grass weeds noted in the table below. Always use a crop oil concentrate plus ammonium nitrogen fertilizer when applying reduced rates of ACCENT® SC.

Weeds Controlled With Reduced Rates of ACCENT® SC.

Grasses	Accent® SC Rate		
	6 fl oz (0.015 lb ai)	9 fl oz (0.023 lb ai)	12 fl oz (0.031 lb ai)
Barnyardgrass	2"	3"	4"
Foxtails (bristly, giant, green)	2"	3"	4"
yellow	--	2"	4"
Itchgrass	2"	4"	6"
Johnsongrass [†] , seedling	--	8"	12"
rhizome	--	8"	18"
Panicum (Texas, browntop)	1"	2"	3"
fall	1"	2"	4"
Sandbur (field, longspine)	--	1"	3"
Timothy	2"	4"	6"
Volunteer cereals	--	2"	6"
Wild oats	2"	3"	4"
Wild proso millet	--	2"	4"
Witchgrass	2"	4"	6"
Woolly cupgrass	--	--	4"

[†] Naturally occurring resistant biotypes are known to occur.

Other Tank Mixtures

Other than the exceptions noted, and in addition to the tank mix partners and rates indicated above, ACCENT® SC may be tank mixed or followed with sequential applications of other products registered for use in field corn. Applications of full or reduced rates of other products registered for use in corn provided:

- The tank mix product is labeled for the same timing, method of application, adjuvants, and use restrictions as ACCENT® SC.
- The tank mixture is not specifically prohibited on the label of the tank mix product.
- The tank mix combination is compatible as determined by a "jar test" described in the TANK MIX COMPATIBILITY TESTING section below.

Weed control and crop response with tank mixtures not specifically directed in this label are the responsibility of the user and manufacturer of the tank mix product.

A corn plant's predisposition to develop fused tissue emerging from the whorl (rattail) after the V-11 stage may increase when a product containing dicamba is applied to small corn under early stressful conditions. Be aware of this when applying tank mixes with dicamba to small corn (V-3 stage or smaller) under stressful conditions. See ENVIRONMENTAL CONDITIONS for a description of these stressful conditions.

SEQUENTIAL APPLICATIONS OF ACCENT® SC

Annual grasses may have more than one flush of emerging seedlings. Also, regrowth of treated annual grasses may occur due to adverse environmental conditions following application. Perennial grasses may regrow from underground stems or roots, depending upon environmental conditions. To control grasses under these conditions, a sequential application of ACCENT® SC may be necessary.

CULTIVATION

A timely cultivation may be necessary to control suppressed weeds or weeds that emerge after an application of ACCENT® SC.

Optimum timing for cultivation is 7–14 days after ACCENT® SC application or upon seeing the establishment of new weeds.

GRAZING / PREHARVEST INTERVALS FOR CORN

Restriction: Do not graze or feed forage, hay or straw from treated areas to livestock within 30 days of ACCENT® SC application.

GRAIN SORGHUM CONTAINING INZEN™ TRAIT

PRODUCT INFORMATION

Apply ACCENT® SC herbicide to grain sorghum containing the INZEN™ trait for postemergence control of certain annual and perennial grass and annual broadleaf weeds.

These application directions are specific for ACCENT® SC applied to grain sorghum containing the INZEN™ trait. **Do not** use ACCENT® SC on grain sorghum that does not contain the INZEN™ trait as severe injury or death will occur.

It is possible that pollen-mediated gene flow from grain sorghum containing the INZEN™ trait to weedy relatives, including shattercane and johnsongrass, may contribute to the development of resistance to ALS herbicides in these species. Plant into fields in which emerged weeds have been controlled by tillage or nonselective herbicides, including glyphosate. Manage johnsongrass and shattercane growth in road ditches, fence rows and nearby places so their flowering does not coincide with the INZEN™ sorghum trait flowering. **Do not** use ACCENT® SC on grain sorghum containing the INZEN™ trait in fields known to have ALS-resistant shattercane or johnsongrass. Adherence to the DuPont Stewardship program, including completion of the certification program and following the Best Management Practices is necessary to reduce the risk of the development of resistance to ALS herbicides in weedy relatives.

RESTRICTIONS

CROP	Maximum Fl Oz of Product/ Acre/ Single Application	Maximum Lb AI/Acre/Single Application	Maximum Number of Applications per Year	Maximum FL Oz of Product/Acre/Year	Maximum Lb AI/A per Year	Retreat Interval (Days)	Last Treatment Preharvest Interval
Grain sorghum containing the INZEN™ Trait	24 fl oz	0.062 lb ai	2	32 fl oz	0.084 lb ai	7	Do not apply to grain sorghum taller than 20 inches. Forage may be cut and livestock may be grazed once the crop has reached the mature forage stage (soft dough growth stage 7). Grain and stover may be harvested once the crop has reached the mature grain stage (physiological maturity growth stage 9).

Do not use ACCENT® SC on grain sorghum that does not contain the INZEN™ trait as severe injury or death will occur.

Do not plant grain sorghum containing the INZEN™ trait in fields known to have ALS resistant johnsongrass or shattercane.

Do not plant sorghum the year following growing of grain sorghum containing the INZEN™ trait in the same field.

The combined dosage of sequential applications cannot exceed 32 fluid ounces (0.084 lb ai) per acre in grain sorghum containing the INZEN™ trait per year.

Allow a minimum of 7 days between applications, but do not make any additional ACCENT® SC application until all herbicide symptomatology including yellowing or reduced plant height has subsided on the grain sorghum containing the INZEN™ trait.

When tank mixing with other products that contain nicosulfuron, do not apply more than 1.34 ounces (0.084 lb) of nicosulfuron in grain sorghum containing the INZEN™ trait per year.

Do not tank mix ACCENT® SC with "Huskie" brands (AI: pyrasulfotole + bromoxynil) as significant grass antagonism, and INZEN™ grain sorghum crop injury can result.

Do not use crop oil concentrate (COC) with ACCENT® SC when tank mixing dicamba or 2,4-D, use only non-ionic surfactant (NIS), in grain sorghum containing the INZEN™ trait.

Do not apply dicamba or 2,4-D if the potential for injury to grain sorghum containing the INZEN™ trait is not acceptable.

APPLICATION DIRECTIONS

WEED RESISTANCE IN GRAIN SORGHUM CONTAINING THE INZEN™ TRAIT

The continued availability and utility of this product depends on the successful management of the weed resistance program; therefore, it is very important to perform the following activities.

The following steps are provided to aid in the prevention of developing weeds resistant to this product:

- Scout fields before application to ensure herbicides and rates will be appropriate for the weed species and weed sizes present.
- Apply the maximum specified labeled use rates of ACCENT® SC for the most difficult to control weeds in the field at the specified time (correct weed size) or when applications are made under challenging environmental conditions to minimize weed escapes.
- Scout fields after application to detect weed escapes or shifts in weed species.
- Report any incidence of non-performance of this product on a weed species listed in the “Weeds Controlled” section to your DuPont retailer, representative or 1-800-258-3033.
- If resistance is suspected in a weed species listed in the “Weeds Controlled” section or to johnsongrass or shattercane, treat the weed escapes with an herbicide having a mode of action other than group 2 and/or use non-chemical methods to remove escapes, as practicable, with the goal of preventing further seed production. Report suspected resistance to your DuPont retailer, representative or 1-800-258-3033.
- Suspected herbicide-resistant weeds may be identified by these indicators:
 - Failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds;
 - A spreading patch of non-controlled plants of a particular weed species; and
 - Surviving plants mixed with controlled individuals of the same species.

Additionally, users should follow as many of the following herbicide resistance management practices as practical:

- Use a broad-spectrum soil-applied herbicide with other modes of action as a foundation in a weed control program.
- Utilize sequential applications of herbicides with alternative modes of action.
- Rotate the use of this product with non-group 2 herbicides.
- Incorporate non-chemical weed control practices, including mechanical cultivation, crop rotation, cover crops and weed-free crop seeds, as part of an integrated weed control program.
- Thoroughly clean plant residues from equipment before leaving fields suspected to contain resistant weeds.
- Avoid using any other group 2 herbicide within a single growing season unless in conjunction with another mode of action herbicide with overlapping spectrum.
- Manage weeds in and around fields, during and after harvest to reduce weed seed production.

Contact the local agricultural extension service, DuPont representative, agricultural retailer or crop consultant for further guidance on weed control practices as needed.

APPLICATION RATE

Apply 12 to 24 fluid ounces ACCENT® SC (0.031 to 0.062 lb ai) by ground or by air per acre per application to grain sorghum containing the INZEN™ trait.

APPLICATION TIMING TO CROP

Apply ACCENT® SC to emerged grain sorghum containing the INZEN™ herbicide tolerance trait that is up to 20 inches tall. Applications made to 4-20 inch tall grain sorghum (approximately five leaf stage (growth stage 2) to flag leaf visible (growth stage 4)) are advised for best crop resiliency. **Do not** apply to grain sorghum taller than 20 inches.

APPLICATION TIMING TO WEEDS

Apply ACCENT® SC when grasses are young and actively growing, but before they exceed the sizes indicated in the table WEEDS CONTROLLED IN INZEN™ TRAIT GRAIN SORGHUM with 12 fluid ounces (0.031 lb ai) ACCENT® SC. Treat heavy infestations of weeds before they become too competitive with the crop, especially where soil moisture and/or fertility are limited. ACCENT® SC provides weed control via foliar absorption.

ACCENT® SC only controls those weeds that have emerged. For later-emerging weeds, a second application or a timely cultivation is required. Applications made to weeds larger than the size indicated on this label or to weeds under stress may result in unsatisfactory control.

As weeds mature, their sensitivity to ACCENT® SC decreases. As grassy weeds become mature (more than 3 tillers), they may not reach the size listed below, due to drought or other environmental factors. Treat grassy weeds that are maturing rapidly before they reach the stages listed in the table Weeds controlled with 12 fluid ounces (0.031 lb ai) ACCENT® SC.

SEQUENTIAL APPLICATIONS

In the event that a subsequent flush of weeds, or a regrowth of previously treated weeds occur, a second application of ACCENT® SC may be applied.

WEEDS CONTROLLED IN INZEN™ TRAIT GRAIN SORGHUM

Weeds controlled with 12 fluid ounces (0.031 lb ai) ACCENT® SC herbicide

Grasses	Maximum Height or Diameter
Barnyardgrass†	4"
Broadleaf signalgrass	2"
Crabgrass (large)*	2"
Foxtails (bristly, giant†, green†, yellow†)	4"
Itchgrass	6"
Panicum (Texas, browntop)	3"
fall	4"
Ryegrass (Italian, perennial) †	6"
Sandbur(field, longspine)*	3"
Wild Oats†	4"
Wild proso millet	4"
Witchgrass	6"

† Naturally occurring resistant biotypes are known to occur. If weed escapes occur, treat with an herbicide having a mode of action other than group 2 and/or use non-chemical methods to remove escapes, as practicable, with the goal of preventing further seed production.

* Refer to **Specific Weed Instructions** section of this label

SPECIFIC WEED INSTRUCTIONS

Crabgrass (large): Requires the application of a soil applied herbicide that is effective in controlling large crabgrass, including CINCH® (AI: S-metolachlor; Reg. No. 352-625) or CINCH® ATZ (AI: S-metolachlor + atrazine; Reg. No. 352-624), followed by the post emergence application of ACCENT® SC herbicide at 12 fluid ounces/acre plus COC and ammonium nitrogen fertilizer. Adequate moisture is required after application of these soil applied herbicides to provide activation for weed control to occur. Cultivation or retreatment with ACCENT® SC plus COC and ammonium nitrogen fertilizer may be required for additional control of later emerging grasses. ACCENT® SC will not control or suppress smooth crabgrass.

Sandbur (field, longspine): Requires the use of COC plus ammonium nitrogen fertilizer. Cultivation or re-treatment may be required.

SPRAY ADJUVANTS FOR USE IN INZEN™ TRAIT GRAIN SORGHUM

ACCENT® SC herbicide contains a proprietary adjuvant system that aids in the control of many difficult weed species. Adjuvants including petroleum crop oil concentrate (COC) or nonionic surfactant (NIS) must be used with ACCENT® SC herbicide. In addition, an ammonium nitrogen fertilizer must be used unless specifically prohibited by tank mix partner labeling. Consult local DuPont fact sheets, technical bulletins, and service policies prior to using other adjuvant systems. If another herbicide is tank mixed with ACCENT® SC, select adjuvants authorized for use with both products, products must contain only EPA-exempt ingredients (40 CFR 1001).

Petroleum Crop Oil Concentrate (COC)

- Petroleum-based crop oil concentrates are the preferred adjuvant systems in arid areas.
- Apply up to 1% v/v (1 gallon per 100 gallons spray solution) or 2% under arid conditions.
- Oil adjuvants must contain at least 80% high quality, petroleum (mineral) or modified vegetable seed Oil with at least 15% surfactant emulsifiers.
- For aerial applications apply 0.5% v/v (2 quarts product per 100 gallons spray solution).

Nonionic Surfactant (NIS)

- Apply up to 0.25% v/v (1 quart per 100 gallons spray solution) or 0.5% v/v under arid conditions.
- Surfactant products must contain at least 60% nonionic surfactant with a hydrophilic/lipophilic balance (HLB) greater than 12.

Ammonium Nitrogen Fertilizer

- Use 2 quarts/acre of a high-quality urea ammonium nitrate (UAN), including 28%N or 32%N, or 2 pounds/acre of a spray- grade ammonium sulfate (AMS). Use 4 quarts/acre UAN or 4 pounds/acre AMS under arid conditions.
- **Do not** use liquid nitrogen fertilizer as the total carrier solution.

Special Adjuvant Types

- Combination adjuvant products may be used at doses that provide the required amount of NIS, COC, MSO and/or ammonium nitrogen fertilizer. Consult product literature for use rates and restrictions.

- In addition to the adjuvants specified above, other adjuvant types may be used if they provide the same functionality and have been evaluated and approved by DuPont Product Management. Consult separate DuPont technical bulletins for detailed information before using adjuvant types not specified on this label.

TANK MIXING

ACCENT® SC herbicide may be tank mixed with 2,4-D low volatile-ester, dicamba, atrazine, "Starane Ultra" (AI: fluroxypyr; Reg. No. 62719-77), and ALLY® XP (AI: metsulfuron; Reg. No. 352-435, 279-9575) herbicides registered for use in grain sorghum. **Do not** use COC (crop oil concentrate) when tank mixing with 2,4-d or dicamba. When tank mixing with 2,4-d or dicamba expect some crop response in the form of rolled leaves, leaning, brace root malformation and/or brittle stems. **Do not** apply 2,4-D or dicamba if this potential for injury is not acceptable. **Do not** tank mix with "Huskie" brands (AI: pyrasulfotole + bromoxynil) as significant grass antagonism, and crop injury can result. Refer to the labels of all tank mix products for information regarding use information (including rates, timing, application information, and sprayer cleanup), product precautions and restrictions. The most restrictive language on either label shall apply. If those instructions conflict with this label, **do not** tank mix the herbicide with ACCENT® SC.

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

CULTIVATION

A timely cultivation may be necessary to control suppressed weeds or weeds that emerge after an application of ACCENT® SC.

Optimum timing for cultivation is 7–14 days after ACCENT® SC application or upon seeing the establishment of new weeds.

GRAZING / PREHARVEST INTERVALS FOR GRAIN SORGHUM CONTAINING INZEN™ TRAIT

Forage may be cut, and livestock may be grazed once the crop has reached the mature forage stage (soft dough growth stage 7). Grain and stover may be harvested once the crop has reached the mature grain stage (physiological maturity growth stage 9).

CROP ROTATION

Rotational crops vary in their response to low concentrations of ACCENT® SC remaining in the soil. ACCENT® SC dissipates rapidly in warm, acidic, microbiologically active soils.

The amount of ACCENT® SC which may be present in the soil depends on application rate, soil pH and organic matter content, elapsed time since application, crop production practices, and environmental factors.

Injury to rotational crops may occur in high-pH, cold soils if dry weather prevails between application and rotational crop planting. Consult your local DuPont representative for additional guidelines.

Determine soil pH by laboratory analysis using the 1:1 soil:water suspension method on representative soil samples taken at 0–4" depth. Soil pH varies within fields; therefore, base recropping on the highest soil pH within each field. Consult local extension publications for soil sampling procedures.

The following rotational intervals must be observed when using ACCENT® SC at a maximum of 24 fluid

ounces (0.062 lb ai):

ACCENT® SC ROTATIONAL CROP GUIDELINE – 1

No soil pH restrictions

Rotational Crop	Interval in Months
Corn (field, seed)	Anytime
Corn (pop, sweet)*	10
Soybeans	0.5 (15 days)
Cereals, spring (barley, oats, rye, wheat)	8
Cereals, winter (barley, oats, rye, wheat)	4
Cotton	10
Dry Beans, Peas, Snap Beans	10
Alfalfa**	12
Red Clover**	12
Sorghum (All types including hybrids containing the INZEN™ trait)	18
Other Crops	See Rotational Crop Guidelines 2 and 3

* Except the sweet corn varieties “Merit”, “Carnival”, and “Sweet Success”, for which the minimum time interval is 15 months.

**Except for the state of Kansas east of highway 75, for Minnesota east and south of the Red River Valley and for the states east of the line formed by the western borders of Iowa, Missouri, Arkansas, and Louisiana, where the minimum time interval is 10 months.

ACCENT® SC ROTATIONAL CROP GUIDELINE – 2

With Soil pH ≤ 7.5 restrictions

Crop	Rotational Interval in months	
	pH 7.5	pH > 7.5
Sunflowers*	11	18
All other crops not listed in Rotational Guidelines 1 or 2	See Rotational Guideline 3	

*Precipitation following application must exceed 14” prior to planting sunflowers.

ACCENT® SC ROTATIONAL CROP GUIDELINE – 3

With soil pH < 6.5 restrictions

Crop	Rotational Interval in Months	
	pH 6.5	pH > 6.5
Sugarbeets*, potatoes**	10	18***
All other crops not listed in rotational guidelines 1 or 2	10	18

* Except on irrigated sites in Colorado, Wyoming, Nebraska, Texas, Michigan, and Ohio, where precipitation following application must exceed 25” prior to planting beets, where the interval is 10 months on soils with pH < 7.5. Sites in Minnesota east and south of the red river Valley may follow these guidelines provided maximum rates of ACCENT® SC do not exceed 12 fluid oz (0.031 lb ai).

**Irrigated potatoes following irrigated corn treated in the States of Washington or Idaho, or Utah can be

planted 10 months after using ACCENT® SC on sprinkler irrigated corn with no soil pH restrictions, providing the maximum use rate on corn does not exceed 18 fluid oz (0.046 lb ai) ACCENT® SC per year. Corn treated with ACCENT® SC must be grown to maturity and receive a minimum of 18 inches of irrigation water before potatoes can be planted at this rotational interval. Injury to potatoes may occur if less than 18 inches of irrigation is used on the previous corn crop. ACCENT® SC may not be used in a tankmix or sequential application program with other ALS-inhibiting herbicides (including “Beacon”; AI: primisulfuron-methyl; Reg. No. 100-705).

***In North Dakota and northwest Minnesota, the cumulative precipitation in the 18 months following application must exceed 28" in order to rotate to sugarbeets or potatoes.

ROTATIONAL CROP GUIDELINES - 4 may be observed when using a single application of ACCENT® SC per year with a maximum use rate of 12 fluid ounces (0.031 lb ai) product. Extend rotational intervals to 12 months if drought conditions prevail after application and before the rotational crop is planted, unless sprinkler irrigation has been applied and totals greater than 15" during the growing season.

ACCENT® SC ROTATIONAL CROP GUIDELINES – 4

With 12 fluid ounces (0.031 lb ai) maximum use rate

Crop	Rotational Interval in Months
Alfalfa*	10
Canola	10
Flax**	10
Sorghum (All types including hybrids containing the INZEN™ trait)	18
Potato	10
Red clover	10
Sunflower	10

*On sprinkler irrigated fields in Idaho, Utah, and Northern Nevada it is best to use deep fall tillage including plowing prior to planting alfalfa. Product degradation may be less on furrow irrigated soils and may result in some crop injury.

** Extend rotational intervals to 18 months if drought conditions prevail after application and before the rotational crop is planted, unless sprinkler irrigation has been applied and totals greater than 15" during the growing season.

GROUND APPLICATION

Broadcast Application

- Use a minimum of 15 gallons of water per acre (15 GPA) for best performance. Use a minimum of 10 gallons of water per acre (GPA) for light, scattered stands of weeds.
- Overlaps or starting, stopping, slowing, and turning while spraying may result in crop injury.

Band Application

For band applications, use proportionately less spray mixture, and carefully calibrate the band applicator to not exceed the labeled rate. Carefully follow the manufacturer’s instructions for nozzle type (flat fans), orientation, distance of nozzles from the crop and weeds, spray volumes, calibration and spray pressure.

AERIAL APPLICATION

Aerial application is not permitted in New York and California.

Use nozzle types and arrangements that will provide optimum spray distribution and maximum coverage at a minimum of 3 GPA.

SPRAY ADJUVANTS FOR USE IN FIELD CORN GROWN FOR GRAIN & SEED, POPCORN, & SWEET CORN

ACCENT® SC contains a proprietary adjuvant system that aids in the control of many difficult weed species. In certain arid climates, the use of additional adjuvants may be required for optimal weed control. In addition, an ammonium nitrogen fertilizer must be used unless specifically prohibited by tank mix partner labeling. Consult local DuPont fact sheets, technical bulletins, and service policies prior to using other adjuvant systems. If another herbicide is tank mixed with ACCENT® SC, select adjuvants authorized for use with both products, products must contain only EPA-exempt ingredients (40 CFR 1001).

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

Crop Oil Concentrate (COC) - Petroleum or Modified Seed Oil (MSO)

- Apply up to 1% v/v (1 gallon per 100 gallons spray solution) under arid conditions.
- MSO adjuvants may be used up to 0.5% v/v (0.5 gallons per 100 gallons spray solution) if specifically noted on adjuvant product labeling or if specified on local DuPont product literature or service policies.
- Oil adjuvants must contain at least 80% high quality, petroleum (mineral) or modified vegetable seed oil with at least 15% surfactant emulsifiers.

Nonionic Surfactant (NIS)

- Apply up to 0.25% v/v (1 quart per 100 gallons spray solution) under arid conditions.
- Surfactant products must contain at least 60% nonionic surfactant with a hydrophilic/lipophilic balance (HLB) greater than 12.

Ammonium Nitrogen Fertilizer

- Use 2 quarts/acre of a high-quality urea ammonium nitrate (UAN), including 28%N or 32%N, or 2 pounds/acre of a spraygrade ammonium sulfate (AMS). Use 4 quarts/acre UAN or 4 pounds/acre AMS under arid conditions.
- **Do not** use liquid nitrogen fertilizer as the total carrier solution.

Special Adjuvant Types

- Combination adjuvant products may be used at doses that provide the required amount of NIS, COC, MSO and/or ammonium nitrogen fertilizer. Consult product literature for use rates and restrictions.
- In addition to the adjuvants specified above, other adjuvant types may be used if they provide the same functionality and have been evaluated and approved by DuPont Product Management. Consult separate DuPont technical bulletins for detailed information before using adjuvant types not specified on this label.

MIXING INSTRUCTIONS

Select a spray volume that will ensure thorough coverage and a uniform spray pattern. If tank mixing with other herbicides, always consult the label of the tank mix partner(s) for minimum spray volume requirements and apply the tank mixture using a water volume as directed for all products.

1. Always start with a clean and empty sprayer tank.
2. Fill the tank with clean water one half of the required spray volume.
3. With the agitator running, add the required amount of ACCENT® SC. Continue to agitate for a minimum of 5 minutes to ensure that ACCENT® SC is completely dispersed.
4. If tank mixing ACCENT® SC with another herbicide, follow this mixing order: dry flowables and soluble granules, followed by liquids, then oil dispersions (OD) or emulsifiable concentrates (EC). Maintain continuous agitation.
5. Add the rest of the water.
6. If required for the tank mixture, add the appropriate adjuvant. If an antifoam agent is required, add last.
7. Continue agitation sufficient enough to maintain a uniform spray solution.
8. Refer to the tank mix sections of this booklet for mixing order and other mixing instructions.

TANK MIX COMPATIBILITY TESTING

Perform a jar test prior to tank mixing to ensure compatibility of ACCENT® SC and other pesticides. Use a clear glass quart jar with lid and mix the tank mix ingredients in their relative proportions. Invert the jar containing the mixture several times and observe the mixture for approximately 1/2 hour. If the mixture balls-up, forms flakes, sludges, gels, oily films or layers, or other precipitates, it is not compatible. **Do not** use the tank mix combination.

ENVIRONMENTAL CONDITIONS AND BIOLOGICAL ACTIVITY

ACCENT® SC provides best results when applied to young, actively growing weeds. Applications made during warm, moist conditions (70°F or more) and adequate soil moisture both before and after application maximizes performance.

The degree and duration of control depend on spray coverage, weed spectrum, weed size, growing conditions before and after treatment, soil moisture, and adjuvant selection.

ACCENT® SC is rainfast in 4 hours.

Treating weeds that exceed maximum label height or that are under stress may result in incomplete control. Poor weed control or crop injury may result from applications made to plants under stress from:

- abnormally hot or cold weather
- environmental conditions including drought, water-saturated soils, hail damage, or frost
- disease, insect, or nematode injury
- prior herbicide or carryover from a previous year's herbicide application

Severe stress from conditions preceding or immediately following application may also result in crop injury

or poor weed control.

Stress affects all weeds, but especially weeds including woolly cupgrass, green and yellow foxtail, and wild proso millet.

ACCENT® SC rapidly inhibits the growth of susceptible weeds, reducing weed competition within as little as 6 hours after application. Susceptible plants are controlled in 7–21 days.

Do not exceed labeled application rates. **Do not** tank mix ACCENT® SC with other products that contain the same active ingredients as ACCENT® SC (nicosulfuron) unless the label of either tank mix partner specifies the maximum rate that may be used.

If the crop or grass weeds are under stress, delay application until stress passes and both weeds and crop resume active growth.

SPRAYER PREPARATION/CLEANUP

It is important that spray equipment is clean and free of previous pesticide deposits before using ACCENT® SC and then properly cleaned out following application. Clean all application equipment before applying ACCENT® SC. Follow the cleanup procedures specified on the label of the product previously sprayed. If no cleanup procedure is provided, use the procedure that follows.

Immediately following applications of ACCENT® SC, thoroughly clean all mixing and spray equipment to avoid subsequent crop injury.

Note:

- When cleaning spray equipment before applying ACCENT® SC, read and follow label directions for proper rinsate disposal of the product previously sprayed.
- Steam cleaning of aerial spray tanks will help to dislodge any visible pesticide deposits.
- When spraying or mixing equipment will be used over an extended period to apply multiple loads of ACCENT® SC, partially fill the tank with fresh water at the end of each day of spraying, flush the boom and hoses, and allow to sit overnight.

Cleanup Procedure

1. Drain the tank and thoroughly hose down the interior surfaces. Flush the tank, hoses, and boom with clean water for a minimum of 5 min.
2. Partially fill the tank with clean water and add one gallon of household ammonia* (containing 3% active) for every 100 gallons of water. Finish filling the tank with water, then flush the cleaning solution through the hoses, boom, and nozzles. Add more water to completely fill the tank and allow to agitate/recirculate for at least 15 min. Again, flush the hoses, boom, and nozzles with the cleaning solution, then drain the tank.
3. Repeat Step 2.
4. Remove the nozzles, screens and the end caps of sprayer booms and clean separately in a bucket containing the cleaning agent and water.
5. Thoroughly rinse the tank with clean water for a minimum of 5 min, flushing the water through the hoses and boom.

* Equivalent amounts of an alternate strength ammonia solution or a tank cleaner may be used.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage and disposal.

Pesticide Storage: Store product in original container only. Store in a cool, dry place.

Pesticide Disposal: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

Container Handling: Refer to the Net Contents section of this product's labeling for the applicable "Nonrefillable Container" or "Refillable Container" designation.

Nonrefillable Rigid Plastic and Metal Containers (Capacity Equal to or Less Than 5 Gallons):

Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then, for plastic containers, offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration. Do not burn, unless allowed by state and local ordinances. For Metal containers, offer for recycling if available or reconditioning if appropriate, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

Nonrefillable Rigid Plastic and Metal Containers (Capacity Greater Than 5 Gallons): Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then, for plastic containers, offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration. Do not burn, unless allowed by state and local ordinances. For Metal containers, offer for recycling if available or reconditioning if appropriate, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

Nonrefillable Rigid Plastic and Metal Containers, e.g., Intermediate Bulk Containers [IBC] (Size or Shape Too Large to be Tipped, Rolled or Turned Upside Down): Nonrefillable container. Do not reuse or refill this container. Clean container promptly after emptying the contents from this container into application equipment or mix tank and before final disposal using the following pressure rinsing procedure. Insert a lance fitted with a suitable tank cleaning nozzle into the container and ensure that the water spray thoroughly covers the top, bottom and all sides inside the container. The nozzle manufacturer generally provides instructions for the appropriate spray pressure, spray duration and/or spray volume. If the manufacturer's instructions are not available, pressure rinse the container for at least 60 seconds using a minimum pressure of 30 PSI with a minimum rinse volume of 10% of the container volume. Drain, pour or pump rinsate into application equipment or rinsate collection system. repeat this pressure rinsing procedure two more times. Then, for plastic containers, offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration. For Metal containers, offer for recycling if available or reconditioning if appropriate, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

All Refillable Containers: Refillable container. *Refilling Container:* refill this container with ACCENT® SC containing nicosulfuron only. do not reuse this container for any other purpose. Cleaning before

refilling is the responsibility of the refiller. Prior to refilling, inspect carefully for damage including cracks, punctures, abrasions, worn out threads and closure devices. If damage is found, do not use container, contact DuPont at the number below for instructions. Check for leaks after refilling and before transporting. If leaks are found, do not reuse or transport container, contact DuPont at the number below for instructions. *Disposing of Container:* Do not reuse this container for any other purpose other than refilling (see preceding). Cleaning the container before final disposal is the responsibility of the person disposing of the container. To clean the container before final disposal, use the following pressure rinsing procedure. Insert a lance fitted with a suitable tank cleaning nozzle into the container and ensure that the water spray thoroughly covers the top, bottom and all sides inside the container. The nozzle manufacturer generally provides instructions for the appropriate spray pressure, spray duration and/or spray volume. If the manufacturer's instructions are not available, pressure rinse the container for at least 60 seconds using a minimum pressure of 30 PSI with a minimum rinse volume of 10% of the container volume. Drain, pour or pump rinsate into application equipment or rinsate collection system. Repeat this pressure rinsing procedure two more times. Then, for plastic containers, offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration. Do not burn, unless allowed by state and local ordinances. For Metal containers, offer for recycling if available or reconditioning if appropriate, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

Do not transport if container is damaged or leaking. If the container is damaged, leaking or obsolete, or in the event of a major spill, fire or other emergency, contact DuPont at 1-800-441-3637, day or night.

NOTICE TO BUYER: Purchase of this material does not confer any rights under patents of countries outside of the United States.

®™ Trademarks of Dow AgroSciences, DuPont or Pioneer and their affiliated companies or respective owners

Ally is a registered trademark of FMC Corp.

"Huskie" and "LibertyLink" are trademarks or registered trademarks of Bayer CropScience

"Basagran", "Clarity", "Marksman", are trademarks or registered trademarks of BASF Corporation

"Bicep", "Callisto", "Exceed", "Force" and "Northstar" are registered trademarks of a Syngenta Group Company

"Aztec", "Counter", "Fortress", "Smart Choice" are registered trademarks of Amvac Chemical Corporation

"Roundup Ready" is a registered trademark of Monsanto Technology LLC

LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read this Limitation of Warranty and Liability before buying or using This product. If the Terms Are Not Acceptable, Return the Product at Once, Unopened, and the Purchase Price Will Be Refunded. It is impossible to eliminate all risks associated with the use of this product. Such risks arise from weather conditions, soil factors, off target movement, unconventional farming techniques, presence of other materials, the manner of use or application, or other unknown factors, all of which are beyond the control of DuPont. These risks can cause: ineffectiveness of the product, crop injury, or injury to non-target crops or plants. WHEN YOU BUY OR USE THIS PRODUCT, YOU AGREE TO ACCEPT THESE RISKS.

DuPont warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for the purpose stated in the directions for use, subject to the inherent risks described above, when used in accordance with the directions for use under normal conditions.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, DUPONT MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR OF MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL DUPONT OR SELLER BE LIABLE FOR ANY INCIDENTAL, CONSEQUENTIAL OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. BUYER'S OR USER'S

BARGAINED-FOR EXPECTATION IS CROP PROTECTION. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER AND THE EXCLUSIVE LIABILITY OF DUPONT OR SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY OR CONTRACT, NEGLIGENCE, TORT OR STRICT LIABILITY), WHETHER FROM FAILURE TO PERFORM OR INJURY TO CROPS OR OTHER PLANTS RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR AT THE ELECTION OF DUPONT OR SELLER THE REPLACEMENT OF THE PRODUCT.

To the extent consistent with applicable law that allows such requirement, DuPont or its Ag retailer must have prompt notice of any claim so that an immediate inspection of buyer's or user's growing crops can be made. Buyer and all users shall promptly notify DuPont or a DuPont Ag retailer of any claims, whether based on contract, negligence, strict liability, other tort or otherwise, or be barred from any remedy.

This Limitation of Warranty and Liability may not be amended by any oral or written agreement.

For product information call: 1-800-258-3033

© 1996-2019 E. I. du Pont de Nemours and Company, Chestnut Run Plaza, 974 Centre Road, Wilmington, DE 19805 U.S.A.

®™ Trademarks of Dow AgroSciences, DuPont or Pioneer and their affiliated companies or respective owners

EPA accepted __/__/__