

352-751

03-01-2011

1/21



U.S. ENVIRONMENTAL PROTECTION AGENCY
Office of Pesticide Programs
Registration Division (7504P)
Ariel Rios Building
1200 Pennsylvania Ave., NW
Washington, D.C. 20460

EPA Registration
 Number:
 352-751

Date of Issuance:
 MAR 1 2011

NOTICE OF PESTICIDE:
 Registration
 Reregistration
 (under FIFRA, as amended)

Term of Issuance: **Unconditional**

Name of Pesticide Product:
DuPont Agility SG Herbicide (with TotalSol soluble granules)

Name and Address of Registrant (include ZIP Code):
E. I. du Pont de Nemours and Company
1007 Market Street
Wilmington, DE 19898

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act. Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.


EPA received a label amendment request submitted on August 25, 2010. EPA grants this request under the authority of section 3(c)(5) of the Federal Insecticide, Fungicide and Rodenticide Act, as amended. With this accepted labeling, all requirements set forth in the Reregistration Eligibility Decision for **Dicamba** have been satisfied. Therefore, EPA reregisters the product listed above. This action is taken under the authority of section 4(g)(2)(c) of the Federal Insecticide, Fungicide, and Rodenticide Act, as amended. Reregistration under this section does not eliminate the need for continual reassessment of pesticides. EPA may require submission of data at any time to maintain the registration of your product.

Submit one (1) copy of final printed labeling. Amended labeling will supersede all previously accepted labels. A copy of your label stamped "Accepted" is enclosed for your records. Products shipped after twelve (12) months from the date of this Notice or the next printing of your label, whichever occurs first, must bear the new revised label.

If you have any questions regarding this Notice, please contact Mindy Ondish at (703) 605-0723 or at ondish.mindy@epa.gov.


Signature of Approving Official:
Kable Bo Davis
Product Manager 25
Herbicide Branch
Registration Division

Date:
 3-1-11



DuPont™ Agility™ SG
herbicide (with TotalSol® soluble granules)

DRAFT LABEL



DUPONT™ AGILITY™ SG HIGHLIGHTS

- For selective postemergence broadleaf weed control in Wheat, Barley, Triticale and Fallow.
- Apply at the rate of 1.6 to 3.2 ounces per acre (see Use Rate, Application Timing and Tank Mixtures).
- May be applied by ground or by air.
- Use in tank mixtures with other registered herbicides for broader spectrum weed control (see Tank Mixtures).
- Consult label text for complete instructions. Always read and follow label "Directions For Use".

TABLE OF CONTENTS

PRECAUTIONARY STATEMENTS1

DIRECTIONS FOR USE3

PRODUCT INFORMATION3

PRECAUTIONS3

ENVIRONMENTAL CONDITIONS AND BIOLOGICAL ACTIVITY4

WEEDS CONTROLLED4

FALLOW5

WHEAT, BARLEY AND TRITICALE5

 USE RATE5

 APPLICATION TIMING5

 TANK MIXTURES5-6

 SPECIFIC WEED PROBLEMS6

APPLICATION INFORMATION7

 PRODUCT MEASUREMENT7

 SPRAY ADJUVANTS7

 GROUND APPLICATION7

 AERIAL APPLICATION7

 SEQUENTIAL APPLICATIONS8

 MIXING INSTRUCTIONS8

 SPRAY EQUIPMENT8

 AT THE END OF THE DAY8

 AFTER SPRAYING AGILITY™ SG AND BEFORE SPRAYING CROPS OTHER THAN WHEAT, BARLEY OR TRITICALE8

CROP ROTATION9

 MINIMUM ROTATIONAL INTERVALS9

 FIELD BIOASSAY9

 GRAZING9

 CROP ROTATION TABLES10-14

SPRAY DRIFT MANAGEMENT15

RESISTANCE16

INTEGRATED PEST MANAGEMENT16

STORAGE AND DISPOSAL16

LIMITATION OF WARRANTY AND LIABILITY ...18



DuPont™ Agility™ SG

herbicide (with TotalSol®
soluble granules)

**For Use on Wheat, Barley, Triticale and
Fallow.**

AGILITY™ SG herbicide is a soluble granule for selective
postemergence weed control.

Active Ingredients	By Weight
Sodium salt of dicamba (3,6-dichloro-2-Methoxybenzoic acid)*	63.6%
Thifensulfuron methyl Methyl 3-[[[(4-methoxy-6-methyl-1,3,5- -triazin-2-yl) amino]carbonyl]amino]- sulfonyl]-2-thiophenecarboxylate	4.7%
Tribenuron methyl Methyl 2-[[[N-(4-methoxy-6-methyl-1,3,5- -triazin-2-yl)methylamino]carbonyl]- amino]sulfonyl]benzoate	2.4%
Metsulfuron methyl Methyl 2-[[[(4-methoxy-6-methyl -1,3,5- -triazin-2-yl)amino]carbonyl] amino]- sulfonyl]benzoate	1.9%
Other Ingredients	27.4%
TOTAL	100.0%

*This product contains 57.8% 3,6-dichloro-2-Methoxybenzoic acid
(dicamba)

EPA Reg. No. 352-751

EPA Est. No. _____


Nonrefillable Container

Net: _____

OR

Refillable Container

Net: _____



KEEP OUT OF REACH OF CHILDREN WARNING/AVISO

Si usted no entiende la etiqueta, busque a alguien para que se
la explique a usted en detalle. (If you do not understand this
label, find someone to explain it to you in detail.)

FIRST AID

IF IN EYES: Hold eye open and rinse slowly and gently
with water for 15-20 minutes. Remove contact lenses, if
present, after the first 5 minutes, then continue rinsing eye.
Call a poison control center or doctor for treatment advice.

IF ON SKIN OR CLOTHING: Take off contaminated
clothing. Rinse skin with plenty of water for 15-20 minutes.
Call a poison control center or doctor for treatment advice.

IF SWALLOWED: Call a poison control center or doctor
immediately for treatment advice. Have person sip a glass of
water if able to swallow. Do not induce vomiting unless told
to do so by a poison control center or doctor. Do not give
anything by mouth to an unconscious person.

Have the product container or label with you when calling a
poison control center or doctor, or going for treatment. You
may also contact 1-800-441-3637 for emergency medical
treatment information.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING! Causes substantial but temporary eye injury.
Do not get in eyes, on skin, or on clothing. Harmful if
swallowed or absorbed through skin. Prolonged or
frequently repeated skin contact may cause allergic
reactions in some individuals.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some of the materials that are chemical-resistant to this
product are made of any waterproof material. If you want
more options, follow the instructions for Category A on an
EPA chemical-resistance category selection chart.

**All mixers, loaders, and applicators and other handlers
must wear:**

- Long-sleeved shirt and long pants.
- Chemical resistant gloves (except for pilots).
- Shoes plus socks.
- Protective eyewear.

See engineering controls for additional requirements and
exceptions.

Discard clothing and other absorbent materials that have
been drenched or heavily contaminated with this product's
concentrate. Do not reuse them. Follow manufacturer's
instructions for cleaning/maintaining PPE. If no such
instructions for washables exist, use detergent and hot water.
Keep and wash PPE separately from other laundry.

4
12

Engineering Control Statements:

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides 40 CFR 170.240(d)(4-6), the handler PPE requirements may be reduced or modified as specified in the WPS. Pilots must use cockpits in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides 40 CFR 170.240(d)(4-6).

USER SAFETY RECOMMENDATIONS

USERS SHOULD: Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

Keep out of lakes, streams, or ponds. For terrestrial uses, do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate. Apply this product only as directed on the label.

Dicamba is known to leach through soil into ground water under certain conditions as a result of agricultural use. Use of this product in areas where soils are permeable, particularly where the water table is shallow, may result in ground water contamination.

DO NOT discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans, or other waters unless accordance with the requirements of a National Pollution Discharge Elimination System (NPDES) permit and the permitting authority has been notified in writing prior to discharge. DO NOT discharge effluent containing this product to sewer systems without previously notifying the local sewage treatment plant authority. For guidance, contact your State Water Board or Regional Office of the EPA.

Ground and Surface Water Protection

Point source contamination: To prevent point source contamination, do not mix, load this pesticide product within 50 feet of wells (including abandoned wells and drainage wells), sink holes, perennial or intermittent streams and rivers, and natural or impounded lakes and reservoirs. Do not apply pesticide product within 50 feet of wells. This setback does not apply to properly capped or plugged abandoned wells and does not apply to impervious pad or properly diked mixing/loading areas as described below. Mixing, loading, rinsing, or washing operations performed within 50 feet of a well are allowed only when conducted on an impervious pad constructed to withstand the weight of the heaviest load that may be on or move across the pad. The pad must be self-contained to prevent surface water flow over or from the pad. The pad capacity

must be maintained at 110% that of the largest pesticide container or application equipment used on the pad and have sufficient capacity to contain all product spills, equipment or container leaks, equipment wash waters, and rainwater that may fall on the pad. The containment capacity does not apply to vehicles delivering pesticide shipments to the mixing/loading site. States may have in effect additional requirements regarding wellhead setbacks and operational containment.

Care must be taken when using this product to prevent: a) back siphoning into wells, b) spills or c) improper disposal of excess pesticide, spray mixtures or rinsates. Check valves or antisiphoning devices must be used on all mixing equipment.

Movement by surface runoff or through soil: Do not apply under conditions which favor runoff. Do not apply to impervious substrates such as paved or highly compacted surfaces in areas with high potential for ground water contamination. Ground water contamination may occur in areas where soils are permeable or coarse and ground water is near the surface. Do not apply to soils classified as sand with less than 3% organic matter and where ground water depth is shallow. To minimize the possibility of ground water contamination, carefully follow application rate recommendations as affected by soil type in the product information section of this label.

Movement by water erosion of treated soil: Do not apply or incorporate this product through any type of irrigation equipment nor by flood or furrow irrigation. Ensure treated areas have received at least one-half inch rainfall (or irrigation) before using tailwater for subsequent irrigation of other fields.

Endangered Species Concerns

The use of any pesticide in a manner that may kill or otherwise harm an endangered species or adversely modify their habitat is a violation of federal law.

PESTICIDE HANDLING

- Calibrate sprayers only with clean water away from the well site.
- Make scheduled checks of spray equipment.
- Ensure that all operation employees accurately measure pesticides.
- Mix only enough product for the job at hand.
- Avoid overfilling of spray tank.
- Do not discharge excess material on the soil at a single spot in the field, grove, or mixing/loading station.
- Avoid storage of pesticides near well sites.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours. Exception: If the product is soil injected or soil incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated areas if there will be no contact with anything that has been treated. PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls worn over short-sleeve shirt and short pants
- Chemical-resistant footwear plus socks
- Chemical-resistant gloves made of any waterproof material.
- Chemical-resistant headgear for overhead exposure
- Protective eyewear

DuPont™ AGILITY™ SG must be used only in accordance with instructions on this label or in supplemental DuPont publications.

DuPont will not be responsible for losses or damages resulting from the use of this product in any manner not specified by DuPont.

AGILITY™ SG is for use on wheat, barley, triticale and fallow in most states, check with your state extension service or Department of Agriculture before use, to be certain AGILITY™ SG is registered in your state. AGILITY™ SG is not registered for use in Alamosa, Conejos, Costilla, Rio Grande, and Saquache counties of Colorado unless use is directed otherwise by supplemental labeling.

PRODUCT INFORMATION

AGILITY™ SG herbicide is a water soluble granule that is used for selective postemergence weed control in wheat (including durum), barley, triticale and fallow.

The best control is obtained when AGILITY™ SG is applied to young, actively growing weeds. The degree and duration of control may depend on the following:

- weed spectrum and infestation intensity
- weed size at application
- environmental conditions at and following treatment

AGILITY™ SG is noncorrosive, nonflammable, and does not freeze. AGILITY™ SG should be mixed and completely dissolved in water and applied as a uniform broadcast spray (See Tank Mixtures and Mixing Instructions sections for use with Liquid Nitrogen Fertilizer Solutions).

RESTRICTIONS AND PRECAUTIONS

- Do not use AGILITY™ SG plus Malathion or Lorsban as crop injury may result.
- Do not harvest sooner than 45 days after the last application of AGILITY™ SG.
- AGILITY™ SG is only registered on wheat, barley, triticale and fallow. Do not use on any other crop.
- Varieties of wheat (including durum), barley and triticale may differ in their response to various herbicides. DuPont recommends that you first consult your state experiment station, university, or extension agent as to sensitivity to any herbicide. If no information is available, limit the initial use of AGILITY™ SG to a small area.
- Under certain conditions such as heavy rainfall, prolonged cold weather, or wide fluctuations in day/night temperatures prior to or soon after AGILITY™ SG application, temporary discoloration and/or crop injury may occur. To reduce the potential of crop injury, tank mix AGILITY™ SG with 2,4-D and apply after the crop is in the tillering stage of growth.
- Do not apply to wheat, barley or triticale that is stressed by severe weather conditions, drought, low fertility, water-saturated soil, disease, or insect damage, as crop injury may result. Risk of injury is greatest when crop is in the 2 to 5- leaf stage. Severe winter stress, drought, disease, or insect damage following application also may result in crop injury.
- Do not apply to wheat, barley or triticale undersown with legumes and grasses, because injury to the forages will result.
- Do not treat irrigation ditches or water used for crop irrigation or domestic uses.

- Do not apply this product through any type of irrigation system.
- For ground applications applied when dry, dusty field conditions exist, control of weeds in wheel track areas may be reduced. The addition of 2,4-D or MCPA may improve weed control under these conditions.
- Injury to or loss of desirable trees or vegetation may result from failure to observe the following:
 - Do not apply, drain or flush equipment on or near desirable trees or other plants or on areas where their roots may extend, or in locations where the chemical may be washed or moved into contact with their roots.
 - Do not use on lawns, walks, driveways, tennis courts, or similar areas. Prevent drift of spray to desirable plants.
- Injury to or loss of adjacent sensitive crops and vegetation may result from failure to observe the following:
 - Take all necessary precautions to avoid all direct or indirect contact (such as spray drift) with non-target plants or areas.
 - Do not apply adjacent to sensitive crops when the temperature on the day of application is expected to exceed 85°F as drift is more likely to occur.
 - Carefully observe all sprayer cleanup instructions both prior to and after using this product, as spray tank residue may damage crops other than wheat, barley or triticale.

ENVIRONMENTAL CONDITIONS AND BIOLOGICAL ACTIVITY

DuPont™ AGILITY™ SG is absorbed through the roots, shoots, and foliage of plants, rapidly inhibiting the growth of susceptible weeds. One to three weeks after postemergence application to weeds (2 to 5 weeks for wild garlic), susceptible plants are controlled. In warm, moist conditions, the expression of herbicide symptoms is accelerated; in cold, dry conditions, expression of herbicide symptoms is delayed.

AGILITY™ SG will provide up to 4 to 6 weeks of residual weed control. Susceptible weeds may germinate and emerge a few days after postemergence applications, but growth then ceases and leaves become chlorotic 3- 5 days after emergence. Death of leaf tissue and growing point will follow in some species, while others may remain green but stunted and noncompetitive.

AGILITY™ SG provides the best control of weeds in vigorously growing crops that shade competitive weeds. Weed control in areas of thin crop stand or seeding skips may not provide satisfactory control. However, a crop canopy that is too dense at application can intercept spray and reduce weed control.

The herbicidal action of AGILITY™ SG may be less effective on weeds stressed from adverse environmental conditions (such as extreme temperatures or moisture), abnormal soil conditions, or cultural practices. In addition, weeds hardened-off by drought stress are less susceptible to AGILITY™ SG.

WEEDS CONTROLLED

AGILITY™ SG effectively controls the following weeds when used according to label directions:

- | | |
|--------------------------------------|---|
| Annual knawel | Lambsquarter |
| Annual sowthistle | (common, slimleaf) |
| Black mustard | London rocket |
| Black nightshade | Marshelder |
| Blue/Purple mustard * | Mayweed, chamomile |
| Broadleaf dock | Miners lettuce |
| Buckwheat (tartary, wild) | Narrowleaf lambsquarters |
| Bur buttercup (testiculate) | Nightflowering catchfly |
| Bushy wallflower/
Treadle mustard | Pennsylvania smartweed |
| Canada thistle * | Pigweed (prostrate, redroot,
rough, smooth, tumble) |
| Carolina geranium | Pineappleweed |
| Clasping pepperweed | Plains coreopsis |
| Coast fiddleneck (tarweed) | Prickly lettuce* |
| Common buckwheat | Redmaids |
| Common chickweed | Russian thistle* |
| Common cocklebur | Scentless chamomile /
mayweed |
| Common mallow | Shepherd's-purse |
| Common purslane | Smallflower buttercup |
| Common radish | Smallseed falseflax |
| Common ragweed | Smartweed (green,
ladysthumb, pale) |
| Common sunflower | Snow speedwell |
| Conical catchfly | Sticky chickweed |
| Corn chamomile | Stinking mayweed /
dogfennel |
| Corn cockle | Swinecress |
| Corn gromwell * | Tansymustard |
| Corn spurry | Tarweed fiddleneck |
| Cowcockle | Tumble/ Jim Hill mustard |
| Cress (mouse-ear) | Velvetleaf |
| Curly dock | Volunteer lentils |
| Cutleaf eveningprimrose | Volunteer peas |
| False chamomile | Volunteer sunflower |
| Field chickweed | Waterpod |
| Field pennycress (fanweed) | Wild buckwheat |
| Filaree (redstem, Texas) | Wild chamomile |
| Flixweed * | Wild garlic * |
| Groundsel (common) | Wild mustard |
| Henbit | Wild radish * |
| Knawel (German moss) | |
| Knotweed (prostrate) | |
| Kochia* | |
| Ladysthumb | |

WEEDS PARTIALLY CONTROLLED**

AGILITY™ SG partially controls the following weeds when used according to label directions:

- | | |
|--------------------|---|
| Broadleaf plantain | Nightshade (cutleaf, hairy,
silverleafs) |
| Catchweed bedstraw | Sowthistle (annual) * |
| Common dandelion | Tall waterhemp |
| Common yarrow | Vetch* (common, hairy) |
| Field bindweed | |
| Mallow (little) | |

* See the Specific Weed Problems section of this label for more information.

**Partial control: A visual reduction of weed population as well as a significant loss of vigor. For better results, use the highest labeled rate of AGILITY™ SG and include a tank mix partner such as 2,4-D, MCPA, or bromoxynil (such as "Buctril", "Bison", "Bronate" or "Bronate Advanced"). Refer to the Tank Mixtures section of this label.

FALLOW

Fallow Use Rate

Apply 1.6 to 3.2 ounces per acre of DuPont™ AGILITY™ SG to fallow fields (postharvest, fallow, crop stubble, set-aside). See SPRAY ADJUVANTS for proper adjuvant rates.

Fallow Application Timing

AGILITY™ SG can be applied either postharvest in the fall, spring, or summer during the fallow period or to crop stubble/set-aside acres. Apply AGILITY™ SG as a broadcast or spot treatment to emerged and actively growing weeds.

Tank Mixtures in Fallow

AGILITY™ SG can be tank mixed with other herbicides that are registered for use in fallow. Read and follow all manufacturers' label instructions for the companion herbicide. If those instructions conflict with this label, do not tank mix the herbicide with AGILITY™ SG.

WHEAT, BARLEY AND TRITICALE

USE RATE

Apply AGILITY™ SG at the rate of 1.6 to 3.2 ounces per acre to wheat, barley and triticale. See SPRAY ADJUVANTS for proper adjuvant rates.

Use 3.2 ounces per acre of AGILITY™ SG for heavy infestation of the weeds listed under Weeds Partially Controlled when application timing and environmental conditions are marginal (refer to Biological Activity and Environmental Conditions section of this label for best performance).

Use 1.6 to 2.4 ounces per acre of AGILITY™ SG for light infestation of the weeds listed under Weeds Controlled. Conditions at application should be optimum for effective treatment of these weeds.

Note: See Tank Mix Section for additional info on required combinations when used at less than 3.2 ounces per acre.

APPLICATION TIMING

Fall-seeded Wheat, Barley, and Triticale

Make applications after the crop is in the 2-leaf stage but before the jointing stage.

Spring-seeded Wheat

Make applications after the crop is in the 2-leaf stage but before it exceeds the 6-leaf stage. Early developing wheat varieties (such as TAM 107, Madison, or Wakefield) must receive application between early tillering and the jointing stage (stagger planting dates with these varieties to be certain that the applications occur before jointing). Durum and Wampum variety must receive application after tillering but before it exceeds the 6-leaf stage and should be made in combination with 2,4-D.

Spring-seeded Barley

Make applications after the crop is in the 2-leaf stage but before it exceeds the 4-leaf stage.

Do not tank mix AGILITY™ SG with 2,4-D in early season applications on spring-seeded barley.

Spring-seeded Triticale

Make applications after the crop is in the 2-leaf stage but before it exceeds the 6-leaf stage.

Additional Timing Information

Weed control may be reduced if rainfall or snowfall occurs soon after application. Six hours of dry weather are needed to allow AGILITY™ SG to be sufficiently absorbed by weed foliage.

If applied to irrigated wheat, barley or triticale the first post-treatment irrigation should be delayed for at least 6 hours after treatment and should not exceed 1 inch of water.

Do not apply AGILITY™ SG to stressed crops, as this may cause crop injury. To reduce the potential of crop injury, tank mix AGILITY™ SG with 2,4-D (ester formulations perform best-see TANK MIXTURES) and apply after the crop is in the tillering stage of growth.

TANK MIXTURES

AGILITY™ SG may be tank mixed with other suitable registered herbicides to control weeds listed as partially controlled, weeds resistant to AGILITY™ SG or weeds not listed under Weeds Controlled. Read and follow all manufacturers label instructions for the companion herbicide. If those instructions conflict with this label, do not tank mix the herbicide with AGILITY™ SG.

AGILITY™ SG can also be mixed with registered fungicides, insecticides, or liquid fertilizer for use on wheat, barley and triticale.

With 2,4-D (amine or ester) or MCPA (amine or ester)

AGILITY™ SG can be tank mixed with 2,4-D and MCPA (preferably ester formulations) herbicides for use on wheat, barley, triticale and fallow. For best results, add 2,4-D or MCPA herbicides to the tank at 1/8 to 3/8 lb active ingredient per acre.

In tank mixes containing 1/8 lb active ingredient 2,4-D or MCPA per acre, add 1 to 2 pt of non-ionic surfactant per 100 gal of spray solution; in tank mixes containing 1/4 to 3/8 lb active ingredient 2,4-D or MCPA per acre, add 1 pt of non-ionic surfactant per 100 gal of spray solution. Higher rates of 2,4-D or MCPA may be used, but do not exceed the highest rate allowed by those respective labels.

Always mix AGILITY™ SG in water prior to adding 2,4-D or MCPA and add the surfactant last. Read and follow all label instructions on timing, precautions, and warnings for these herbicides before using these tank mixtures.

With Bromoxynil containing products (such as "Buctril", "Bronate" or "Bison")

AGILITY™ SG may be tank mixed with bromoxynil containing herbicides registered for use on wheat, barley, triticale or fallow. For best results, add bromoxynil-containing herbicides to the tank at 3 to 6 oz active ingredient per acre (such as "Bronate" or "Bison" at 3/4 - 1 1/2 pt per acre).

Tank mixes of AGILITY™ SG plus Bromoxynil may result in reduced control of Canada thistle.

Read and follow all label instructions on timing, precautions, and warnings for these herbicides before using these tank mixtures. Follow the most restrictive labeling.

With "Stinger" or "Curtail"

DuPont™ AGILITY™ SG can be tank mixed with "Stinger" or "Curtail" herbicide for improved control of weeds in wheat, barley and triticale. Refer to the "Stinger" and "Curtail" labels for information regarding use restrictions, labeled crops, rotational cropping intervals, sprayer cleanup, use precautions and other information. The most restrictive provisions on either label will apply. Do not use the tank mix if any restrictions on the "Stinger" or "Curtail" labels conflict with instructions on the AGILITY™ SG label.

With "Puma"

AGILITY™ SG at 1.6 ounces per acre can be tank mixed with "Puma" herbicide for use in wheat and barley for control of green foxtail, foxtail millets, volunteer corn, and light infestations of broadleaf weeds on WEEDS CONTROLLED list. Refer to the "Puma" label for information regarding use restrictions, labeled crops, rotational cropping intervals, sprayer cleanup, use precautions and other information. The most restrictive provisions on either label will apply. Do not use the tank mix if any restrictions on the "Puma" label conflict with instructions on the AGILITY™ SG label.

With "Discover" NG

AGILITY™ SG at 1.6 -2.4 ounces per acre can be tank mixed with "Discover" NG herbicide for improved control of weeds in wheat. Refer to the "Discover" NG label for information regarding use restrictions, labeled crops, rotational cropping intervals, sprayer cleanup, use precautions and other information. The most restrictive provisions on either label will apply. Do not use the tank mix if any restrictions on the "Discover" NG label conflict with instructions on the AGILITY™ SG label.

With "Everest"

AGILITY™ SG can be tank mixed with "Everest" herbicide for improved control of weeds in wheat. Refer to the "Everest" label for information regarding use restrictions, labeled crops, rotational cropping intervals, sprayer cleanup, use precautions and other information. The most restrictive provisions on either label will apply. Do not use the tank mix if any restrictions on the "Everest" label conflict with instructions on the AGILITY™ SG label.

With Other Herbicides

AGILITY™ SG may be tank mixed with other suitable registered cereal or fallow herbicides to control weeds listed as suppressed, weeds resistant to AGILITY™ SG, or weeds not listed under Weeds Controlled. Read and follow all manufacturer's label instructions for the companion herbicide. If those instructions conflict with this label, do not tank mix the herbicide with AGILITY™ SG. Tank mixes of AGILITY™ SG plus metribuzin may result in reduced control of wild garlic.

Do not tank mix AGILITY™ SG with 'Hoelon' 3EC, because grass control may be reduced.

Do not tank mix AGILITY™ SG with DuPont™ ALLY®, ALLY® EXTRA, or other products containing metsulfuron methyl.

Do not tank mix AGILITY™ SG with other dicamba containing products (such as Clarity or Banvel) for use on cereal crops.

With Fungicides

AGILITY™ SG may be tank mixed or used sequentially with fungicides registered for use on cereal crops.

With Insecticides

Do not use AGILITY™ SG plus Malathion or Lorsban as crop injury may result.

AGILITY™ SG may be tank mixed or used sequentially with insecticides registered for use on cereal crops. However, under certain conditions (drought stress, or if the crop is in the 2-4 leaf stage), tank mixes or sequential applications of AGILITY™ SG with organophosphate insecticides (such as parathion) may produce temporary crop yellowing or, in severe cases, crop injury. The potential for crop injury is greatest when wide fluctuations in day/night temperatures occur just prior to or soon after application. Test these mixtures in a small area before treating large areas.

Do not apply AGILITY™ SG within 60 days of crop emergence where an organophosphate insecticide has been applied as an in-furrow treatment, because crop injury may result.

With Liquid Nitrogen Fertilizer Solution

Liquid nitrogen fertilizer solutions (e.g., 28-0-0, 32-0-0) may be used as a carrier in place of water. Run a tank mix compatibility test before mixing AGILITY™ SG in fertilizer solution.

AGILITY™ SG must first be dissolved with water and then added to liquid nitrogen solutions. Ensure that the agitator is running while the AGILITY™ SG is added. Use of this mixture may result in temporary crop yellowing and stunting.

If using low rates of liquid nitrogen fertilizer in the spray solution (less than 50% of the spray solution volume), the addition of surfactant is necessary. Add surfactant at 1/2 pt -1 qt per 100 gal of spray solution (0.06 -0.25% v/v) based on local guidance.

When using high rates of liquid nitrogen fertilizer in the spray solution, adding surfactant increases the risk of crop injury. If 2,4-D or MCPA is included with AGILITY™ SG and fertilizer mixture, ester formulations tend to be more compatible (See manufacturer's label). Additional surfactant may not be needed when using AGILITY™ SG in tank mix with 2,4-D ester or MCPA ester and liquid nitrogen fertilizer solutions. Consult your agricultural dealer, consultant, field advisor, or DuPont representative for specific instructions before adding an adjuvant to these tank mixtures.

Note: In certain areas east of the Mississippi river unacceptable crop response may occur with use of straight or dilute nitrogen fertilizer carrier solutions where cold temperatures or widely fluctuating day/night temperatures exist. In these areas consult your agricultural dealer, consultant, field advisor, or DuPont representative for specific instructions before using nitrogen fertilizer carrier solutions.

Do not use low rates of liquid fertilizer as a substitute for a surfactant.

Do not use with liquid fertilizer solutions with a pH less than 3.0.

SPECIFIC WEED PROBLEMS

Note: Thorough spray coverage of all weed species listed below is very important.

Blue Mustard: For best results, use 2.4-3.2 ounces per acre and apply AGILITY™ SG in tank mixtures with 2,4-D or MCPA postemergence to mustards, but before bloom (refer to Tank Mixtures section of this label for additional details).

Flixweed: For best results, use 2.4-3.2 ounces per acre and apply DuPont™ AGILITY™ SG in tank mixtures with 2,4-D or MCPA postemergence, but before bloom (refer to Tank Mixtures section of this label for additional details).

Canada Thistle: For best results, use 3.2 ounces per acre and apply AGILITY™ SG plus 2,4-D, or MCPA (refer to Tank Mixtures for additional details) in the spring after the majority of thistles have emerged and are small (rosette stage to 6" elongating stems) and actively growing. The application will inhibit the ability of emerged thistles to compete with the crop.

Corn Gromwell: For best results, use 2.4-3.2 ounces per acre and apply AGILITY™ SG when weeds are actively growing, are no larger than 2" tall, and when crop canopy will allow thorough coverage. Tank mixing 2,4-D, MCPA, or bromoxynil containing products (such as "Buctril", "Bronate", "Bison" or "Bronate Advanced") with AGILITY™ SG usually improves results (refer to Tank Mixtures section of this label for additional details).

Kochia, Prickly Lettuce, and Russian Thistle: Naturally occurring resistant biotypes of these weeds are known to occur. For best results, use AGILITY™ SG at 2.4 to 3.2 ounces per acre in a tank mix with 2,4-D and apply in the spring when weeds are less than 2" tall or wide and are actively growing (refer to Tank Mixtures section of this label for additional details).

Vetch (common and hairy): For best results, use 3.2 ounces per acre and apply AGILITY™ SG when vetch is less than 6" in length. For severe infestations of vetch, or when vetch is greater than 6" in length, use AGILITY™ SG in combination with 2,4-D, or MCPA (refer to Tank Mixtures section of this label for additional details).

Wild garlic: For best results, use 3.2 ounces per acre and apply AGILITY™ SG when wild garlic plants are less than 12" tall with 2" to 4" of new growth. Plants hardened-off by cold weather and/or drought stress may be more difficult to control. Thorough spray coverage of all garlic plants is essential. Typical symptoms of dying garlic plants may not be noticeable for 2 to 5 weeks.

Control will be improved by using AGILITY™ SG in combination with 2,4-D or MCPA (refer to Tank Mixtures section of this label for additional details).

Wild radish: For best results, use 3.2 ounces per acre applied in the fall to wild radish rosettes less than 6" in diameter and before plants harden-off. Alternatively, AGILITY™ SG can be applied in the spring for control of wild radish. Control will be improved by using AGILITY™ SG in combination with 2,4-D or MCPA (refer to Tank Mixtures section of this label for additional details) when wild radish rosettes are less than 6" in diameter. Applications made later than 30 days after weed emergence, either in the fall or spring, will result in partial control.

APPLICATION INFORMATION

PRODUCT MEASUREMENT

AGILITY™ SG can be measured using the AGILITY™ SG volumetric measuring cylinder included in the case. The degree of accuracy of this cylinder varies by ± 7.5%. For more precise measurement, use scales calibrated in ounces.

SPRAY ADJUVANTS

Always include a spray adjuvant with applications of AGILITY™ SG. In addition to a spray adjuvant, an ammonium nitrogen fertilizer may be used. Antifoaming agents may be needed.

Consult your Ag dealer or applicator, local DuPont fact sheets, technical bulletins, and service policies prior to using an adjuvant system. If another herbicide is tank mixed with AGILITY™ SG, select adjuvants authorized for use with both products. Products must contain only EPA-exempt ingredients (40CFR 1001).

Nonionic Surfactant (NIS)

- Apply 0.125 to 0.50% volume/volume (1 pt to 4 pt per 100 gal of spray solution).
- Surfactant products must contain at least 60% nonionic surfactant with a hydrophilic/lipophilic balance (HLB) greater than 12. – See the Tank Mixtures section of this label for additional information.

Crop Oil Concentrate (COC) or Modified Seed Oil (MSO)

- Only use adjuvants that contain COC or MSO in fallow applications. Do not use for postemergence applications in cereals.
- Apply at 1% volume/volume (1 gal per 100 gal spray solution) or 2% volume/volume under arid conditions.
- Oil adjuvants must contain at least 80% high quality, petroleum (mineral) or modified vegetable seed oil with at least 15% surfactant emulsifiers.
- Combination adjuvant products may be used at doses that provide the required amount of NIS, COC, MSO and/or ammonium nitrogen fertilizer. Consult product literature for use rates and restrictions.
- In addition to the adjuvants specified above, other adjuvant types may be used if they provide the same functionality and have been evaluated and approved by DuPont product management. Consult separate DuPont technical bulletins for detailed information before using adjuvant types not specified on this label.

Ammonium Nitrogen Fertilizer

- Use 2 qt/acre of a high-quality urea ammonium nitrate (UAN), such as 28%N or 32%N, or 2 lb/acre of a spray-grade ammonium sulfate (AMS). Use 4 qt/acre UAN or 4 lb/acre AMS under arid conditions.

GROUND APPLICATION

For optimum spray distribution and thorough coverage, use flat-fan or low-volume flood nozzles.

For flat-fan nozzles, use a spray volume of at least 5 gal per acre (GPA).

For flood nozzles on 30" spacing, use flood nozzles no larger than TK10 (or the equivalent), a pressure of at least 30 psi and a spray volume of at least 10 GPA only. For 40" nozzle spacing, use at least 13 GPA; for 60" spacing use at least 20 GPA. It is essential to overlap the nozzles 100% for all spacings.

"Raindrop RA" nozzles are not recommended for AGILITY™ SG applications, because weed control performance may be reduced.

Use screens that are 50-mesh or larger.

AERIAL APPLICATION

Use nozzle types and arrangements that provide optimum spray distribution and maximum coverage at 1 to 5 GPA. Use at least 3 GPA in Idaho, Oregon, Washington, or Utah.

When applying AGILITY™ SG by air in areas near sensitive crops, use solid-stream nozzles oriented straight back. Adjust swath to avoid spray drift damage to downwind sensitive crops

and/or use ground equipment to treat border edge of field. See the **Spray Drift Management** section of this label.

For aerial application in Washington, follow the directions in the Spray Drift Management Section of this label and the following Washington state restrictions:

Applications of DuPont™ AGILITY™ SG must be made in equipment that meets the most restrictive Washington Agricultural Codes (WAC) for the prevention of herbicide drift for the respective county.

Do not apply in equipment that does not meet these WAC standards.

SEQUENTIAL APPLICATIONS

AGILITY™ SG can be applied either before or after applications of other products registered for use in wheat, barley, triticale or fallow. Read and follow all label instructions on timing, precautions, and warnings for these herbicides before using these in sequence with AGILITY™ SG. If those instructions conflict with this label, do not use that product in sequence with AGILITY™ SG.

- DuPont™ ALLY®, ALLY® EXTRA, and other products containing metsulfuron methyl should not be used as a sequential treatment with AGILITY™ SG.
- If using DuPont™ HARMONY® EXTRA as a sequential treatment with AGILITY™ SG, do not exceed 0.7 ounce of HARMONY® EXTRA per acre per crop season.
- If using DuPont™ EXPRESS® as a sequential treatment with AGILITY™ SG, do not exceed 0.25 ounce of EXPRESS® per acre per crop season.
- If using products that contain dicamba as a sequential treatment with AGILITY™ SG, do not exceed the following amounts of acid equivalent dicamba per acre per season: 8.4 ounces in wheat, 5.6 ounces in barley, 2.1 ounces in triticale, and 16.8 ounces in fallow. Do not exceed 2.0 pounds acid equivalent dicamba per acre per year.

MIXING INSTRUCTIONS

1. Fill the tank 1/4 to 1/3 full of water.
2. While agitating, add the required amount of AGILITY™ SG.
3. Continue agitation until the AGILITY™ SG is fully dissolved, at least 5 minutes.
4. Once the AGILITY™ SG is fully dissolved, maintain agitation and continue filling tank with water. AGILITY™ SG should be thoroughly dissolved with water before adding any other material.
5. As the tank is filling, add tank mix partners (if desired) then add the required volume of surfactant. Always add surfactant last. Antifoaming agents may be used. Do not use with spray additives that alter the pH of the spray solution below pH 5.0 or above pH 9.0 as rapid product degradation can occur. Spray solutions of pH 6.0 - 8.0 allow for optimum stability of AGILITY™ SG.
6. Dispersed tank mix partners can settle if the tank mixture is not continually agitated.
7. Apply AGILITY™ SG spray mixture within 24 hours of mixing to avoid product degradation.

8. If AGILITY™ SG and a tank mix partner are to be applied in multiple loads, fully dissolve the AGILITY™ SG in clean water prior to adding to the tank. This will prevent the tank mix partner from interfering with the dissolution of the AGILITY™ SG.

SPRAY EQUIPMENT

For specific application equipment, refer to the manufacturer's instructions for additional information on GPA, pressure, speed, nozzle types and arrangements, nozzle heights above the target canopy, etc.

Be sure to calibrate air or ground equipment properly before application. Select a spray volume and delivery system that will ensure thorough coverage and a uniform spray pattern with minimum drift. Use higher spray volumes to obtain better coverage when crop canopy is dense. Avoid swath overlapping, and shut off spray booms while starting, turning, slowing, or stopping, to avoid injury to the crop.

Do not make applications using equipment and/or spray volumes or during weather conditions that might cause spray to drift onto nontarget sites. For additional information on spray drift refer to Spray Drift Management section of label.

Sprayer Precaution

The spray equipment must be cleaned before AGILITY™ SG is sprayed. Follow the cleanup procedures specified on the labels of the previously applied products.

AT THE END OF THE DAY

It is recommended that during periods when multiple loads of AGILITY™ SG herbicide are applied, at the end of each day of spraying the interior of the tank be rinsed with fresh water and then partially filled, and the boom and hoses flushed. This will prevent the buildup of dried pesticide deposits which can accumulate in the application equipment.

AFTER SPRAYING AGILITY™ SG AND BEFORE SPRAYING CROPS OTHER THAN WHEAT, BARLEY OR TRITICALE

To avoid subsequent injury to desirable crops, thoroughly clean all mixing and spray equipment immediately following applications of AGILITY™ SG as follows:

1. Empty the tank and drain the sump completely.
2. Spray the tank walls with clean water using a minimum volume of 10% of the tank volume. Circulate the water through the lines, including all by-pass lines, for at least two minutes. Flush the boom well and empty the sprayer. Completely drain the sump.
3. Fill the tank with water while adding 1 quart of household ammonia for every 25 gallons of water. Operate the pump to circulate the ammonia solution through the sprayer system for 15 to 20 minutes and discharge a small amount of the ammonia solution through the boom and nozzles. Let the solution stand for several hours, preferably overnight. Flush the boom well and empty the sprayer. Completely drain the sump.
4. Remove the nozzles and screens and clean separately in a bucket containing water.

The rinsate solution may be applied to the crop(s) specified on this label. Do not exceed the maximum-labeled use rate. If cleaners are used, consult the cleaner label for rinsate

disposal instructions. If no instructions are given, dispose of the rinsate on site or at an approved waste disposal facility.

Notes:

1. Always start with a clean spray tank.
 2. Steam-cleaning aerial spray tanks is recommended to facilitate the removal of any caked deposits.
 3. When DuPont™ AGILITY™ SG is tank mixed with other pesticides, all cleanout procedures for each product should be examined and the most rigorous procedure should be followed.
 4. In addition to this cleanout procedure, all pre-cleanout guidelines on subsequently applied products should be followed as per the individual labels.
- If using products that contain dicamba as a sequential treatment with AGILITY™ SG, do not exceed the following amounts of acid equivalent dicamba per acre per season: 8.4 ounces in wheat, 5.6 ounces in barley, 2.1 ounces in triticale, and 16.8 ounces in fallow.

CROP ROTATION

Before using AGILITY™ SG carefully consider your crop rotation plans and options. For rotational flexibility, do not treat all of your acres at the same time.

MINIMUM ROTATIONAL INTERVALS

Minimum rotation intervals* are determined by the rate of breakdown of AGILITY™ SG applied. AGILITY™ SG breakdown in the soil is affected by soil pH, presence of soil microorganisms, soil temperature, and soil moisture. Low soil pH, high soil temperature, and high soil moisture increase AGILITY™ SG breakdown in soil, while high soil pH, low soil temperature, and low soil moisture slow AGILITY™ SG breakdown.

Of these 3 factors, only soil pH remains relatively constant. Soil temperature, and to a greater extent, soil moisture, can vary significantly from year to year and from area to area. For this reason, soil temperatures and soil moisture should be monitored regularly when considering crop rotations.

* The minimum rotation interval represents the period of time from the last application to the anticipated date of the next planting. **Minimum rotation intervals must be extended 1 crop season if drought conditions prevail after application and before the rotational crop is planted.**

Soil pH Limitations

AGILITY™ SG should not be used on soils having a pH above 7.9, because extended soil residual activity could extend crop rotation intervals beyond normal. Under certain conditions, AGILITY™ SG could remain in the soil for 34 months or more, injuring wheat, barley or triticale. In addition, other crops planted in high-pH soils can be extremely sensitive to low concentrations of AGILITY™ SG.

Checking Soil pH

Before using AGILITY™ SG, determine the soil pH of the areas of intended use. To obtain a representative pH value for the test area, take several 0” to 4” samples from different areas of the field and analyze them separately. Consult local extension publications for additional information on recommended soil sampling procedures.

FIELD BIOASSAY

A field bioassay is necessary if crops other than wheat, barley or those listed on this label are to be planted on land previously treated with AGILITY™ SG. To conduct a field bioassay, grow test strips of the crop or crops you plan to grow the following year in fields previously treated with AGILITY™ SG. Crop response to the bioassay will indicate whether or not to rotate to the crop(s) grown in the test strips. If a field bioassay is planned, check with your local DuPont representative for information detailing field bioassay procedure.

GRAZING

Allow at least 7 days between application and grazing of treated forage. In addition, allow at least 7 days between application and feeding of forage from treated areas to livestock. Allow at least 30 days between application and feeding of hay from treated areas to livestock. Harvested straw may be used for bedding and/or feed. Allow at least 45 days between application and harvesting of grain.

CROP ROTATION TABLES

See next five pages for specific crop rotation guidance based on AGILITY™ SG use rate, location, soil pH, cumulative precipitation, and crop.

12
21

All Areas - Following Use of DuPont™ AGILITY™ SG at 1.6 to 3.2 Ounces Per Acre

Crop	Soil pH	Minimum Cumulative Precipitation (inches)	Minimum Rotation Interval (months)
Winter wheat, spring wheat and Triticale	7.9 or lower	No restrictions	1
Durum wheat, barley, spring/winter oat	7.9 or lower	No restrictions	10

Rotation Intervals For Crops in Non-Irrigated Land Following Use of AGILITY™ SG at 1.6 to 3.2 Ounces Per Acre on Wheat, Barley, Triticale or Fallow

Location		Crop	Soil pH	Minimum Cumulative Precipitation (inches)	Minimum Rotation Interval (months)
State	County or Area				
Colorado	Statewide	Grain sorghum	7.9 or lower	No restrictions	4
		Flax, Safflower	7.9 or lower	No restrictions	22
	Generally N of I-70	Field corn	7.9 or lower	15	12
	Statewide	STST™ Soybean	7.9 or lower	No restrictions	4
		IR Corn	7.9 or lower	No restrictions	4
		Proso millet	7.9 or lower	No restrictions	4
Idaho	Southern Idaho	Flax, Safflower	7.9 or lower	No restrictions	22
	Statewide	Peas, Lentils, Canola	6.8 or lower	18	10
		Peas	6.9 to 7.9	18	15
		Lentils	6.9 to 7.9	18	34
		Canola	6.9 to 7.9	18	22
		Condiment mustard	7.3 or lower	10	10
		Chickpeas (Garbanzo beans)	7.3 or lower	10	10
		Condiment mustard	7.4 or higher	28	34
		Chickpeas (Garbanzo beans)	7.4 or higher	28	34
Kansas	Statewide	STST™ Soybean	7.9 or lower	No restrictions	4
		IR Corn	7.9 or lower	No restrictions	4
		Proso millet	7.9 or lower	No restrictions	4
		Grain sorghum	7.9 or lower	No restrictions	4
		Flax, Safflower	7.9 or lower	No restrictions	22
	Central and Western Kansas (West of the Flint Hills)	Field corn	7.9 or lower	15	12
	Western Kansas W. of Hwy. 183	Soybeans	7.5 or lower	22	22
			7.6-7.9	33	34
	Central Kansas; generally E. of Hwy. 183 and W. of the Flinthills	Soybeans	7.9 or lower	15	12

Continued on next page

14
21

Rotation Intervals For Crops in Non-Irrigated Land (continued)
Following Use of DuPont™ AGILITY™ SG at 1.6 to 3.2 Ounces Per Acre on Wheat,
Barley, Triticale or Fallow

Location		Crop	Soil pH	Minimum Cumulative Precipitation (inches)	Minimum Rotation Interval (months)
State	County or Area				
Montana	Statewide	Grain sorghum, Proso millet, Field corn	7.9 or lower	22	22
		Alfalfa (hay only)	7.6-7.9	No restrictions	34
			7.5 or lower	No restrictions	22
		Flax, Safflower	7.9 or lower	No restrictions	22
Nebraska	Statewide	STS™ Soybean	7.9 or lower	No restrictions	4
		IR Corn	7.9 or lower	No restrictions	4
		Proso millet	7.9 or lower	No restrictions	4
		Grain sorghum	7.9 or lower	No restrictions	4
		Flax, Safflower	7.9 or lower	No restrictions	22
	Generally W. of Hwy. 77 and E. of the Panhandle	Field corn	7.9 or lower	15	12
Soybeans		7.5 or lower	22	22	
		7.6-7.9	33	34	
New Mexico	Statewide	Grain sorghum, Proso millet	7.9 or lower	No restrictions	10
		Flax, Safflower	7.9 or lower	No restrictions	22
	Eastern New Mexico	Cotton (dryland only)	7.9 or lower	30	22
North Dakota	W. of Hwy. 1	Grain sorghum, Proso millet, Field corn, Dry beans, Flax, Safflower	7.9 or lower	22	22
	E. of Hwy. 1	Grain sorghum, Proso millet, Field corn, Dry beans, Flax, Safflower	7.9 or lower	34	34
Oklahoma	Statewide	STS™ Soybean	7.9 or lower	No restrictions	4
		IR Corn	7.9 or lower	No restrictions	4
		Proso millet	7.9 or lower	No restrictions	4
		Grain sorghum	7.9 or lower	No restrictions	4
		Flax, Safflower	7.9 or lower	No restrictions	22
		Field corn	7.9 or lower	15	12
	Panhandle	Cotton (dryland only)	7.9 or lower	30	22
	E. of the Panhandle	Cotton (dryland only)	7.9 or lower	25	14

Continued on next page

15
21

Rotation Intervals For Crops in Non-Irrigated Land (continued)
Following Use of DuPont™ AGILITY™ SG at 1.6 to 3.2 Ounces Per Acre on Wheat, Barley, Triticale or Fallow

Location		Crop	Soil pH	Minimum Cumulative Precipitation (inches)	Minimum Rotation Interval (months)
State	County or Area				
Oregon	Statewide	Peas Lentils Canola	6.8 or lower	18	10
		Peas	6.9 to 7.9	18	15
		Lentils	6.9 to 7.9	18	34
		Canola	6.9 to 7.9	18	22
		Condiment mustard	7.3 or lower	10	10
		Chickpeas (Garbanzo beans)	7.3 or lower	10	10
		Condiment mustard	7.4 or higher	28	34
		Chickpeas (Garbanzo beans)	7.4 or higher	28	34
South Dakota	Statewide	Flax, Safflower	7.9 or lower	No restrictions	22
	S. of Hwy. 212 & E. of the Missouri River, & S. of Hwy. 34 & W. of Missouri River	Grain sorghum, Proso millet	7.9 or lower	13	12
	Generally E. of Missouri River & S. of Hwy. 14, & W. of Missouri River	Field corn	7.9 or lower	15	12
Texas	Statewide	STS™ Soybeans	7.9 or lower	No restrictions	4
		IR Corn	7.9 or lower	No restrictions	4
		Proso millet	7.9 or lower	No restrictions	4
		Grain sorghum	7.9 or lower	No restrictions	4
		Flax, Safflower	7.9 or lower	No restrictions	22
	Panhandle	Field corn	7.9 or lower	15	12
		Cotton (dryland only)	7.9 or lower	30	22
	N. Central Texas*	Field corn	7.9 or lower	15	12
		Cotton (dryland only)	7.9 or lower	25	14
* The counties of N. Central Texas are: Archer, Baylor, Bell, Bosque, Bowie, Callahan, Camp, Cass, Clay, Collin, Cooke, Coryell, Dallas, Delta, Denton, Eastland, Ellis, Falls, Fannin, Foard, Franklin, Grayson, Hardeman, Haskell, Hill, Hood, Hopkins, Hunt, Jack, Johnson, Kaufman, Knox, Lamar, Limestone, McLennan, Milam, Montague, Morris, Nafarro, Palo Pinto, Parker, Rains, Red River, Robertson, Rockwall, Shackelford, Somervell, Stephens, Tarrant, Throckmorton, Titus, Upshur, Van Zandt, Wilbarger, Wichita, Williamson, Wise, Wood, Young.					
Utah	Statewide	Flax, Safflower	7.9 or lower	No restrictions	22

Continued on next page

16
21

Rotation Intervals For Crops in Non-Irrigated Land (continued)
Following Use of DuPont™AGILITY™ SG at 1.6 to 3.2 Ounces Per Acre on Wheat, Barley, Triticale or Fallow

Location		Crop	Soil pH	Minimum Cumulative Precipitation (inches)	Minimum Rotation Interval (months)
State	County or Area				
Washington	Statewide	Condiment mustard	7.3 or lower	10	10
		Chickpeas (Garbanzo beans)	7.3 or lower	10	10
		Condiment mustard	7.4 or higher	28	34
		Chickpeas (Garbanzo beans)	7.4 or higher	28	34
		Peas Lentils Canola	6.8 or lower	18	10
		Peas	6.9 to 7.9	18	15
		Lentils	6.9 to 7.9	18	34
		Canola	6.9 to 7.9	18	22
Wyoming	Statewide	Flax, Safflower	7.9 or lower	No restrictions	22
	Southern Wyoming	Grain sorghum, Proso millet	7.9 or lower	No restrictions	10
	Southern Wyoming (Goshen, Laramie, and Platte counties only)	Field corn	7.9 or lower	15	12
	Northern Wyoming	Grain sorghum, Proso millet, Field corn	7.9 or lower	22	22

Rotation Intervals for crops not covered above - The minimum rotation interval is 34 months with at least 28" of cumulative precipitation during the period:

- to any major field crop not listed (See the Rotation Intervals table)
- if the soil pH is not in the specified range
- if the use rate applied is not specified in the table
- or if the minimum cumulative precipitation has not occurred since application.

To rotate to a major field crop at an interval shorter than specified, a field bioassay must be successfully completed to that crop. A field bioassay must be successfully completed before rotation to any minor crops (as determined by the USDA criteria). See section on Field Bioassay for further information.

Rotation Intervals For Crops in Non-Irrigated Land
Following Use of AGILITY™ SG up to 2.4 Ounces Per Acre on Wheat, Barley, Triticale or Fallow in the states of Colorado, Kansas, Nebraska, New Mexico, Oklahoma, South Dakota, Texas and Wyoming

Crop	Soil pH	Minimum Cumulative Precipitation (inches)	Minimum Rotation Interval (months)
Sunflower	7.9 or lower	No restrictions	10

Rotation Intervals for crops not covered above (up to 2.4 ounces per acre) - The minimum rotation interval is 34 months with at least 28" of cumulative precipitation during the period:

- to any major field crop not listed (See the Rotation Intervals table)
- if the soil pH is not in the specified range
- if the use rate applied is not specified in the table
- or if the minimum cumulative precipitation has not occurred since application.

To rotate to a major field crop at an interval shorter than specified, a field bioassay must be successfully completed to that crop. A field bioassay must be successfully completed before rotation to any minor crops (as determined by the USDA criteria). See section on Field Bioassay for further information.

17
21

**Rotation Intervals For Crops in Non-Irrigated Land
Following Use of DuPont™ AGILITY™ SG at 2.4 to 3.6 Ounces Per Acre on Wheat,
Barley, Triticale or Fallow**

Location		Crop	Soil pH	Minimum Cumulative Precipitation (inches)	Minimum Rotation Interval (months)
State	County or Area				
Colorado Idaho Kansas Montana Nebraska New Mexico Oklahoma South Dakota Texas Utah Wyoming	Statewide	Sunflower	7.9 or lower	No restrictions	22
North Dakota	W. of Hwy. 1	Sunflower	7.9 or lower	22	22
	E. of Hwy. 1	Sunflower	7.9 or lower	34	34

Rotation Intervals for crops not covered above (2.4 to 3.2 ounces per acre) - The minimum rotation interval is 34 months with at least 28" of cumulative precipitation during the period:

- to any major field crop not listed (See the Rotation Intervals table)
- if the soil pH is not in the specified range
- if the use rate applied is not specified in the table
- or if the minimum cumulative precipitation has not occurred since application.

To rotate to a major field crop at an interval shorter than specified, a field bioassay must be successfully completed to that crop. A field bioassay must be successfully completed before rotation to any minor crops (as determined by the USDA criteria). See section on Field Bioassay for further information.

**Rotation Intervals For Crops in Non-Irrigated Land
Following Use of AGILITY™ SG at 1.6 Ounces Per Acre on Wheat, Barley Triticale or Fallow**

Crop	Soil pH	Minimum Cumulative Precipitation (inches)	Minimum Rotation Interval (months)
Sorghum, Grain	7.9 or lower	No restrictions	4
Cotton	7.9 or lower	No restrictions	10
Safflower	7.9 or lower	No restrictions	10
Peas, Dry /Green	6.8 or lower	No restrictions	10
	6.9 to 7.9	No restrictions	22
Lentils	6.8 or lower	No restrictions	10
	6.9 to 7.9	No restrictions	22
Alfalfa	6.8 or lower	No restrictions	10
	6.9 to 7.9	No restrictions	22
Beans, Dry	6.8 or lower	No restrictions	10
	6.9 to 7.9	No restrictions	22
Sunflower	7.9 or lower	No restrictions	10

Rotation Intervals for crops not covered above (1.6 ounces per acre) - The minimum rotation interval is 22 months with at least 18" of cumulative precipitation during the period:

- to any major field crop not listed (See the Rotation Intervals table)
- if the soil pH is not in the specified range
- if the use rate applied is not specified in the table
- or if the minimum cumulative precipitation has not occurred since application.

To rotate to a major field crop at an interval shorter than specified, a field bioassay must be successfully completed to that crop. A field bioassay must be successfully completed before rotation to any minor crops (as determined by the USDA criteria). See section on Field Bioassay for further information.

SPRAY DRIFT MANAGEMENT

The interaction of many equipment and weather-related factors determines the potential for spray drift. The applicator is responsible for considering all these factors when making application decisions.

AVOIDING SPRAY DRIFT IS THE RESPONSIBILITY OF THE APPLICATOR.

Ground or aerial application equipment which will give good spray coverage of weed foliage should be used. **HOWEVER, DO NOT USE AERIAL APPLICATION EQUIPMENT IF SPRAY PARTICLES CAN BE CARRIED BY WIND INTO AREAS WHERE SENSITIVE CROPS OR PLANTS ARE GROWING.**

Apply 3 to 50 gallons of diluted spray per treated acre when using ground application equipment, or 2 to 10 gallons of diluted spray per treated acre when using aerial application equipment. Use the higher level of the listed spray volumes when treating dense or tall vegetation. Use coarse sprays.

Select nozzles designed to produce minimal amounts of fine spray particles. Spray with nozzles as close to the weeds as is practical for good weed coverage.

DuPont™ AGILITY™ SG should not be applied during periods of gusty wind or when wind is in excess of 15 mph as uneven spray coverage may occur,

Avoid disturbing (e.g. cultivating or mowing) treated areas for at least 7 days following application.

IMPORTANCE OF DROPLET SIZE

The most effective way to reduce drift potential is to apply large droplets. The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. The presence of sensitive species nearby, the environmental conditions, and pest pressure may affect how an applicator balances drift control and coverage. **APPLYING LARGER DROPLETS REDUCES DRIFT POTENTIAL, BUT WILL NOT PREVENT DRIFT IF APPLICATIONS ARE MADE IMPROPERLY OR UNDER UNFAVORABLE ENVIRONMENTAL CONDITIONS!** See **Wind, Temperature and Humidity, and Temperature Inversions** sections of this label.

Controlling Droplet Size - General Techniques

- **Volume** - Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets.
- **Pressure** - Use the lower spray pressures recommended for the nozzle. Higher pressure reduces droplet size and does not improve canopy penetration. **WHEN HIGHER FLOW RATES ARE NEEDED, USE A HIGHER-CAPACITY NOZZLE INSTEAD OF INCREASING PRESSURE.**
- **Nozzle Type** - Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles.

Controlling Droplet Size - Aircraft

- **Number of Nozzles** - Use the minimum number of nozzles with the highest flow rate that provide uniform coverage.
- **Nozzle Orientation** - Orienting nozzles so that the spray is emitted backwards, parallel to the airstream will produce larger droplets than other orientations.

- **Nozzle Type** - Solid stream nozzles (such as disc and core with swirl plate removed) oriented straight back produce larger droplets than other nozzle types.
- **Boom Length** - The boom length should not exceed 3/4 of the wing or rotor length - longer booms increase drift potential.
- **Application Height** - Application more than 10 ft above the canopy increases the potential for spray drift.

BOOM HEIGHT

Setting the boom at the lowest referenced height (if specified) which provides uniform coverage reduces the exposure of droplets to evaporation and wind. For ground equipment, the boom should remain level with the crop and have minimal bounce.

WIND

Drift potential increases at wind speeds of less than 3 mph (due to inversion potential) or more than 10 mph. However, many factors, including droplet size and equipment type determine drift potential at any given wind speed. **AVOID GUSTY AND WINDLESS CONDITIONS.**

Note: Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they affect spray drift.

TEMPERATURE AND HUMIDITY

When making applications in hot and dry conditions, set up equipment to produce larger droplets to reduce effects of evaporation.

TEMPERATURE INVERSIONS

Drift potential is high during a temperature inversion. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain close to the ground and move laterally in a concentrated cloud. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

SHIELDED SPRAYERS

Shielding the boom or individual nozzles can reduce the effects of wind. However, it is the responsibility of the applicator to verify that the shields are preventing drift and not interfering with uniform deposition of the product.

AIR ASSISTED (AIR BLAST) FIELD CROP SPRAYERS

Air assisted field crop sprayers carry droplets to the target via a downward directed air stream. Some may reduce the potential for drift, but if a sprayer is unsuitable for the application and/or set up improperly, high drift potential can result. It is the responsibility of the applicator to determine that a sprayer is suitable for the intended application, is configured properly, and that drift is not occurring.

RESISTANCE

When herbicides that affect the same biological site of action are used repeatedly over several years to control the same weed species in the same field, naturally-occurring resistant biotypes may survive a correctly applied herbicide treatment, propagate, and become dominant in that field. Adequate control of these resistant weed biotypes cannot be expected. If weed control is unsatisfactory, it may be necessary to retreat the problem area using a product affecting a different site of action.

To better manage herbicide resistance through delaying the proliferation and possible dominance of herbicide resistant weed biotypes, it may be necessary to change cultural practices within and between crop seasons such as using a combination of tillage, retreatment, tank-mix partners and/or sequential herbicide applications that have a different site of action. Weed escapes that are allowed to go to seed will promote the spread of resistant biotypes. See the Weeds Controlled section of this label for additional information on managing herbicide resistant weed biotypes.

It is advisable to keep accurate records of pesticides applied to individual fields to help obtain information on the spread and dispersal of resistant biotypes. Consult your agricultural dealer, consultant, applicator, and/or appropriate state agricultural extension service representative for specific alternative cultural practices or herbicide recommendations available in your area.

INTEGRATED PEST MANAGEMENT

DuPont recommends the use of Integrated Pest Management (IPM) programs to control pests. This product may be used as part of an Integrated Pest Management (IPM) program that can include biological, cultural, and genetic practices aimed at preventing economic pest damage. Application of this product should be based on IPM principles and practices including field scouting or other detection methods, correct target pest identification, population monitoring, and treating when target pest populations reach locally determined action thresholds. Consult your state cooperative extension service, professional consultants or other qualified authorities to determine appropriate action treatment threshold levels for treating specific pest/crop systems in your area.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Pesticide Storage: Store product in original container only.

Pesticide Disposal: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

Container Handling: Refer to the Net Contents section of this product's labeling for the applicable "Nonrefillable Container" or "Refillable Container" designation.

Nonrefillable Plastic and Metal Containers (Capacity Equal to or Less Than 50 Pounds): Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then, for Plastic Containers, offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration. Do not burn, unless allowed by state and local ordinances. For Metal Containers, offer for recycling if available or reconditioning if appropriate, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

Nonrefillable Plastic and Metal Containers (Capacity Greater Than 50 Pounds): Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then, for Plastic Containers, offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration. Do not burn, unless allowed by state and local ordinances. For Metal Containers, offer for recycling if available or reconditioning if appropriate, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

19
21

Nonrefillable Plastic and Metal Containers, e.g., Intermediate Bulk Containers [IBC] (Size or Shape Too Large to be Tipped, Rolled or Turned Upside Down):

Nonrefillable container. Do not reuse or refill this container. Clean container promptly after emptying the contents from this container into application equipment or mix tank and before final disposal using the following pressure rinsing procedure. Insert a lance fitted with a suitable tank cleaning nozzle into the container and ensure that the water spray thoroughly covers the top, bottom and all sides inside the container. The nozzle manufacturer generally provides instructions for the appropriate spray pressure, spray duration and/or spray volume. If the manufacturer's instructions are not available, pressure rinse the container for at least 60 seconds using a minimum pressure of 30 PSI with a minimum rinse volume of 10% of the container volume. Drain, pour or pump rinsate into application equipment or rinsate collection system. Repeat this pressure rinsing procedure two more times. Then, for Plastic Containers, offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration. For Metal Containers, offer for recycling if available or reconditioning if appropriate, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

Nonrefillable Paper or Plastic Bags, Fiber Sacks including Flexible Intermediate Bulk Containers (FIBC) or Fiber Drums With Liners:

Nonrefillable container. Do not reuse or refill this container. Completely empty paper or plastic bag, fiber sack or drum liner by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application or manufacturing equipment. Then offer for recycling if available or dispose of empty paper or plastic bag, fiber sack or fiber drum and liner in a sanitary landfill, or by incineration. Do not burn, unless allowed by state and local ordinances.

Refillable Fiber Drums With Liners: Refillable container (fiber drum only). Refilling Fiber Drum: Refill this fiber drum with DuPont™ AGILITY™ SG herbicide (with TotalSol® soluble granules) containing dicamba, thifensulfuron methyl, tribenuron methyl and metsulfuron methyl only. Do not reuse this fiber drum for any other purpose. Cleaning before refilling is the responsibility of the refiller. Completely empty liner by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application or manufacturing equipment. Disposing of Fiber Drum and/or Liner: Do not reuse this fiber drum for any other purpose other than refilling (see preceding). Cleaning the container (liner and/or fiber drum) before final disposal is the responsibility of the person disposing of the container. Offer the liner for recycling if available or dispose of liner in a sanitary landfill, or by incineration. Do not burn, unless allowed by state and local ordinances. If drum is contaminated and cannot be reused, dispose of it in the manner required for its liner. To clean the fiber drum before final disposal, completely empty the fiber drum by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application or manufacturing equipment. Then offer the fiber drum for recycling if available or dispose of in a sanitary landfill, or by incineration. Do not burn, unless allowed by state and local ordinances.

All Other Refillable Containers:

Refillable container. Refilling Container: Refill this container with AGILITY™ SG herbicide (with TotalSol® soluble granules) containing dicamba, thifensulfuron methyl, tribenuron methyl and metsulfuron methyl only. Do not reuse this container for any other purpose. Cleaning before refilling is the responsibility of the refiller. Prior to refilling, inspect carefully for damage such as cracks, punctures, abrasions, worn out threads and closure devices. If damage is found, do not use the container, contact DuPont at the number below for instructions. Check for leaks after refilling and before transporting. If leaks are found, do not reuse or transport container, contact DuPont at the number below for instructions. Disposing of Container: Do not reuse this container for any other purpose other than refilling (see preceding). Cleaning the container before final disposal is the responsibility of the person disposing of the container. To clean the container before final disposal, use the following pressure rinsing procedure. Insert a lance fitted with a suitable tank cleaning nozzle into the container and ensure that the water spray thoroughly covers the top, bottom and all sides inside the container. The nozzle manufacturer generally provides instructions for the appropriate spray pressure, spray duration and/or spray volume. If the manufacturer's instructions are not available, pressure rinse the container for at least 60 seconds using a minimum pressure of 30 PSI with a minimum rinse volume of 10% of the container volume. Drain, pour or pump rinsate into application equipment or rinsate collection system. Repeat this pressure rinsing procedure two more times. Then, for Plastic Containers, offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration. Do not burn, unless allowed by state and local ordinances. For Metal Containers, offer for recycling if available or reconditioning if appropriate, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

Outer Foil Pouches of Water Soluble Packets (WSP):

Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available or, dispose of the empty outer foil pouch in the trash as long as WSP is unbroken. If the outer pouch contacts the formulated product in any way, the pouch must be triple rinsed with clean water. Add the rinsate to the spray tank and dispose of the outer pouch as described previously.

Do not transport if this container is damaged or leaking. If the container is damaged, leaking or obsolete, or in the event of a major spill, fire or other emergency, contact DuPont at 1-800-441-3637, day or night.

NOTICE TO BUYER: Purchase of this material does not confer any rights under patents of countries outside of the United States.

The DuPont Oval Logo, DuPont™, AGILITY™, ALLY®, HARMONY®, EXPRESS®, STS™, and TotalSol® are trademarks or registered trademarks of E. I. du Pont de Nemours & Company

"Clarity" and "Banvel" are registered trademarks of BASF
"Bronate", "Buctril", "Hoelon" and "Puma" are registered trademarks of Bayer CropScience
"Raindrop RA" is a registered trademark of GP Companies; Inc.
"Everest" is a registered trademark of Arysta Lifescience North America
"Discover" is a registered trademark of Syngenta Participations AG
"Stinger", "Curtail", and "Lorsban" are registered trademarks of Dow AgroSciences
"Bison" is a registered trademark of Agrilience, LLC

D - 1440 082510

21
21

LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read this Limitation of Warranty and Liability Before Buying or Using This Product. If the Terms Are Not Acceptable, Return the Product at Once, Unopened, and the Purchase Price Will Be Refunded.

It is impossible to eliminate all risks associated with the use of this product. Such risks arise from weather conditions, soil factors, off target movement, unconventional farming techniques, presence of other materials, the manner of use or application, or other unknown factors, all of which are beyond the control of DuPont. These risks can cause: ineffectiveness of the product, crop injury, or injury to non-target crops or plants. **WHEN YOU BUY OR USE THIS PRODUCT, YOU AGREE TO ACCEPT THESE RISKS.**

DuPont warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for the purpose stated in the Directions for Use, subject to the inherent risks described above, when used in accordance with the Directions for Use under normal conditions.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, DUPONT MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR OF MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL DUPONT OR SELLER BE LIABLE FOR ANY INCIDENTAL, CONSEQUENTIAL OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. BUYER'S OR USER'S BARGAINED-FOR EXPECTATION IS CROP PROTECTION. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER AND THE EXCLUSIVE LIABILITY OF DUPONT OR SELLER, FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY OR CONTRACT, NEGLIGENCE, TORT OR STRICT LIABILITY), WHETHER FROM FAILURE TO PERFORM OR INJURY TO CROPS OR OTHER PLANTS, AND RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT, OR AT THE ELECTION OF DUPONT OR SELLER, THE REPLACEMENT OF THE PRODUCT.

To the extent consistent with applicable law that allows such requirement, DuPont or its Ag Retailer must have prompt notice of any claim so that an immediate inspection of buyer's or user's growing crops can be made. Buyer and all users shall promptly notify DuPont or a DuPont Ag Retailer of any claims, whether based on contract, negligence, strict liability, other tort or otherwise, or be barred from any remedy.

This Limitation of Warranty and Liability may not be amended by any oral or written agreement.

For product information call: 1-888-6-DUPONT

Internet address: <http://cropprotection.dupont.com/>

© 2007-2010 E. I. du Pont de Nemours and Company, 1007 Market Street, Wilmington, Delaware 19898.

All rights reserved.