

352-757

2/25/2010

1 of 3



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

FEB 25 2010

OFFICE OF
PREVENTION, PESTICIDES AND
TOXIC SUBSTANCES

Richard J. Ambrose
E.I. DuPont de Nemours and Company
DuPont Crop Protection
Stine-Haskell Research Center
P.O. Box 30
Newark, DE 19714-0030

Subject: Supplemental Label – Add the NE and SW corners of Nebraska for soybean applications
Product Name: DuPont Enlite Herbicide
EPA Reg. No. 352-757
Application dated: January 15, 2010

Dear Mr. Ambrose:

The supplemental labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act, as amended, is acceptable.

At your next label printing, or within eighteen months of the date of this letter, whichever comes first, you must incorporate this supplemental labeling into the main product labeling.

A stamped copy of the label is enclosed for your records. Please submit one copy of your final printed label before you release the product for shipment.

Sincerely,

A handwritten signature in black ink, appearing to read "Jim Tompkins", written over a circular stamp or mark.

Jim Tompkins
Product Manager 25
Herbicide Branch
Registration Division (7505P)

203

SUPPLEMENTAL LABELING

DuPont Crop Protection

DUPONT™ ENLITE® HERBICIDE

DUPONT™ Enlite® Herbicide

EPA Reg. No. 352-757

FOR USE ON ALL pH SOILS IN THE STATE OF NEBRASKA

ENLITE® will provide burndown and residual control of labeled winter annuals, perennials, and spring annuals when applied pre-plant / preemergence to soybeans.

Geographic Use Region Addition to the Northern Region –

- Add the northeast corner of Nebraska that includes fields north of HWY 30 and east of HWY 281 and the southwest corner of Nebraska that includes fields south of HWY 30 and west of HWY 281.

The geographical use region for ENLITE® in the state of Nebraska is now defined below:

Northern Region – In Nebraska: All fields north of HWY 30 or west of HWY 281.

DIRECTIONS FOR USE

ENLITE® herbicide is a dispersible granule formulation to be mixed with water at a rate of 2.8 ounces per acre, and applied for selective burndown and limited residual weed control in soybeans. ENLITE® will control many broadleaf weeds and provide partial control of certain annual grasses.

Apply ENLITE® using one of the following application methods:

- Early pre-plant or pre-plant in conservation tillage, no-till or stale seedbed systems.
- Preemergence application must be made within 3 days after planting and prior to soybean emergence. Do not apply ENLITE® to cracking soybeans or after the soybeans crop has emerged because severe injury or death of crop will occur.

Degree of control and duration of effect depend on: rate used, weed spectrum, growing conditions at and following time of

treatment, soil pH, texture, organic matter, moisture and precipitation.

Best residual control is obtained if ENLITE® is applied to moist soil and followed by rainfall or irrigation (~1") before weeds germinate. Several small rainfalls of less than 1/4" each are not as beneficial as one large rainfall of 1/2- 1".

On dry soil, more moisture is required for activation (1- 2") before weed emergence. If moisture is insufficient to activate the herbicide, a rotary hoeing or shallow cultivation should be made after emergence of the crop while weeds are small enough to be controlled by mechanical means. Deep cultivation reduces the effectiveness of ENLITE® and should be avoided.

A maximum of 2.8 oz/acre ENLITE® can applied during the use season (any soil pH).

ROTATIONAL INTERVALS

Follow Recrop Interval 1 if the field is in the Nebraska (Northern Region) and:

- A maximum of 2.8 oz/acre ENLITE™ was applied during the use season (any soil pH).

ACCEPTED
~~XXXXXXXXXX~~
in EPA Letter Dated FEB 25 2010

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No.

352-757

3 of 3

Rotational Cropping Intervals		
For all recommended Fall and Spring DuPont™ ENLITE® uses, including sequentials with DuPont™ CLASSIC® or DuPont™ SYNCHRONY® XP ENLITE® Crop Rotational Interval in Months		
Crop	Interval 1	Interval 2
Soybeans	0	0
Barley, Ryegrass, Wheat, Winter Rye	4	4
Dry Beans, Kidney Beans Peas, Snap Beans	9	9
Field Corn*(States in Northern and Central Regions)	9	9
Popcorn	9	9
Sorghum	9	9
Tobacco (transplant)	9	9
Tomato (transplant)	9	9
Peanuts	15	6
Rice	15	9
Cotton	9	9
Alfalfa	12	12
Clover	12	12
Oats	12	12
Cabbage	18	18
Canola (Rapeseed)	18	18
Cucumber	18	9
Flax	18	18
Lentils	18	18
Mustard	18	18
Pumpkins	18	9
Sunflower	18	9
Sweet Corn	18	9†
Watermelon	18	9
Carrots	30	30
Onions	30	30
Potatoes	30	30
Sugar Beets	30	30
Any crop not listed	30	30

*The term "Field Corn" is defined to include only that corn grown for grain or silage or for seed corn relative to the Rotational Crop Guidelines section of this label.
 † Rotational crop intervals are for processing Sweet Corn varieties only.
 The rotational crop interval for other Sweet Corn varieties is 18 months

IMPORTANT
BEFORE USING ENLITE®, READ AND FOLLOW ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA-REGISTERED LABEL.

This bulletin contains new or supplemental instructions for use of this product which do not appear on the EPA-registered package label. Follow the instructions carefully. This labeling must be in the possession of the user at the time of pesticide application.

Expiration Date: October 8, 2011
 DR-1063 020310-3