

352-768

2/2/2012

1 of 3



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY  
Washington, D.C. 20460

OFFICE OF  
CHEMICAL SAFETY AND  
POLLUTION PREVENTION

FEB 2 2012

Tim McPherson  
E.I. du Pont de Nemours and Company  
1007 Market Street  
Wilmington, DE 19898

Subject: Supplemental Label (new uses on Caneberry Subgroup 13-07A and  
Bushberry Subgroup 13-07B)  
DuPont Matrix SG Herbicide  
EPA Reg. No. 352-768  
Application Dated September 28, 2010

Dear Mr. McPherson:

The supplemental labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act, as amended, is acceptable.

The supplemental labeling must be incorporated into the main product labeling within eighteen (18) months of the date of this letter.

A stamped copy of your label is enclosed for your records. You must submit one (1) copy of the final printed label before you release the product for shipment. Products shipped after eighteen (18) months from the date of this letter must bear the new revised label. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA §6(e). Your release for shipment of the product constitutes acceptance of these conditions.

If you have any questions, please contact Mindy Ondish at (703)605-0723 or at [ondish.mindy@epa.gov](mailto:ondish.mindy@epa.gov).

Sincerely,

Kable Bo Davis  
Product Manager 25  
Herbicide Branch  
Registration Division (7505P)



DuPont Crop Protection

ACCEPTED  
FEB 2 2012  
Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 352-768

SUPPLEMENTAL LABELING

DUPONT™ MATRIX® SG  
HERBICIDE  
FOR USE IN BUSHBERRY  
SUBGROUP 13-07B AND  
CANE BERRY SUBGROUP 13-07A

DUPONT™ MATRIX® SG Herbicide

EPA Reg. No. 352-768

FOR USE IN BUSHBERRY SUBGROUP 13-07B  
AND CANEBERRY SUBGROUP 13-07A

This supplemental labeling expires on February 1, 2015 and must not used or distributed after this date.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling. DuPont™ MATRIX® SG must be used only in accordance with instructions on this label or in separate published DuPont instructions.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

APPLICATION INFORMATION

BLUEBERRY (High Bush)

For broadcast applications, make a single application of MATRIX® SG preemergence or early postemergence to actively growing weeds at 4 ounces per acre per year. Use a directed spray application adjusted to provide complete coverage of the weeds while minimizing the amount of spray coming into contact with the blueberry plants. When applied as a banded treatment (50% band or less), MATRIX® SG may be applied twice per year.

Allow a minimum of 30 days between applications.

Application made after bud break may cause temporary chlorosis and/or stunting of leaves contacted by the spray.

Use MATRIX® SG on high bush blueberries that have gone through at least one growing season and are in good health and vigor. MATRIX® SG may be applied in tank mixture with other

herbicides registered for use in high bush blueberries.

Do not apply by air.  
Do not use on soils classified as Sand.  
Do not apply within 21 days of first harvest.  
Do not apply more than 4 ounces per acre on a broadcast application basis per year.

BLUEBERRY (Low Bush)

All applications of MATRIX® SG are to be applied in the Vegetative Year growth stage of low bush blueberries. Make a single broadcast application of MATRIX® SG preemergence or early postemergence to actively growing weeds at 4 ounces per acre per year. When applied as a banded treatment (50% treated band or less) MATRIX® SG may be applied twice per year.

Allow a minimum of 30 days between applications.

For broadcast treatments, make the application prior to bud break of the blueberries. After bud break, use a directed spray application adjusted to provide complete coverage of the weeds while minimizing spray contact with the blueberry plants.

Application made after bud break may cause temporary chlorosis and/or stunting of foliage contacted by the spray.

Use MATRIX® SG on low bush blueberries that have gone through at least one growing season and are in good health and vigor.

MATRIX® SG may be applied in tank mixture with other herbicides registered for use in low bush blueberries.

Do not apply by air.  
Do not use on soils classified as Sand.  
Do not apply within 21 days of first harvest.

Do not apply more than 4 ounces per acre on a broadcast application basis per year.

**CANEBERRY (Raspberry, Blackberry)**

For broadcast applications, make a single application of MATRIX® SG preemergence or early postemergence to actively growing weeds at 4 ounces per acre per year. Use a directed spray application adjusted to provide complete coverage of the weeds while minimizing the amount of spray coming into contact with the caneberry plants. When applied as a banded treatment (50% band or less), MATRIX® SG may be applied twice per year.

Allow a minimum of 30 days between applications.

If primocanes are up at time of treatment, temporary chlorosis of foliage and/or stunting of primocane growth may occur. These symptoms are temporary and do not affect the overall health and vigor of primocane development.

Use MATRIX® SG on caneberries that have gone through at least one growing season and are in good health and vigor.

MATRIX® SG may be applied in tank mixture with other herbicides registered for use in caneberries.

Do not apply by air.

Do not use on soils classified as Sand.

Do not apply within 21 days of first harvest.

Do not apply more than 4 ounces per acre on a broadcast application basis per year.

**IMPORTANT  
BEFORE USING DUPONT™ MATRIX® SG  
HERBICIDE, READ AND FOLLOW ALL  
APPLICABLE DIRECTIONS, RESTRICTIONS  
AND PRECAUTIONS ON THE EPA  
REGISTERED LABEL.**

This bulletin contains new or supplemental instructions for use of the product which may not appear on the package label, and may contain recommendations which are not appropriate for all geographies and weed control situations. Follow the instructions carefully.

This labeling must be in the possession of the user at the time of pesticide application.