



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, DC 20460

**OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION**

April 18, 2023

Wes Marchione
Regulatory Leader – Regulatory Affairs
Corteva Agriscience
9330 Zionsville Road
Indianapolis, IN 46268

Subject: Notification per PRN 98-10 – Updated primary and alternate brand names,
company details, and warranty statement
Product Name: Accent XP
EPA Registration Number: 352-817
Application Date: 06/23/2022
Decision Number: 585689

Dear Wes Marchione:

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10 for the above referenced product. The Registration Division (RD) has conducted a review of this request for its applicability under PRN 98-10 and finds that the action requested falls within the scope of PRN 98-10.

The label submitted with the application has been stamped “NOTIFICATION” and placed in our records.

The alternate brand name, “*DuPont Accent XP*” has been added to the product record.

Should you wish to add/retain a reference to the company’s website on your label, then please be aware that the website becomes labeling under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA) and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) lists examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product’s label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA’s Office of Enforcement and Compliance.

Page 2 of 2
EPA Reg. No. 352-817
Decision No. 585689

If you have any questions, please contact Jennifer Drobish at 202-566-2642 or at Drobish.jennifer@epa.gov.

Sincerely,

A handwritten signature in cursive script that reads "Aswathy Balan".

Aswathy Balan
Senior Regulatory Specialist
Fungicide and Herbicide Branch
Registration Division
Office of Pesticide Programs

(Base Label):

NICOSULFURON | GROUP 2 | HERBICIDE

DuPont™ Accent® XP**[Alternate Brand Name: DuPont Accent® XP]
HERBICIDE****For use on Field Corn grown for grain, silage or seed, Popcorn, and Sweet Corn.****Water-Dispersible Granule**

Active Ingredients	By Weight
Nicosulfuron 2-[[[(4,6-dimethoxypyrimidin-2-yl)aminocarbonyl]aminosulfonyl]-N,N-dimethyl-3-pyridinecarboxamide	75%
Other Ingredients	25%
TOTAL	100%

Keep Out of Reach of Children**CAUTION**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you **DO NOT** understand this label, find someone to explain it to you in detail.)

Precautionary Statements**Hazards to Humans and Domestic Animals****CAUTION**

Causes moderate eye irritation. Harmful if absorbed through skin. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

PERSONAL PROTECTIVE EQUIPMENT (PPE)**Applicators and other handlers must wear:**

- Long-sleeved shirt and long pants
- Chemical resistant gloves made of any water proof material including butyl rubber, natural rubber, neoprene rubber, or nitrile rubber
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENGINEERING CONTROL STATEMENTS

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR part 170.240 (d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

Important: When reduced PPE is worn because a closed system is being used, handlers must be provided all PPE specified above for "Applicators and Other Handlers" and have such PPE immediately available for use in an emergency, including a spill or equipment breakdown.

NOTIFICATION

352-817

The applicant has certified that no changes, other than those reported to the Agency have been made to the labeling. The Agency acknowledges this notification by letter dated:

04/18/2023

USER SAFETY RECOMMENDATIONS

USERS SHOULD:-

- Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of the gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

STORAGE AND DISPOSAL

DO NOT contaminate water, food or feed by storage and disposal.

Pesticide Storage: Store product in original container only. Store in a cool, dry place.

Pesticide Disposal: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

Container Handling: Refer to the Net Contents section of this product's labeling for the applicable "Nonrefillable Container" or "Refillable Container" designation.

Nonrefillable Plastic and Metal Containers (Capacity Equal to or Less Than 50 Pounds):

Nonrefillable container. **DO NOT** reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then, for Plastic Containers, offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration. **DO NOT** burn, unless allowed by state and local ordinances. For Metal Containers, offer for recycling if available or reconditioning if appropriate, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

Nonrefillable Plastic and Metal Containers (Capacity Greater Than 50 Pounds): Nonrefillable container. **DO NOT** reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then, for Plastic Containers, offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration. **DO NOT** burn, unless allowed by state and local ordinances. For Metal Containers, offer for recycling if available or reconditioning if appropriate, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

Nonrefillable Plastic and Metal Containers, e.g., Intermediate Bulk Containers [IBC] (Size or Shape Too Large to be Tipped, Rolled or Turned Upside Down): Nonrefillable container. **DO NOT** reuse or refill this container. Clean container promptly after emptying the contents from this container into application equipment or mix tank and before final disposal using the following pressure rinsing procedure. Insert a lance fitted with a suitable tank cleaning nozzle into the container and ensure that the water spray thoroughly covers the top, bottom and all sides inside the container. The nozzle manufacturer generally provides instructions for the appropriate spray pressure, spray duration and/or

spray volume. If the manufacturer's instructions are not available, pressure rinse the container for at least 60 seconds using a minimum pressure of 30 PSI with a minimum rinse volume of 10% of the container volume. Drain, pour or pump rinsate into application equipment or rinsate collection system. Repeat this pressure rinsing procedure two more times. Then, for Plastic Containers, offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration. For Metal Containers, offer for recycling if available or reconditioning if appropriate, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

Nonrefillable Paper or Plastic Bags, Fiber Sacks including Flexible Intermediate Bulk

Containers (FIBC) or Fiber Drums With Liners: Nonrefillable container. **DO NOT** reuse or refill this container. Completely empty paper or plastic bag, fiber sack or drum liner by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application or manufacturing equipment. Then offer for recycling if available or dispose of empty paper or plastic bag, fiber sack or fiber drum and liner in a sanitary landfill, or by incineration. **DO NOT** burn, unless allowed by state and local ordinances.

Refillable Fiber Drums With Liners: Refillable container (fiber drum only). Refilling Fiber Drum: Refill this fiber drum with **ACCENTAccent®** XP containing nicosulfuron only. **DO NOT** reuse this fiber drum for any other purpose. Cleaning before refilling is the responsibility of the refiller. Completely empty liner by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application or manufacturing equipment. Disposing of Fiber Drum and/or Liner: **DO NOT** reuse this fiber drum for any other purpose other than refilling (see preceding). Cleaning the container (liner and/or fiber drum) before final disposal is the responsibility of the person disposing of the container. offer the liner for recycling if available or dispose of liner in a sanitary landfill, or by incineration. **DO NOT** burn, unless allowed by state and local ordinances. If drum is contaminated and cannot be reused, dispose of it in the manner required for its liner. To clean the fiber drum before final disposal, completely empty the fiber drum by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application or manufacturing equipment. Then offer the fiber drum for recycling if available or dispose of in a sanitary landfill, or by incineration. **DO NOT** burn, unless allowed by state and local ordinances.

All Other Refillable Containers: Refillable container. Refilling Container: Refill this container with **ACCENTAccent®** XP containing nicosulfuron only. **DO NOT** reuse this container for any other purpose. Cleaning before refilling is the responsibility of the refiller. Prior to refilling, inspect carefully for damage including cracks, punctures, abrasions, worn out threads and closure devices. If damage is found, **DO NOT** use container, contact **DuPont-Corteva Agriscience** at the number below for instructions. Check for leaks after refilling and before transporting. If leaks are found, **DO NOT** reuse or transport container, contact **DuPont-Corteva Agriscience** at the number below for instructions. Disposing of Container: **DO NOT** reuse this container for any other purpose other than refilling (see preceding). Cleaning the container before final disposal is the responsibility of the person disposing of the container. To clean the container before final disposal, use the following pressure rinsing procedure. Insert a lance fitted with a suitable tank cleaning nozzle into the container and ensure that the water spray thoroughly covers the top, bottom and all sides inside the container. The nozzle manufacturer generally provides instructions for the appropriate spray pressure, spray duration and/or spray volume. If the manufacturer's instructions are not available, pressure rinse the container for at least 60 seconds using a minimum pressure of 30 PSI with a minimum rinse volume of 10% of the container volume. Drain, pour or pump rinsate into application equipment or rinsate collection system. Repeat this pressure rinsing procedure two more times. Then, for Plastic Containers, offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration. **DO NOT** burn, unless allowed by state and local ordinances. For Metal Containers, offer for recycling if available or reconditioning if appropriate, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

DO NOT transport if container is damaged or leaking. If the container is damaged, leaking or obsolete, or in the event of a major spill, fire or other emergency, call 1-800-441-3637/992-5994, day or night.

Refer to inside of label booklet for additional precautionary information and Directions for Use.

Notice: Read the entire label. Use only according to label directions. Before using this product, read Warranty Disclaimer, Inherent Risks of Use and Limitation of Remedies at end of label booklet. If terms are unacceptable, return at once unopened.

~~NOTICE TO BUYER: Purchase of this material does not confer any rights under patents of countries outside of the United States.~~

In case of emergency endangering health or the environment involving this product, call: 1-800-~~441-3637~~992-5994

Agricultural Chemical: **DO NOT** ship or store with food, feeds, drugs, or clothing.

EPA Reg. No. 352-817

EPA Est. No. _____

~~™ Trademarks of Corteva Agriscience and its affiliated companies™ Trademarks of Dow AgroSciences, DuPont or Pioneer and their affiliated companies or respective owners~~

For product information call: 1-800-258-3033

Produced for

Corteva Agriscience LLC

9330 Zionsville Road

Indianapolis, IN 46268

~~E. I. du Pont de Nemours and Company, Chestnut Run Plaza, 974 Centre~~

~~Road, Wilmington, DE 19805 U.S.A.~~

NET CONTENTS _____

(Cover/Shipping Label):

NICOSULFURON	GROUP	2	HERBICIDE
--------------	-------	---	-----------

DuPont™ Accent® XP[Alternate Brand Name: [DuPont Accent® XP](#)]**HERBICIDE****For use on Field Corn grown for grain, silage or seed, Popcorn, and Sweet Corn.****Water-Dispersible Granule**

Active Ingredients	By Weight
Nicosulfuron 2-[[[4,6-dimethoxypyrimidin-2-yl)aminocarbonyl]aminosulfonyl]-N,N-dimethyl-3-pyridinecarboxamide	75%
Other Ingredients	25%
TOTAL	100%

Keep Out of Reach of Children**CAUTION**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you **DO NOT** understand this label, find someone to explain it to you in detail.)

Refer to inside of label booklet for additional precautionary information and Directions for Use.

Notice: [Read the entire label. Use only according to label directions. Before using this product, read Warranty Disclaimer, Inherent Risks of Use and Limitation of Remedies at end of label booklet. If terms are unacceptable, return at once unopened.](#)

NOTICE TO BUYER: ~~Purchase of this material does not confer any rights under patents of countries outside of the United States.~~

In case of emergency endangering health or the environment involving this product, call: 1-800-441-3637 [992-5994](#)

Agricultural Chemical: **DO NOT** ship or store with food, feeds, drugs, or clothing.

EPA REG. NO. 352-854

EPA Est. _____

~~™ Trademarks of Corteva Agriscience and its affiliated companies.™ Trademarks of Dow AgroSciences, DuPont or Pioneer and their affiliated companies or respective owners~~

For product information call: 1-800-258-3033**Produced for****[Corteva Agriscience LLC](#)****[9330 Zionsville Road](#)****[Indianapolis, IN 46268](#) ~~E. I. du Pont de Nemours and Company, Chestnut Run Plaza, 974 Centre Road, Wilmington, DE 19805 U.S.A.~~**

NET CONTENTS_____

(Page 1 through end):

FIRST AID

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

IF SWALLOWED: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. **DO NOT** induce vomiting unless told to do so by the poison control center or doctor. **DO NOT** give anything by mouth to an unconscious person.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-800-~~441-3637~~[992-5994](tel:1-800-992-5994) for emergency medical treatment information.

Precautionary Statements

Hazards to Humans and Domestic Animals

CAUTION

Causes moderate eye irritation. Harmful if absorbed through skin. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical resistant gloves made of any water proof material including butyl rubber, natural rubber, neoprene rubber, or nitrile rubber
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENGINEERING CONTROL STATEMENTS

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR part 170.240 (d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

Important: When reduced PPE is worn because a closed system is being used, handlers must be provided all PPE specified above for "Applicators and Other Handlers" and have such PPE immediately available for use in an emergency, including a spill or equipment breakdown.

USER SAFETY RECOMMENDATIONS

USERS SHOULD:-

- Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet

- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of the gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

DO NOT apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. **DO NOT** contaminate water when disposing of equipment washwaters or rinsate. **DO NOT** apply where/when conditions favor runoff.

Groundwater Advisory

Nicosulfuron is known to leach through soil into groundwater under certain conditions as a result of label use. This chemical may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow.

Surface Water Advisory

This product may impact surface water quality due to runoff of rain water. This is especially true for poorly draining soils and soils with shallow ground water. This product is classified as having high potential for reaching surface water via runoff for several months or more after application. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features including ponds, streams, and springs will reduce the potential loading of this product from runoff water and sediment. Runoff of this product will be greatly reduced by avoiding applications when rainfall or irrigation is expected to occur within 48 hours.

Windblown Soil Particles Advisory

This product has the potential to move off-site due to wind erosion. Soils that are subject to wind erosion usually have a high silt and/or fine to very fine sand fractions and low organic matter content. Other factors which can affect the movement of windblown soil include the intensity and direction of prevailing winds, vegetative cover, site slope, rainfall, and drainage patterns. Avoid applying this product if prevailing local conditions may be expected to result in off-site movement.

Non-target Organism Advisory

This product is toxic to plants and may adversely impact the forage and habitat of non-target organisms, including pollinators, in areas adjacent to the treated area. Protect the forage and habitat of non-target organisms by minimizing spray drift. For further guidance and instructions on how to minimize spray drift, refer to the Mandatory Spray Drift Management section of this label.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

DO NOT apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective

equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

DO NOT enter or allow worker entry into treated areas during the restricted entry interval (REI) of 4 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, including plants, soil, or water, is:

Coveralls

Chemical resistant gloves made of any water proof material including butyl rubber, natural rubber, neoprene rubber, or nitrile rubber

Shoes plus socks

ACCENT@Accent XP herbicide must be used only in accordance with directions on this label, or as otherwise permitted by FIFRA. . To the extent consistent with applicable law, **DuPont-Corteva Agriscience** will not be responsible for losses or damage resulting from use of this product in any manner not specifically directed by **DuPont-Corteva Agriscience**. Always read the entire label, including the Limitation of Warranty and Liability.

STORAGE AND DISPOSAL

DO NOT contaminate water, food or feed by storage and disposal.

Pesticide Storage: Store product in original container only. Store in a cool, dry place.

Pesticide Disposal: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

Container Handling: Refer to the Net Contents section of this product's labeling for the applicable "Nonrefillable Container" or "Refillable Container" designation.

Nonrefillable Plastic and Metal Containers (Capacity Equal to or Less Than 50 Pounds):

Nonrefillable container. **DO NOT** reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then, for Plastic Containers, offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration. **DO NOT** burn, unless allowed by state and local ordinances. For Metal Containers, offer for recycling if available or reconditioning if appropriate, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

Nonrefillable Plastic and Metal Containers (Capacity Greater Than 50 Pounds): Nonrefillable container. **DO NOT** reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then, for Plastic Containers, offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration. **DO NOT** burn, unless allowed by state and local ordinances. For Metal Containers, offer for recycling if available or reconditioning if appropriate, or puncture and dispose of in a sanitary landfill, or by other procedures

approved by state and local authorities.

Nonrefillable Plastic and Metal Containers, e.g., Intermediate Bulk Containers [IBC] (Size or Shape Too Large to be Tipped, Rolled or Turned Upside Down): Nonrefillable container. **DO NOT** reuse or refill this container. Clean container promptly after emptying the contents from this container into application equipment or mix tank and before final disposal using the following pressure rinsing procedure. Insert a lance fitted with a suitable tank cleaning nozzle into the container and ensure that the water spray thoroughly covers the top, bottom and all sides inside the container. The nozzle manufacturer generally provides instructions for the appropriate spray pressure, spray duration and/or spray volume. If the manufacturer's instructions are not available, pressure rinse the container for at least 60 seconds using a minimum pressure of 30 PSI with a minimum rinse volume of 10% of the container volume. Drain, pour or pump rinsate into application equipment or rinsate collection system. Repeat this pressure rinsing procedure two more times. Then, for Plastic Containers, offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration. For Metal Containers, offer for recycling if available or reconditioning if appropriate, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

Nonrefillable Paper or Plastic Bags, Fiber Sacks including Flexible Intermediate Bulk Containers (FIBC) or Fiber Drums With Liners: Nonrefillable container. **DO NOT** reuse or refill this container. Completely empty paper or plastic bag, fiber sack or drum liner by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application or manufacturing equipment. Then offer for recycling if available or dispose of empty paper or plastic bag, fiber sack or fiber drum and liner in a sanitary landfill, or by incineration. **DO NOT** burn, unless allowed by state and local ordinances.

Refillable Fiber Drums With Liners: Refillable container (fiber drum only). Refilling Fiber Drum: Refill this fiber drum with [AccentACCENT®](#) XP containing nicosulfuron only. **DO NOT** reuse this fiber drum for any other purpose. Cleaning before refilling is the responsibility of the refiller. Completely empty liner by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application or manufacturing equipment. Disposing of Fiber Drum and/or Liner: **DO NOT** reuse this fiber drum for any other purpose other than refilling (see preceding). Cleaning the container (liner and/or fiber drum) before final disposal is the responsibility of the person disposing of the container. offer the liner for recycling if available or dispose of liner in a sanitary landfill, or by incineration. **DO NOT** burn, unless allowed by state and local ordinances. If drum is contaminated and cannot be reused, dispose of it in the manner required for its liner. To clean the fiber drum before final disposal, completely empty the fiber drum by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application or manufacturing equipment. Then offer the fiber drum for recycling if available or dispose of in a sanitary landfill, or by incineration. **DO NOT** burn, unless allowed by state and local ordinances.

All Other Refillable Containers: Refillable container. Refilling Container: Refill this container with [AccentACCENT®](#) XP containing nicosulfuron only. **DO NOT** reuse this container for any other purpose. Cleaning before refilling is the responsibility of the refiller. Prior to refilling, inspect carefully for damage including cracks, punctures, abrasions, worn out threads and closure devices. If damage is found, **DO NOT** use container, contact [DuPont-Corteva Agriscience](#) at the number below for instructions. Check for leaks after refilling and before transporting. If leaks are found, **DO NOT** reuse or transport container, contact [DuPont-Corteva Agriscience](#) at the number below for instructions. Disposing of Container: **DO NOT** reuse this container for any other purpose other than refilling (see preceding). Cleaning the container before final disposal is the responsibility of the person disposing of the container. To clean the container before final disposal, use the following pressure rinsing procedure. Insert a lance fitted with a suitable tank cleaning nozzle into the container and ensure that the water spray thoroughly covers the top, bottom and all sides inside the container. The nozzle manufacturer generally provides instructions for the appropriate spray pressure, spray duration and/or spray volume. If the manufacturer's instructions are not available, pressure rinse the container for at least 60 seconds using a minimum pressure of 30 PSI with a minimum rinse volume of 10% of the container volume. Drain,

pour or pump rinsate into application equipment or rinsate collection system. Repeat this pressure rinsing procedure two more times. Then, for Plastic Containers, offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration. **DO NOT** burn, unless allowed by state and local ordinances. For Metal Containers, offer for recycling if available or reconditioning if appropriate, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

DO NOT transport if container is damaged or leaking. If the container is damaged, leaking or obsolete, or in the event of a major spill, fire or other emergency, call 1-800-~~441-3637~~[992-5994](tel:992-5994), day or night.

PRODUCT INFORMATION

[AccentACCENT®](#) XP herbicide is a water-dispersible granule used at a rate of 0.33-1.33 ounces (0.015 - 0.062 lb ai) per acre for selective postemergence grass weed control in field corn grown for silage, seed or grain, popcorn and sweet corn,

RESTRICTIONS

CROPS	Maximum Oz of Product/Acre/ Single Application	Maximum Lb AI/Acre/Single Application	Maximum Number of Applications Per Year	Maximum Oz of Product /Acre/Year	Maximum Lb AI/A per Year	Retreat Interval (Days)	Last Treatment Preharvest Interval
Field Corn – Grown for Grain, Silage; Seed, Popcorn, or Sweet Corn	1.33 oz	0.062 lb ai	2*	1.33 oz	0.062 lb ai	7	DO NOT harvest corn grain within 70 days or harvest corn forage or stover within 45 days of an AccentACCENT® XP application

* Make only 1 application of [AccentACCENT®](#) XP to sweet corn. See Popcorn, Field Corn Grown for Seed and Sweet Corn use directions.

The combined dosage of sequential applications cannot exceed 1.33 ounces (0.062 lb ai) per acre of [AccentACCENT®](#) XP in corn.

When tank mixing with other products that contain nicosulfuron, **DO NOT** apply more than 1.0 ounce (0.062 lb) of nicosulfuron in corn per year.

The combined dosage of sequential applications cannot exceed 1.33 ounces (0.062 lb ai) per acre of [AccentACCENT®](#) in corn.

DO NOT tank mix [AccentACCENT®](#) XP with products containing bentazon or severe corn crop injury may occur.

DO NOT tank mix [AccentACCENT®](#) XP with foliar-applied organophosphate insecticides including chlorpyrifos and malathion, etc, as severe crop injury may occur. To avoid crop injury or antagonism, apply these products at least seven days before or 3 days after the application of [AccentACCENT®](#) XP.

DO NOT apply [AccentACCENT®](#) XP within 45 days of crop emergence where the organophosphate insecticide, terbufos was applied since corn crop injury may occur. Applications made to corn previously treated with chlorpyrifos or other similar organophosphate insecticides may result in unacceptable crop injury. Any crop injury or yield loss resulting from these applications are the responsibility of the grower.

DO NOT make a late application of [AccentACCENT®](#) XP to field corn grown for grain or silage that is taller than 36 inches or that exhibits 10 or more collars (V10), whichever is more restrictive.

DO NOT apply [AccentACCENT®](#) XP to any white popcorn inbred, or white popcorn hybrid.

DO NOT apply [AccentACCENT®](#) XP to sweet corn taller than 18 inches or which exhibits 6 or more leaf collars (V6).

DO NOT graze, feed forage, grain or fodder (stover) from corn treated areas to livestock within 30 days of an [AccentACCENT®](#) XP application

DO NOT apply [AccentACCENT®](#) XP to corn that exhibits herbicide injury from previous applications made to the current or preceding crop.

DO NOT use liquid nitrogen fertilizer as the total carrier solution for postemergence applications.

Injury or loss of desirable trees or vegetation may result from failure to observe the following:

DO NOT apply [AccentACCENT®](#) XP or drain or flush application equipment on or near desirable trees or other plants, or on areas where their roots may extend, or in locations where the chemical may be washed or moved into contact with their roots.

DO NOT use on lawns, walks, driveways, tennis courts or other residential areas.

DO NOT contaminate any body of water

DO NOT apply this product through any type of irrigation system.

PRECAUTIONS

[AccentACCENT®](#) XP may interact with certain insecticides previously applied to the crop. Crop response varies with field corn type, insecticide used, insecticide application methods, and soil type.

[AccentACCENT®](#) XP may be applied to corn previously treated with chlorethoxyfos, bifenthrin + chlorethoxyfos, cyfluthrin + tebupirimphos, tefluthrin, or other non-organophosphate soil insecticides regardless of soil type.

Prevent drift or spray onto desirable plants.

Thoroughly clean application equipment immediately after use (See Sprayer Cleanup section of this label).

For all application systems, use 50-mesh or larger strainer screens.

WEED RESISTANCE MANAGEMENT

[AccentACCENT®](#) XP, which contains the active ingredient nicosulfuron, is a group 2 herbicide based on the mode of action classification system of the Weed Science Society of America.

Proactively implementing diversified weed control strategies to minimize selection for weed populations resistant to one or more herbicides is a best practice. A diversified weed management program may include the use of multiple herbicides with different sites of action and overlapping weed spectrum with or without tillage operations and/or other cultural practices.

Research has demonstrated that using the labeled rate and directions for use is important to delay the selection for resistance.

The continued effectiveness of this product depends on the successful implementation of a weed resistance management program.

To aid in the prevention of developing weeds resistant to this product, users should:

- Scout fields before application to ensure herbicides and rates will be appropriate for the weed species and weed sizes present.
- Start with a clean field, using either a burndown herbicide application or tillage.
- Control weeds early when they are relatively small (less than 4 inches).
- Apply full rates of this product for the most difficult to control weed in the field at the specified time (correct weed size) to minimize weed escapes.
- Scout fields after application to detect weed escapes or shifts in control of weed species.
- Control weed escapes before they reproduce by seed or proliferate vegetatively.
- Report any incidence of non-performance of this product against a particular weed to your [DuPont company](#) representative, local retailer, or county extension agent.
- Contact your sales representative, crop advisor, or extension agent to find out if suspected resistant weeds to this MOA have been found in your region. If resistant biotypes of target weeds have been reported, use the application rates of this product specified for your local conditions. Tank mix products so that there are multiple effective sites of actions for each target weed.
- If resistance is suspected, treat weed escapes with an herbicide having a site of action other than group 2 and/or use nonchemical methods to remove escapes, as practical, with the goal of preventing further seed production.
- Suspected herbicide-resistant weeds may be identified by these indicators:
 - Failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds;
 - A spreading patch of non-controlled plants of a particular weed species; and
 - Surviving plants mixed with controlled individuals of the same species.

Additionally, users should follow as many of the following herbicide resistance management practices as practical:

- Use a broad-spectrum soil-applied herbicide with other sites of action as a foundation in a weed control program.
- Utilize sequential applications of herbicides with alternative sites of action.
- Rotate the use of this product with non-group 2 herbicides.
- Avoid making more than two applications of 2 and any other group 2 herbicides within a single growing season unless mixed with an herbicide with a different site of action with an overlapping spectrum for the difficult-to-control weeds.
- Incorporate non-chemical weed control practices, including mechanical cultivation, crop rotation, cover crops and weed-free crop seeds, as part of an integrated weed control program.
- Use good agronomic principles that enhance crop development and crop competitiveness.
- Thoroughly clean plant residues from equipment before leaving fields suspected to contain resistant weeds.
- Manage weeds in and around fields, during and after harvest to reduce weed seed production.

INTEGRATED PEST MANAGEMENT

This product may be used as part of an Integrated Pest Management (IPM) program that can include biological, cultural, and genetic practices aimed at preventing economic pest damage. IPM principles and practices include field scouting or other detection methods, correct target pest identification, population monitoring, and treating when target pest populations reach locally determined action thresholds. Consult your state cooperative extension service, professional consultant or other qualified authorities to determine appropriate action treatment threshold levels for treating specific pest / crop systems in your

area.

MANDATORY SPRAY DRIFT MANAGEMENT

Aerial Applications:

- **DO NOT** release spray at a height greater than 10 feet above the vegetative canopy, unless a greater application height is necessary for pilot safety.
- For applications prior to the emergence of crops and target weeds, applicators are required to use a Coarse or coarser droplet size (ASABE S572.1).
- For all other applications, applicators are required to use a Medium or coarser droplet size (ASABE S572.1).
- The boom length must not exceed 65% of the wingspan for airplanes or 75% of the rotor blade diameter for helicopters.
- Applicators must use one-half swath displacement upwind at the downwind edge of the field.
- Nozzles must be oriented so the spray is directed toward the back of the aircraft.
- **DO NOT** apply when wind speeds exceed 10 miles per hour at the application site.
- **DO NOT** apply during temperature inversions.

Ground Boom Applications:

- Apply with the nozzle height recommended by the manufacturer, but no more than 3 feet above the ground or crop canopy
- For applications prior to the emergence of crops and target weeds, applicators are required to use a Coarse or coarser droplet size (ASABE S572.1).
- For all other applications, applicators are required to use a Medium or coarser droplet size (ASABE S572.1).
- **DO NOT** apply when wind speeds exceed 10 miles per hour at the application site.
- **DO NOT** apply during temperature inversions.

SPRAY DRIFT ADVISORIES

THE APPLICATOR IS RESPONSIBLE FOR AVOIDING OFF-SITE SPRAY DRIFT.

BE AWARE OF NEARBY NON-TARGET SITES AND ENVIRONMENTAL CONDITIONS.

IMPORTANCE OF DROPLET SIZE

An effective way to reduce spray drift is to apply large droplets. Use the largest droplets that provide target pest control. While applying larger droplets will reduce spray drift, the potential for drift will be greater if applications are made improperly or under unfavorable environmental conditions.

Controlling Droplet Size – Ground Boom

- **Volume** - Increasing the spray volume so that larger droplets are produced will reduce spray drift. Use the highest practical spray volume for the application. If a greater spray volume is needed, consider using a nozzle with a higher flow rate.
- **Pressure** - Use the lowest spray pressure recommended for the nozzle to produce the target spray volume and droplet size.
- **Spray Nozzle** - Use a spray nozzle that is designed for the intended application. Consider using nozzles designed to reduce drift.

Controlling Droplet Size – Aircraft

- **Adjust Nozzles** - Follow nozzle manufacturers recommendations for setting up nozzles. generally, to reduce fine droplets, nozzles should be oriented parallel with the airflow in flight.

BOOM HEIGHT – Ground Boom

Use the lowest boom height that is compatible with the spray nozzles that will provide uniform coverage. For ground equipment, the boom should remain level with the crop and have minimal bounce.

RELEASE HEIGHT - Aircraft

Higher release heights increase the potential for spray drift. When applying aerially to crops, **DO NOT** release spray at a height greater than 10 feet above the crop canopy, unless a greater application height is necessary for pilot safety.

SHIELDED SPRAYERS

Shielding the boom or individual nozzles can reduce spray drift. Consider using shielded sprayers. Verify that the shields are not interfering with the uniform deposition of the spray on the target area.

TEMPERATURE AND HUMIDITY

When making applications in hot and dry conditions, use larger droplets to reduce effects of evaporation.

TEMPERATURE INVERSIONS

Drift potential is high during a temperature inversion. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. The presence of an inversion can be indicated by ground fog or by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing. Avoid applications during temperature inversions.

WIND

Drift potential generally increases with wind speed. **AVOID APPLICATIONS DURING GUSTY WIND CONDITIONS.**

- Applicators need to be familiar with local wind patterns and terrain that could affect spray drift.

DRIFT CONTROL ADDITIVES

Using product compatible drift control additives can reduce drift potential. When a drift control additive is used, read and carefully observe cautionary statements and all other information on the additive's label. If using an additive that increases viscosity, ensure that the nozzles and other application equipment will function properly with a viscous spray solution.

Preferred drift control additives have been certified by the Chemical Producers and Distributors Association (CPDA).

FIELD CORN GROWN FOR GRAIN OR SILAGE

Timing to Crop

[AccentACCENT®](#) XP is registered for use on field corn, high lysine, waxy, white or other food grade corn hybrids.

[AccentACCENT®](#) XP may be broadcast to corn up to 20" tall (free standing) or that is exhibiting up to and including 6 leaf collars (V6), whichever is more restrictive.

While [AccentACCENT®](#) XP has a wide application window, research has shown best results are obtained when applications are made early postemergence when corn and weeds are small. Target applications to corn that is less than 12" tall for best overall performance.

Timing to Weeds

Apply [AccentACCENT®](#) XP when grasses are young and actively growing, but before they exceed the sizes indicated in the “Weeds Controlled” table with 0.67 ounces (0.031 lb ai) [AccentACCENT®](#) XP. Treat heavy infestations of weeds before they become too competitive with the crop, especially where soil moisture and/or fertility are limited. [AccentACCENT®](#) XP provides weed control via foliar absorption. [AccentACCENT®](#) XP only controls those weeds that have emerged. For later-emerging weeds, a second application or a timely cultivation is required. Applications made to weeds larger than the size indicated on this label or to weeds under stress may result in unsatisfactory control. Refer to LATE APPLICATIONS.

On glyphosate-resistant corn, glyphosate products including “ABUNDIT™ Edge” and Durango® DMA®, may be applied with [AccentACCENT®](#) XP after weeds emerge but before they reach the maximum size listed on the glyphosate herbicide label.

On glufosinate-resistant corn, glufosinate containing products may be applied with [AccentACCENT®](#) XP after weeds emerge but before they reach the maximum size listed on the glufosinate herbicide label.

LATE APPLICATIONS IN FIELD CORN GROWN FOR GRAIN OR SILAGE

[AccentACCENT®](#) XP may be applied to field corn as a directed postemergence application on corn that is taller than 20” or which has more than 6 collars (whichever occurs first).

- For corn 20” to 36” tall, apply [AccentACCENT®](#) XP with drop nozzles only and avoid spraying into the whorl of cornstalks.
- **DO NOT** apply to corn that is taller than 36” or that exhibits 10 or more collars (V10), whichever is most restrictive.

Applications made to weeds larger than those listed on this label may vary from complete control to suppression. Level of control will depend on the weed species, stage of growth, and environmental conditions.

Choices must be made between the risks that arise from applications made beyond the proper time for [AccentACCENT®](#) XP use, and the effects of season long grass competition and/or harvest complications. These choices must balance risks from improperly timed [AccentACCENT®](#) XP use that include, but are not limited to:

- Yield loss due to competition: Research indicates competition from foxtail exceeding 4 inches in height may reduce corn yields. Applications to foxtail and other annual grasses that exceed the sizes stated on the label increases the risk of yield losses due to prolonged competition with the crop even though control may be acceptable.
- Incomplete control of grasses beyond labeled size: Applications to grasses that exceed the labeled sizes can result in reduced control. This incomplete control may reduce corn yield.
- Incomplete grass control due to herbicide stress: Grasses under stress from previous herbicide applications may not be actively growing and susceptible to [AccentACCENT®](#) XP.
- Ear malformation: Applications of [AccentACCENT®](#) XP on corn that has 7 to 10 collars (V7 to V10) increases the potential for ear malformation (pinching). This risk may be greatly reduced, but not eliminated, by using drop nozzles properly adjusted so as to not apply [AccentACCENT®](#) XP into the

corn whorl.

APPLICATION RATE

Optimum control of the weeds listed can be achieved with 0.67 ounces (0.031 lb ai) of [AccentACCENT® XP](#). Weeds that exceed the listed weed sizes by up to 50% may be partially controlled with rates between 0.67 and 1.33 ounces (0.031 and 0.062 lb ai) of [AccentACCENT® XP](#) per acre.

[AccentACCENT® XP](#) may be applied at 0.33 - 0.67 ounces (0.015 – 0.031 lb ai) for limited control of certain small grass weeds. See the table for reduced rates under ADDITIONAL DIRECTIONS for details.

As weeds mature, their sensitivity to [AccentACCENT® XP](#) decreases. As grassy weeds become mature (more than 3 tillers), they may not reach the size listed below, due to drought or other environmental factors. Treat grassy weeds that are maturing rapidly before they reach the stages listed below.

When applied as directed, [AccentACCENT® XP](#) will control the following weeds:

Weeds controlled with 0.67 ounces (0.031 lb ai) [AccentACCENT® XP](#).

Grasses	Maximum Height or Diameter
Barnyardgrass [†]	4"
Broadleaf signalgrass	2"
Foxtails (bristly, giant [†] green [†] , yellow [†])	4"
Itchgrass	6"
Johnsongrass [†] seedling	12"
rhizome	18"
Panicum (Texas, browntop)	3"
Fall	4"
Quackgrass*	10"
Ryegrass (Italian, perennial) [†]	6"
Sandbur (field, longspine)*	3"
Timothy	6"
Volunteer cereals (barley, oats, rye, triticale, wheat)	6"***
Wild oats [†]	4"
Wild proso millet	4"
Wirestem muhly*	8"
Witchgrass	6"
Woolly Cupgrass* [†]	4"

* Requires the use of COC plus ammonium nitrogen fertilizer. Cultivation or re-treatment may be required. See "FOR ADDITIONAL CONTROL OF LATER EMERGING GRASSES"

**10 inches in the states of WA, OR, ID, and MT, where the use of MSO adjuvants are preferred. See SPRAY ADJUVANTS.

† Naturally occurring resistant biotypes are known to occur.

Broadleaves	Maximum Height or Diameter
Burcucumber	3"
Dandelion	6"
Hemp dogbane*	4"
Jimsonweed	3"
Morningglory (ivy leaf, pitted)	3"
Tall	2"
Pigweed (redroot, smooth)	4"
Pokeweed*	4"
Smartweeds (ladysthumb, PA)	4"

Thistle, Canada*	4"
------------------	----

*Suppression

POPCORN, FIELD CORN GROWN FOR SEED AND SWEET CORN

[AccentACCENT®](#) XP may be broadcast or applied with drop nozzles to popcorn or field corn grown for seed that is less than 20" tall (free-standing) or that exhibits up to and including 5 leaf-collars (V5), whichever is most restrictive. **DO NOT** apply to corn that is taller than 20" or that exhibits more than 5 leaf-collars (V5), whichever is more restrictive.

Many seed companies have tested seed corn inbreds or yellow popcorn hybrids for sensitivity to [AccentACCENT®](#) XP and have reported excellent safety. **DO NOT** apply [AccentACCENT®](#) XP to any white popcorn inbred, or white popcorn hybrid unless specifically approved by the seed company. This includes "White Dynamite" popcorn.

[AccentACCENT®](#) XP may be applied to certain sweet corn hybrids grown for fresh markets or under contract for processing. Applications of [AccentACCENT®](#) XP may be applied broadcast or with drop nozzles (post-directed) on sweet corn up to 12 inches tall or up to and including 5 leaf-collars (V5). For sweet corn 12 - 18 inches tall, apply only with drop nozzles. **DO NOT** apply to sweet corn taller than 18 inches or those which exhibit 6 or more leaf-collars (V6). Make only one application of [AccentACCENT®](#) XP to sweet corn per year.

Sweet corn hybrid sensitivity to [AccentACCENT®](#) XP is highly variable, and not all hybrids have been tested for crop sensitivity. Contact your sales representative for information on local sweet corn hybrids that have been evaluated with [AccentACCENT®](#) XP.

Not all seed corn inbreds, popcorn or sweet corn hybrids have been tested, nor does [DuPont-Corteva Agriscience](#) have access to all seed company data. Consequently, to the extent consistent with the law, [DuPont-Corteva Agriscience](#) is not responsible for any crop injury arising from the use of [AccentACCENT®](#) XP on field corn grown for seed, popcorn or sweet corn. When tank mixing, check the tank mix partner label for sensitivities and instructions for use. It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements for each product in the tank mix.

SEQUENTIAL APPLICATIONS FOLLOWING PREEMERGENCE HERBICIDES

[AccentACCENT®](#) XP is best used in a planned postemergence weed control program as part of a sequential application herbicide program, following a before-planting application of: BASIS® Blend, CINCH®_®, CINCH ATZ, CINCH ATZ Lite, FulTime®_® NXT, INSTIGATE®_®, Keystone®_® NXT, Keystone®_® LA NXT, LEADOFF®_®, PREQUEL®_®, Resicore®_®, RESOLVE®_® Q, SureStart®_® II, Surpass®_® NXT herbicides and/or other pre-applied corn herbicides. Refer to the label of the respective corn herbicide partner for specific use directions.

DO NOT apply [AccentACCENT®](#) XP to corn that exhibits herbicide injury from previous applications made to the current or preceding crop.

TANK MIX APPLICATIONS

For Additional Control of Broadleaf Weeds

[AccentACCENT®](#) XP may be tank mixed with herbicides registered for postemergence application in corn for additional control of broadleaf weeds. It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive

directions for use and precautionary statements of each product in the tank mixture.

See SPRAY ADJUVANTS for adjuvant rate specification.

In addition to the tank mixtures noted above, [AccentACCENT®](#) XP may be tank mixed with the labeled rates of atrazine*, dicamba, dicamba + diflufenpyr †, dicamba + primisulfuron-methyl† and mesotrione for improved control of many broadleaf weeds, including cocklebur, dandelion, Eastern black nightshade, lambsquarters, pigweeds, ragweeds, PA smartweed and velvetleaf.

* Make applications to emerged corn before the corn reaches 12" tall.

† **DO NOT** apply to sweet corn, seed corn, or popcorn.

†† **DO NOT** apply to sweet corn.

If other brands or formulations are used, adjust rates of active ingredients to correspond to the products indicated. Formulations of products other than those listed may not have been tested with [AccentACCENT®](#) XP. Check with the manufacturer for information on tank mix compatibility prior to using (see TANK MIX COMPATIBILITY TESTING).

Crop oil concentrate plus ammonium nitrogen fertilizer is the preferred adjuvant for tank mixtures when using products at the low end of the rate range indicated in the table. The use of nonionic surfactant is permitted in place of crop oil concentrate for tank mixtures containing dicamba, however, overall weed control may be reduced. See SPRAY ADJUVANTS for adjuvant rate specification.

DO NOT use MSO adjuvants when tank mixing [AccentACCENT®](#) XP with a mesotrione-containing product.

ADDITIONAL DIRECTIONS AND/OR DIRECTIONS FOR SPECIFIC WEED PROBLEMS

Reduced Rates of [AccentACCENT®](#) XP

[AccentACCENT®](#) XP may be applied at 0.33 - 0.67 ounces (0.015 – 0.031 lb ai) for control of the small grass weeds noted in the table below. Always use a crop oil concentrate plus ammonium nitrogen fertilizer when applying reduced rates of [AccentACCENT®](#) XP.

Weeds controlled with reduced rates of [AccentACCENT®](#) XP.

XP Rate Maximum Height or Diameter [AccentACCENT®](#)

Grasses	0.33 ounces (0.015 lb ai)	0.5 ounces (0.023 lb ai)	0.67 ounces (0.031 lb ai)
Barnyardgrass [†]	2"	3"	4"
Foxtails (bristly, giant [†] , green [†])	2"	3"	4"
yellow [†]	--	2"	4"
Itchgrass	2"	4"	6"
Johnsongrass [†] , seedling	--	8"	12"
rhizome	--	8"	18"
Panicum (Texas, browntop)	1"	2"	3"
fall	1"	2"	4"
Sandbur (field, longspine)	--	1"	3"
Timothy	2"	4"	6"
Volunteer cereals	--	2"	6"
Wild oats [†]	2"	3"	4"
Wild proso millet	--	2"	4"
Witchgrass	2"	4"	6"

Woolly cupgrass	--	--	4"
-----------------	----	----	----

† Naturally occurring resistant biotypes are known to occur.

Tank Mixtures with Atrazine

AccentACCENT® XP may be tank mixed with up to 2 lb ai atrazine* for additional control of many broadleaf weeds, including:

	<u>Maximum Height or Diameter</u>
Sicklepod	2"
Prickly sida	2"
Wild radish	12"
Cutleaf evening primrose	6"
Florida pusley	2"

* Products containing atrazine are restricted use products.

AccentACCENT® XP + atrazine tank mixtures may result in reduced control of grasses (antagonism) if applied to grasses under low moisture stress or to grasses exceeding the maximum labeled height. Before applying AccentACCENT® XP + atrazine tank mixtures, refer to the atrazine product label for information regarding the maximum amount of atrazine that may be applied in a year.

Tank Mixtures with Mesotrione + Atrazine”

AccentACCENT® XP may be tank mixed with mesotrione herbicide for control of the following weeds:

Cocklebur
Dandelion
Jimsonweed
Kochia
Lambsquarters, common
Morningglory, ivyleaf
Mustard, wild
Nightshade, black, Eastern black
Pigweed, palmer
Pigweed, redroot, smooth
Ragweed, common
Ragweed, giant
Smartweed, ladysthumb
Smartweed, Pennsylvania
Sunflower, common
Velvetleaf
Waterhemp, tall

and common

For improved grass and broadleaf weed control, [AccentACCENT®](#) XP tank mixtures with mesotrione (with or without atrazine) may be applied a petroleum-based crop oil concentrate (COC) + an ammonium nitrogen fertilizer. Refer to the mesotrione herbicide label for additional information regarding specific weed control, tank mixtures, adjuvants and rotational crops.

Tank Mixtures with Topramezone” + Atrazine

[AccentACCENT®](#) XP may be tank mixed with topramezone herbicide plus atrazine for control of the following weeds:

Amaranth, Palmer
Cocklebur, common
Jimsonweed
Kochia
Lambsquarter, common
Morningglory, annual
Mustard, wild
Nightshade (black, Eastern black)
Pigweed (redroot, smooth)
Ragweed, common
Ragweed, giant
Smartweed, Pennsylvania
Smartweed, Ladysthumb
Sunflower, common
Thistle, Canada
Velvetleaf
Waterhemp (tall, common)

Refer to the topramezone label for additional information regarding specific weed control, tank mixtures, adjuvants and rotational crops.

Tank Mixtures with glyphosate

Glyphosate may be tank mixed with postemergence applications of [AccentACCENT®](#) XP at 0.67 oz (0.031 lb ai) per acre when made to glyphosate-resistant corn hybrids. Refer to the Spray Adjuvants section for additional information on proper adjuvant selection.

Tank Mixtures with glufosinate

Glufosinate may be tank mixed with applications of [AccentACCENT®](#) XP at 0.67 oz (0.031 lb ai) per acre if applications are made to glufosinate-resistant corn hybrids. Consult with your seed supplier to confirm the corn hybrid is glufosinate-resistant before applying any herbicide containing glufosinate.

For Additional Control of Later Emerging Grasses

[AccentACCENT®](#) XP may be tank mixed with full or reduced rates of preemergence grass herbicides labeled for early postemergence application to field corn (including CINCH®, CINCH® ATZ, Keystone® NXT, Keystone® LA NXT, Resicore®, SureStart®, Surpass® NXT, pendimethalin or dimethenamid) for residual activity on later-emerging flushes of grass.

Application must be made before the grass emerges and before other grass weeds on the [AccentACCENT®](#) XP label exceed their labeled sizes.

The use of nonionic surfactant is advised in place of crop oil concentrate for tank mixtures with preemergence grass herbicides where applications are made early postemergence to small grass weeds. See SPRAY ADJUVANTS for adjuvant rate specifications.

When tank mixing [AccentACCENT®](#) XP with EC formulated preemergence grass herbicides including CINCH® or pendimethalin, **DO NOT** add mesotrione herbicides to the tank mixture. When other formulations of preemergence grass herbicides are tank mixed with [AccentACCENT®](#) XP + “mesotrione, including CINCH® ATZ or Keystone® NXT, limit preemergence herbicide.

Tank mixes of [AccentACCENT®](#) XP and preemergence grass herbicides must be broadcast applied postemergence to field corn before the crop exceeds the heights listed on the preemergence grass herbicide label. Refer to WHEN TO APPLY- POSTEMERGENCE and the preemergence grass herbicide label for complete postemergence application information, rates, and restrictions.

Tank Mixtures with Insecticides

[AccentACCENT®](#) XP may be tank mixed with diamide, pyrethroid or carbamate insecticides including LANNATE® brand insecticides. See **RESTRICTIONS (Corn) and PRECAUTIONS sections** for information on use of [AccentACCENT®](#) XP following soil insecticides application. It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

Other Tank Mixtures

Other than the exceptions noted, and in addition to the tank mix partners and rates indicated above, [AccentACCENT®](#) XP may be tank mixed or followed with sequential applications of other products registered for use in field corn. Applications of full or reduced rates of other products registered for use in corn provided:

- The tank mix product is labeled for the same timing, method of application, adjuvants, and use restrictions as [AccentACCENT®](#) XP.
- The tank mixture is not specifically prohibited on the label of the tank mix product.
- The tank mix combination is compatible as determined by a “jar test” described in the TANK MIX COMPATIBILITY TESTING section below.

Weed control and crop response with tank mixtures not specified in this label are the responsibility of the user and manufacturer of the tank mix product.

A corn plant's predisposition to develop fused tissue emerging from the whorl (rattail) after the V-11 stage may increase when a product containing dicamba is applied to small corn under early stressful conditions. Be aware of this when applying tank mixes with dicamba to small corn (V-3 stage or smaller) under stressful conditions. See ENVIRONMENTAL CONDITIONS for a description of these stressful conditions.

To avoid crop injury or antagonism, apply the products indicated below at least seven days before or three days after the application of [AccentACCENT®](#) XP.

- **DO NOT** tank mix [AccentACCENT®](#) XP with bentazon or severe crop injury may occur.
- **DO NOT** tank mix [AccentACCENT®](#) XP with 2,4-D -containing products as severe grass control antagonism may occur.
- **DO NOT** tank mix [AccentACCENT®](#) XP with foliar-applied organophosphate insecticides including chlorpyrifos, malathion, parathion, etc., as severe crop injury may occur.

DO NOT exceed labeled application rates. **DO NOT** tank mix [AccentACCENT®](#) XP with other products that contain the same active ingredients as [AccentACCENT®](#) XP (nicosulfuron) unless the label of either tank mix partner specifies the maximum rate that may be used.

SEQUENTIAL ACCENT® XP APPLICATIONS

Annual grasses may have more than one flush of emerging seedlings. Also, regrowth of treated annual grasses may occur due to adverse environmental conditions following application. Perennial grasses may regrow from underground stems or roots, depending upon environmental conditions. To control grasses under these conditions, a sequential application of [AccentACCENT®](#) XP may be necessary.

CULTIVATION

A timely cultivation may be necessary to control suppressed weeds or weeds that emerge after an application of [AccentACCENT®](#) XP.

Optimum timing for cultivation is 7–14 days after [AccentACCENT®](#) XP application or upon seeing the establishment of new weeds.

GRAZING/ PREHARVEST INTERVALS FOR CORN

DO NOT graze or feed forage, hay, or straw from treated areas to livestock within 30 days of [AccentACCENT®](#) XP application.

CROP ROTATION

Rotational crops vary in their response to low concentrations of [AccentACCENT®](#) XP remaining in the soil. [AccentACCENT®](#) XP dissipates rapidly in warm, acidic, microbiologically active soils.

The amount of [AccentACCENT®](#) XP which may be present in the soil depends on application rate, soil pH and organic matter content, elapsed time since application, crop production practices, and environmental factors.

Injury to rotational crops may occur in high-pH, cold soils if dry weather prevails between application and

rotational crop planting. Consult your local [DuPont company](#) representative for additional guidelines.

Determine soil pH by laboratory analysis using the 1:1 soil: water suspension method on representative soil samples taken at 0–4" depth. Soil pH varies within fields; therefore, base recropping on the highest soil pH within each field. Consult local extension publications for specified soil sampling procedures.

Observe the following rotational intervals when using [AccentACCENT®](#) XP at a maximum of 1.33 ounces (0.062 lb ai):

DUPONT™-ACCENT® XP ROTATIONAL CROP GUIDELINE – 1

No soil pH restrictions		
Rotational Crop	Interval in Months	
Corn (field, pop, seed, sweet*)	Anytime	
Soybeans	0.5 (15 days)	
Cereals, spring (barley, oats, rye, wheat)	8	
Cereals, winter (barley, oats, rye wheat)	4	
Cotton	10	
Dry beans, Peas, Snap beans	10	
Alfalfa**	12	
Red Clover**	12	
Sorghum (All types including hybrids containing the INZEN™ trait)	18	
Other Crops	See rotational Crop guidelines 2 and 3	

* Except the sweet corn varieties "Merit", "Carnival", and "Sweet Success", for which the minimum time interval is 15 months.

**Except for the state of Kansas east of Highway 75, for Minnesota east and south of the Red River Valley and for the states east of the line formed by the western borders of Iowa, Missouri, Arkansas, and Louisiana, where the minimum time interval is 10 months.

ACCENT® XP ROTATIONAL CROP GUIDELINE – 2

With soil pH < 7.5 restrictions		
Crop	Rotational Interval in Months	
	pH 7.5	pH >7.5
Sunflowers*	11	18
All other crops not listed in Rotational guidelines 1 or 2	See Rotational Guidelines 3	

*Precipitation following application must exceed 14" prior to planting sunflowers.

ACCENT® XP ROTATIONAL CROP GUIDELINE – 3

With soil pH < 6.5 restrictions		
Crop	Rotational Interval in Months	
	pH 6.5	pH >6.5
Sugarbeets*, Potatoes**	10	18***
All other crops not listed in Rotational Guidelines 1 or 2	10	18

* Except on irrigated sites in Colorado, Wyoming, Nebraska, Texas, Michigan, and Ohio, where precipitation following application must exceed 25" prior to planting beets, where the interval is 10 months on soils with pH < 7.5. Sites in Minnesota east and south of the Red River Valley may follow these guidelines provided maximum rates of [AccentACCENT®](#) XP **DO NOT** exceed 0.67 ounces (0.031 lb ai).

**Irrigated potatoes following irrigated corn treated in the States of Washington, Oregon, Idaho, or Utah can be planted 10 months after using [AccentACCENT®](#) XP on sprinkler irrigated corn with no soil pH restrictions, providing the maximum use rate on corn does not exceed 1.0 ounce product (0.046 lb ai) per year. Corn treated with [AccentACCENT®](#) XP must be grown to maturity and receive a minimum of 18 inches of irrigation water before potatoes can be planted at this rotation interval. Injury to potatoes may occur if less than 18 inches of irrigation is used on the previous corn crop. [AccentACCENT®](#) XP may not be used in a tankmix or sequential application program with other ALS-inhibiting herbicides including primisulfuron-methyl.

***In North Dakota and northwest Minnesota, the cumulative precipitation in the 18 months following application must exceed 28" in order to rotate to sugarbeets or potatoes.

ROTATIONAL CROP GUIDELINES - 4 may be observed when using a single application of [AccentACCENT®](#) XP per year with a maximum use rate of 0.67 ounces product (0.031 lb ai). Extend rotational intervals to 12 months if drought conditions prevail after application and before the rotational crop is planted, unless sprinkler irrigation has been applied and totals greater than 15" during the growing season.

ACCENT® XP ROTATIONAL CROP GUIDELINES – 4

With 0.67 ounces (0.031 lb ai) maximum use rate	
Crop	Rotational Interval in Months
Alfalfa*	10
Canola	10
Flax**	10
Sorghum (All types including hybrids containing the INZEN™ trait)	18
Potato	10
Red clover	10
Sunflower	10

*On sprinkler irrigated fields in Idaho, Utah, and Northern Nevada it is best to use deep fall tillage including plowing prior to planting alfalfa. Product degradation may be less on furrow irrigated soils and may result in some crop injury.

**Extend rotational intervals to 18 months if drought conditions prevail after application and before the rotational crop is planted, unless sprinkler irrigation has been applied and totals greater than 15" during the growing season.

GROUND APPLICATION

Broadcast Application

- Use a minimum of 15 gallons of water per acre (15 GPA) for best performance. Use a minimum of 10 gallons of water per acre (GPA) for light, scattered stands of weeds.
- Overlaps or starting, stopping, slowing, and turning while spraying may result in crop injury.

Band Application

For band applications, use proportionately less spray mixture, and carefully calibrate the band applicator to not exceed the labeled rate. Carefully follow the manufacturer's instructions for nozzle type (flat fans), orientation, distance of nozzles from the crop and weeds, spray volumes, calibration and spray pressure.

Refer to MANDATORY SPRAY DRIFT MANAGEMENT section

AERIAL APPLICATION

In New York state and California aerial application is not permitted.

Use nozzle types and arrangements that will provide optimum spray distribution and maximum coverage at a minimum of 3 GPA.

Refer to MANDATORY SPRAY DRIFT MANAGEMENT section

SPRAY ADJUVANTS FOR USE IN FIELD CORN GROWN FOR GRAIN & SEED, POPCORN, & SWEET CORN

Applications of [AccentACCENT®](#) XP must include either a crop oil concentrate or a nonionic surfactant. In addition, an ammonium nitrogen fertilizer must be used unless specifically prohibited by tank mix partner labeling. Crop oil concentrate plus ammonium nitrogen fertilizer is the preferred adjuvant system

for activity on difficult to control species including woolly cupgrass, quackgrass, sandbur and wirestem muhly. Consult local [DuPont company](#) fact sheets, technical bulletins, and service policies prior to using other adjuvant systems. If another herbicide is tank mixed with [Accent ACCENT® XP](#), select adjuvants authorized for use with both products. Products must contain only EPA-exempt ingredients (40 CFR 1001). It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

When applied in tank mix combination with a glyphosate containing product that contains a built-in adjuvant, ensure the total adjuvant load is equivalent to the directions on this label.

Crop Oil Concentrate (COC) - Petroleum or Modified Seed Oil (MSO)

- Apply up to 1% v/v (1 gallon per 100 gallons spray solution) under arid conditions.
- MSO adjuvants may be used up to 0.5% v/v (0.5 gallons per 100 gallons spray solution) if specifically noted on adjuvant product labeling or if specified on local [DuPont company](#) product literature or service policies.
- Oil adjuvants must contain at least 80% high quality, petroleum (mineral) or modified vegetable seed oil with at least 15% surfactant emulsifiers.

Nonionic Surfactant (NIS)

- Apply up to 0.25% v/v (1 quart per 100 gallons spray solution) under arid conditions.
- Surfactant products must contain at least 60% nonionic surfactant with a hydrophilic/lipophilic balance (HLB) greater than 12.

Ammonium Nitrogen Fertilizer

- Use 2 quarts/acre of a high-quality urea ammonium nitrate (UAN), including 28%N or 32%N, or 2 pounds/acre of a spraygrade ammonium sulfate (AMS). Use 4 quarts/acre UAN or 4 pounds/acre AMS under arid conditions.
- **DO NOT** use liquid nitrogen fertilizer as the total carrier solution.

Special Adjuvant Types

- Combination adjuvant products may be used at doses that provide the required amount of NIS, COC, MSO and/or ammonium nitrogen fertilizer. Consult product literature for use rates and restrictions.
- In addition to the adjuvants specified above, other adjuvant types may be used if they provide the same functionality. Consult separate [DuPont company](#) technical bulletins for detailed information before using adjuvant types not specified on this label.

MIXING INSTRUCTIONS

Select a spray volume that will ensure thorough coverage and a uniform spray pattern. If tank mixing with other herbicides, always consult the label of the tank mix partner(s) for minimum spray volume requirements and apply the tank mixture using a water volume advised for all products.

1. Fill the tank 1/4 to 1/3 full of water.

2. While agitating, add the required amount of [DuPont™ AccentACCENT®](#) XP.
3. Continue agitation until the [AccentACCENT®](#) XP is fully dispersed, at least 5 minutes.
4. Once the [AccentACCENT®](#) XP is fully dispersed, maintain agitation and continue filling tank with water. Thoroughly mix [AccentACCENT®](#) XP with water before adding any other material.
5. If tank mixing [AccentACCENT®](#) XP with another herbicide, follow this mixing order: dry flowables (DF) and soluble granules (SG), followed by liquids, then oil dispersions (OD) or emulsifiable concentrates (EC). Maintain continuous agitation.
6. As the tank is filling, add the required spray adjuvants (crop oil concentrate, nonionic surfactant, or ammonium nitrogen fertilizer).
7. If the mixture is not continuously agitated, settling will occur. If settling occurs, thoroughly re-agitate before using.
8. Apply [AccentACCENT®](#) XP spray mixture within 24 hours of mixing to avoid product degradation.
9. If [AccentACCENT®](#) XP and a tank mix partner are to be applied in multiple loads, pre-slurry the [AccentACCENT®](#) XP in clean water prior to adding to the tank. This will prevent the tank mix partner from interfering with the dissolution of the [AccentACCENT®](#) XP.

TANK MIX COMPATIBILITY TESTING

Perform a jar test prior to tank mixing to ensure compatibility of [AccentACCENT®](#) XP and other pesticides. Use a clear glass quart jar with lid and mix the tank mix ingredients in their relative proportions. Invert the jar containing the mixture several times and observe the mixture for approximately 1/2 hour. If the mixture balls-up, forms flakes, sludges, gels, oily films or layers, or other precipitates, it is not compatible. **DO NOT** use the tank mix combination.

ENVIRONMENTAL CONDITIONS AND BIOLOGICAL ACTIVITY

[AccentACCENT®](#) XP provides best results when applied to young, actively growing weeds. Applications made during warm, moist conditions (70°F or more) and adequate soil moisture both before and after application maximizes performance.

The degree and duration of control depend on spray coverage, weed spectrum, weed size, growing conditions before and after treatment, soil moisture, and adjuvant selection.

[AccentACCENT®](#) XP is rainfast in 4 hours.

Treating weeds that exceed maximum label height or that are under stress may result in incomplete control. Poor weed control or crop injury may result from applications made to plants under stress from:

- abnormally hot or cold weather
- environmental conditions including drought, water-saturated soils, hail damage, or frost
- disease, insect, or nematode injury
- prior herbicide or carryover from a previous year's herbicide application

Severe stress from conditions preceding or immediately following application may also result in crop injury

or poor weed control. Stress affects all weeds, but especially weeds including woolly cupgrass, green and yellow foxtail, and wild proso millet.

If the crop or grass weeds are under stress, delay application until stress passes and both weeds and crop resume active growth.

[AccentACCENT®](#) XP rapidly inhibits the growth of susceptible weeds, reducing weed competition within as little as 6 hours after application. Susceptible plants are controlled in 7–21 days.

DO NOT exceed labeled application rates. **DO NOT** tank mix [AccentACCENT®](#) XP with other products that contain the same active ingredients as [AccentACCENT®](#) XP (nicosulfuron) unless the label of either tank mix partner specifies the maximum rate that may be used.

If the crop or grass weeds are under stress, delay application until stress passes and both weeds and crop resume active growth.

SPRAYER PREPARATION/ CLEANUP

It is important that spray equipment is clean and free of previous pesticide deposits before using [AccentACCENT®](#) XP and then properly cleaned out following application. Clean all application equipment before applying [AccentACCENT®](#) XP. Follow the cleanup procedures specified on the label of the product previously sprayed. If no cleanup procedure is provided, use the procedure that follows.

Immediately following applications of [AccentACCENT®](#) XP, thoroughly clean all mixing and spray equipment to avoid subsequent crop injury.

Note:

- When cleaning spray equipment before applying [AccentACCENT®](#) XP, read and follow label directions for proper rinsate disposal of the product previously sprayed.
- Steam cleaning of aerial spray tanks will help to dislodge any visible pesticide deposits.
- When spraying or mixing equipment will be used over an extended period to apply multiple loads of [DuPont™ AccentACCENT®](#) XP, partially fill the tank with fresh water at the end of each day of spraying, flush the boom and hoses, and allow to sit overnight.

Cleanup Procedure

1. Drain the tank and thoroughly hose down the interior surfaces. Flush the tank, hoses, and boom with clean water for a minimum of 5 min.
2. Partially fill the tank with clean water and add one gallon of household ammonia* (containing 3% active) for every 100 gallons of water. Finish filling the tank with water, then flush the cleaning solution through the hoses, boom, and nozzles. Add more water to completely fill the tank and allow to agitate/recirculate for at least 15 min. Again, flush the hoses, boom, and nozzles with the cleaning solution, then drain the tank.
3. Repeat Step 2.
4. Remove the nozzles, screens and the end caps of sprayer booms and clean separately in a bucket containing the cleaning agent and water.
5. Thoroughly rinse the tank with clean water for a minimum of 5 min, flushing the water through the hoses and boom.

* Equivalent amounts of an alternate strength ammonia solution or a tank cleaner may be used.

USEPA REGISTERED PRODUCTS MENTIONED IN THIS LABEL FOR USE IN TANK MIXTURES OR OTHER REASONS		
PRODUCT BRAND NAME	ACTIVE INGREDIENT(S)	EPA REGISTRATION NUMBER
Abundit® Edge	glyphosate	352-922
Basis® Blend	rimsulfuron + thifensulfuron methyl	352-854
Cinch®	s-metolachlor	352-625
Cinch® ATZ	atrazine + S-metolachlor + atrazine related compounds	352-624
Cinch® ATZ Lite	atrazine + S-metolachlor + atrazine related compounds	352-623
Instigate®	mesotrione + rimsulfuron	352-873
Keystone® LA NXT	acetochlor + atrazine	62719-670
Keystone®	acetochlor + atrazine	62619-671
Lannate® SP	methomyl	352-342
Lannate® LV	methomyl	352-384
Leadoff®	rimsulfuron + thifensulfuron methyl	352-853
Prequel®	isoxaflutole + rimsulfuron	352-779
Resicore®	acetochlor + clopyralid + mesotrione	62719-693
Resolve® Q	rimsulfuron + thifensulfuron methyl	352-777
SureStart® II	acetochlor + clopyralid + flumetsulam	62719-679
Surpass® NXT	acetochlor	62719-672

NOTICE TO BUYER: Purchase of this material does not confer any rights under patents of countries outside of the United States.

LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read this Limitation of Warranty and Liability before buying or using This Product. If the Terms Are Not Acceptable, Return the Product at once, unopened, and the Purchase Price Will be Refunded.

It is impossible to eliminate all risks associated with the use of this product. Such risks arise from weather conditions, soil factors, off target movement, unconventional farming techniques, presence of other materials, the manner of use or application, or other unknown factors, all of which are beyond the control of DuPont. These risks can cause: ineffectiveness of the product, crop injury, or injury to non-target crops or plants. WHEN YOU BUY OR USE THIS PRODUCT, YOU AGREE TO ACCEPT THESE RISKS.

DuPont warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for the purpose stated in the Directions for use, subject to the inherent risks described above, when used in accordance with the Directions for use under normal conditions.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, DUPONT MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR OF MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL DUPONT OR SELLER BE LIABLE FOR ANY INCIDENTAL, CONSEQUENTIAL OR SPECIAL

~~DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. BUYER'S OR USER'S BARGAINED-FOR EXPECTATION IS CROP PROTECTION. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER AND THE EXCLUSIVE LIABILITY OF DUPONT OR SELLER, FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY OR CONTRACT, NEGLIGENCE, TORT OR STRICT LIABILITY), WHETHER FROM FAILURE TO PERFORM OR INJURY TO CROPS OR OTHER PLANTS, AND RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT, OR AT THE ELECTION OF DUPONT OR SELLER, THE REPLACEMENT OF THE PRODUCT.~~

~~To the extent consistent with applicable law that allows such requirement, DuPont or its Ag Retailer must have prompt notice of any claim so that an immediate inspection of buyer's or user's growing crops can be made. buyer and all users shall promptly notify DuPont or a DuPont Ag Retailer of any claims, whether based on contract, negligence, strict liability, other tort or otherwise, or be barred from any remedy.~~

~~This Limitation of Warranty and Liability may not be amended by any oral or written agreement.~~

Terms and Conditions of Use

If terms of the following Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid. To the extent consistent with applicable law, use by the buyer or any other user constitutes acceptance of the terms under Warranty Disclaimer, Inherent Risks of Use and Limitations of Remedies.

Warranty Disclaimer

Corteva Agriscience warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions for use, subject to the inherent risks set forth below. To the extent consistent with applicable law, Corteva Agriscience MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

Inherent Risks of Use

It is impossible to eliminate all risks associated with use of this product. Plant injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperatures, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application or other factors, all of which are beyond the control of Corteva Agriscience or the seller. To the extent consistent with applicable law, Corteva Agriscience will not be responsible for losses or damages resulting from the use of this product in any manner not specifically directed by Corteva Agriscience. To the extent consistent with applicable law, all such risks associated with non-directed use shall be assumed by buyer and/or user.

Limitation of Remedies

To the extent consistent with applicable law, the exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, tort, strict liability, or other legal theories), shall be limited to, at Corteva Agriscience's election, one of the following:

1. Refund of purchase price paid by buyer or user for product bought, or
2. Replacement of product used.

To the extent consistent with applicable law, Corteva Agriscience shall not be liable for losses or damages resulting from handling or use of this product unless Corteva Agriscience is promptly notified of such loss or damage in writing. To the extent consistent with applicable law, in no case shall Corteva Agriscience be liable for consequential, incidental or special damages or losses.

The terms of the Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of Corteva Agriscience or the seller is authorized to vary or exceed the terms of the Warranty Disclaimer or this Limitation of Remedies in any manner.

For product information call: 1-800-258-3033

Produced for
Corteva Agriscience LLC
9330 Zionsville Road
Indianapolis, IN 46268~~E. I. du Pont de Nemours and Company, Chestnut Run Plaza, 974 Centre Road, Wilmington, DE 19805 U.S.A.~~

~~™Trademarks of Corteva Agriscience and its affiliated companies[®]™Trademarks of Dow AgroSciences, DuPont or Pioneer and their affiliated companies or respective owners~~

EPA accepted 07/13/2020