

352-835

2/29/2012

10f 11



U S ENVIRONMENTAL PROTECTION AGENCY

Office of Chemical Safety and Pollution Prevention
Office of Pesticide Programs
Registration Division (7504P)
1200 Pennsylvania Ave, N W
Washington, DC 20460

EPA Reg Number

352-835

Date of Issuance

FEB 29 2012

Term of Issuance
Unconditional

Name of Pesticide Product
**Dupont™ Velista™
Fungicide**

NOTICE OF PESTICIDE

- Registration
 - Reregistration
- Under FIFRA as amended

Name and Address of Registrant (include ZIP Code)

**DuPont Crop Protection
Stine Haskell Research Center
P O Box 30
Newark, DE 19714**

Mailed to

**Richard H Collier, Ph D
Landis International, Inc
P O Box 5126
Valdosta, GA 31603-5126**

Note Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce In any correspondence on this product always refer to the above EPA registration number

On the basis of information furnished by the registrant, the above named pesticide is hereby registered under the Federal Insecticide, Fungicide and Rodenticide Act Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others

This product is registered in accordance with FIFRA section 3(c)(5) provided that you

- 1 Submit and/or cite all data required for registration of your product under FIFRA sec 3(c)(5) when the Agency requires all registrants of similar products to submit such data, and submit acceptable responses required for reregistration of your product under FIFRA section 4

Page 1 of 2

Signature of Approving Official

Date


FEB 29 2012

Tony Kish, Product Manager (22)
Fungicide Branch/Registration Division/OPP/OCSP (7504P)

- 2 Make the following change to the label
 - a Change the product registration number to 'EPA Reg No 352-835'
- 3 Submit one copy of the revised final printed label for the record before the product is released for shipment

Your release for shipment of the product constitutes acceptance of these conditions


A copy of the label stamped "Accepted" is enclosed for your records



Tony Kish
Product Manager (22)
Fungicide Branch
Registration Division (7504P)

Enclosure

Label stamped 'Accepted'
Product Chemistry Review DP373541
Acute Toxicity Review DP373543



Professional Products

GROUP	7	FUNGICIDE
-------	---	-----------

ACCEPTED
FEB 29 2012
 Under the Federal Insecticide Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg No 352-835

INTENDED FOR USE BY COMMERCIAL APPLICATORS ONLY

A water dispersible granular fungicide for turf disease control on golf courses residential institutional commercial and municipal lawns park and recreation areas and sod farms and for disease control on landscape flowering plants and trees

Active Ingredient	By Weight
Penthiopyrad	50%
Inert Ingredients	50%
TOTAL	100%

EPA Reg No 352 835 EPA Est No _____

[Refillable]

or

[Nonrefillable]

Net Contents _____

E I du Pont de Nemours and Company
 1007 Market Street
 Wilmington Delaware 19898

FIRST AID

IF IN EYES Hold eye open and rinse slowly and gently with water for 15 20 minutes Remove contact lenses if present after the first 5 minutes then continue rinsing Call a poison control center or doctor for treatment advice

IF SWALLOWED Call a poison control center or doctor immediately for treatment advice Have person sip a glass of water if able to swallow Do not induce vomiting unless told to do so by the poison control center or a doctor Do not give anything by mouth to an unconscious person

Have the product container or label with you when calling a poison control center or doctor or going for treatment You may also contact 1 800 441 3637 for emergency medical treatment information

For information on the use of this product call DuPont Professional Products at 1 888 6 DUPONT (1 888 638 7668)

PERSONAL PROTECTIVE EQUIPMENT

- Mixers/loaders/applicators and other handlers must wear
- Long sleeved shirt
 - Long pants
 - Shoes plus socks

Follow manufacturer s instructions for cleaning/ maintaining PPE If no such instructions for washables are available use detergent and hot water Keep and wash PPE separately from other laundry

ENGINEERING CONTROL STATEMENTS

When handlers use closed systems or enclosed cabs in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170 240(d)(4 6)] the handler PPE requirements may be reduced or modified as specified in the WPS

KEEP OUT OF REACH OF CHILDREN CAUTION

Si usted no entiende la etiqueta busque a alguien para que se la explique a usted en detalle (If you do not understand the label find someone to explain it to you in detail)

Refer to back/side panels for additional precautionary statements and First Aid

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Causes moderate eye irritation Avoid contact with eyes or clothing May be harmful if swallowed Wash thoroughly with soap and water after handling and before eating drinking chewing gum using tobacco or using the toilet



The miracles of science™

USER SAFETY RECOMMENDATIONS

Users should wash hands before eating drinking chewing gum using tobacco or using the toilet Users should remove clothing/PPE immediately if pesticide gets inside Then wash thoroughly and put on clean clothing Users should remove PPE immediately after handling this product Wash the outside of gloves before removing As soon as possible wash thoroughly and change into clean clothing

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish For terrestrial uses Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark Do not contaminate water when disposing of equipment washwater or rinsate Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas

This product may impact surface water quality due to runoff of rain water This is especially true for poorly draining soils and soils with shallow ground water This product is classified as having high potential for reaching surface water via runoff several weeks after application A level well maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds streams and springs will reduce the potential loading of this chemical from runoff water and sediment Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 48 hours

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling

Read entire label before using this product

DuPont™ VELISTA™ fungicide (i.e. VELISTA™) must be used only in accordance with specifications on this label DuPont will not be responsible for losses or damages resulting from the use of this product in any manner not specifically recommended by DuPont User assumes all risks associated with such non specified use

Do not formulate this product into other end use products without written permission of DuPont Professional Products

RESTRICTIONS [optional restrictions based on use sites included]

Do not apply this product in commercial nurseries and greenhouses

[Not for use on ornamental plants being grown for sale or other commercial use or for commercial seed production]

[Do not use this product on any bearing fruit and nut trees]

Do not apply this product through any type of irrigation system

- Do not apply more than 2.2 oz of VELISTA™ per 1 000 sq ft per year or 6 lbs of VELISTA™ (equivalent to 3 lbs of active ingredient) per acre per year
 - Keep people and pets away from areas treated with VELISTA™ until treated area has dried
- Do not apply this product using aerial application

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are not within the scope of the Worker Protection Standard (WPS) for agricultural pesticides 40 CFR Part 170 The WPS applies when this product is used to produce agricultural plants on farms forests nurseries and greenhouses

Professional applications to golf courses residential industrial institutional and commercial lawns/landscape areas and sports fields are not within the scope of the Worker Protection Standard

For applications to turfgrass do not enter or allow others to enter the treated area until treatments have dried

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard 40 CFR Part 170 This standard contains requirements for the protection of agricultural workers on farms (sod farms included) forests nurseries and greenhouses and handlers of agricultural pesticides It contains requirements for training decontamination notification and emergency assistance It also contains specific instructions and exceptions pertaining to the statements on the label about personal protective equipment (PPE) and restricted entry interval and notification to workers (as applicable) The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated such as plants soil or water is

- Coveralls
- Shoes and socks
- Chemical resistant gloves (made of any waterproof material)

PRODUCT INFORMATION

VELISTA™ fungicide is a water dispersible granule applied as a foliar or broadcast treatment to control listed diseases on the landscape flowering plants turf/turf use sites and non bearing fruit/nut trees which are specified on this label

VELISTA™ fungicide may be used on landscape flowering plants and trees or turfgrasses that are being grown for aesthetic or recreational purposes or climatic modification in or around home lawns residential dwellings business and office complexes shopping complexes multi family residential complexes institutional buildings airports cemeteries interior plantscapes ornamental gardens parks playgrounds schools day care centers golf courses (tee box areas roughs fairways greens collars) athletic fields and commercial sod farms

The active ingredient in VELISTA™ fungicide is penthiopyrad and it is a carboxamide fungicide Penthiopyrad inhibits fungal respiration by binding to mitochondrial respiratory complex II Penthiopyrad is not cross

resistant with other classes of fungicide chemistry like triazoles and other ergosterol biosynthesis inhibitors stro bilurins and other complex III (QoI) inhibitors anilnopy rimidines dicarboximides benzimidazoles and N phenyl carbamates

DuPont™ VELISTA™ fungicide is in GROUP 7 of the EPA's Fungicide groups based on Target Site of Action and may be used in rotational and mixture resistance management programs. VELISTA™ fungicide must be diluted with water before application according to the following use directions

Not all turf and landscape flowering plant and tree varieties cultivars or hybrids have been individually tested for plant safety. It is not possible to evaluate for plant safety all applications of VELISTA™ on all turf and landscape flowering plants and trees or their varieties cultivars or hybrids or under all environmental conditions and growing circumstances. To test for turf and plant safety apply the product in accordance with the label instructions to a small area of the target turf or landscape flowering plant or tree to ensure that a phytotoxic response will not occur especially where the application is a new use of the product by the applicator.

APPLICATION TIMING

VELISTA™ fungicide is a broad spectrum fungicide recommended for the control of many important listed diseases. Apply VELISTA™ preventatively. Begin applications when conditions favor disease development. Apply curatively according to the instructions in Table 2.

MIXING DIRECTIONS

Apply VELISTA™ fungicide alone or in tank mixes with other registered fungicides. Do not exceed the recommended application rates for VELISTA™ and/or the tank mix partners given in the label(s). Follow the use directions provided in the Resistance Management section of this label. Follow all use directions given in this label. Failure to follow the use directions can result in injury to the plant and/or inadequate disease control.

Basic Mixing Instructions

Important: Application equipment must be clean and free of previous pesticide deposits before mixing VELISTA™ fungicide.

- 1 Use clean, well-maintained application equipment.
- 2 Fill sprayer tank 1/4 to 1/2 full of water.
- 3 Start agitation, then add VELISTA™ directly to the mixing or sprayer tank. Use mechanical or hydraulic agitation.
- 4 Mix thoroughly to fully disperse the VELISTA™ fungicide and continue filling the tank with water while agitating to keep the product in suspension.
- 5 It is recommended that the treatment mixture not be stored in the mixing or spray tank overnight. If treatment mixture sits in the tank for extended periods, re-agitate prior to application.

Tank mixtures

VELISTA™ fungicide may be tank mixed with other pesticides and/or spray additives. When tank mixing VELISTA™ with other pesticides, observe all precautions

and limitations on each separate product label. No label application rates may be exceeded. VELISTA™ fungicide cannot be mixed with any product containing a label prohibition against such mixing. The physical compatibility of VELISTA™ fungicide may vary with different sources of pesticide products and local cultural practices. Any tank mixture that has not been previously tested should be prepared on a small scale (pint or quart jar) using the proper proportions of pesticides and water to ensure the physical compatibility of the mixture and for potential phytotoxicity on turf and landscape ornamentals.

Tank mixing Sequence

Add different formulation types in the sequence indicated below. Allow time for complete mixing and dispersion after the addition of each product.

- 1 Water soluble bags
- 2 VELISTA™ fungicide and other water dispersible granules
- 3 Wettable powders
- 4 Water based suspension concentrates
- 5 Water soluble concentrates
- 6 Oil based suspension concentrates
- 7 Emulsifiable concentrates
- 8 Adjuvants, surfactants, oils
- 9 Soluble fertilizers
- 10 Drift retardants

APPLICATION EQUIPMENT CLEANING

Prior to application, start with clean, well-maintained application equipment. Immediately following application, thoroughly clean all application equipment to reduce the risk of forming hardened deposits that might become difficult to remove. Drain application equipment. Thoroughly rinse application equipment and flush hoses, boom and nozzles with clean water. Clean all other associated application equipment. Take all necessary safety precautions when cleaning equipment. Do not clean equipment near wells, water sources or desirable vegetation. Dispose of waste rinse water in accordance with local regulations.

INTEGRATED PEST MANAGEMENT

DuPont recommends the use of Integrated Pest Management (IPM) programs to control diseases. VELISTA™ may be used as part of an Integrated Disease and Pest Management (IPM) program which can include biological, cultural and genetic practices aimed at reducing potential disease incidence. Follow cultural practices known to reduce disease development. Application of this product should be based on IPM principles and practices including but not limited to field scouting or other detection methods, correct target disease identification, population monitoring, advisory (disease forecasting) programs that recommend treatment timing based on environmental factors favorable for disease development, and treating when disease pressure reaches locally determined action levels. Consult your state cooperative extension service, professional consultants or other qualified authorities to determine the appropriate management, cultural practice and treatment threshold levels for the specific crop, geography and diseases.

RESISTANCE MANAGEMENT

Repeated use of fungicides with the same mode of action for control of specific plant pathogens may lead to selection of resistant strains of fungi and result in a reduction of disease control. Penthiopyrad, the active ingredient in DuPont™ VELISTA™, is one of EPA's Target Site of Action Group 7 fungicides (carboxamides). A disease management program that includes rotation and/or tank mixing with non Group 7 fungicides is essential to reduce the risk of fungicide resistance development. Do not make more than two successive applications of VELISTA™; these two applications should be followed by applications of fungicides from a different FRAC mode of action group. For mixture applications, use the recommended rate of VELISTA™ mixed with a labeled rate of a fungicide from a different FRAC mode of action group. For guidance on a particular disease control situation, consult your Cooperative Extension specialist for the latest information on resistance management in your area.

DIRECTIONS FOR TURFGRASS

Apply in sufficient water to ensure thorough coverage. Apply the recommended rates as indicated in Table 1 in up to 10 gallons of water per 1,000 sq. ft. Apply with a properly calibrated sprayer.

Apply after mowing OR allow treated area to dry before mowing.

- Allow treated area to dry before irrigation.

Evaluate spray additives prior to use for phytotoxicity or antagonism. Label directions are based on data obtained with no additives.

Before use of any fungicide, proper diagnosis of the organism causing the disease is important. Use of diagnostic kits or other means of identification of the disease organism is essential to determine the best control measures.

SPECIFIC USE RATES FOR TURFGRASS DISEASES

- Use the low rate of 0.3 oz/1,000 sq. ft. for preventative and for light to moderate disease situations and for use in tank mixtures.

Use the high rate of 0.5 oz/1,000 sq. ft. for high disease pressure situations and for use in curative situations and for use in tank mixtures.

Do not apply more than 0.7 oz of VELISTA™ per 1,000 sq. ft. per application (equivalent to 0.95 lbs of active ingredient per acre per single application) to turf grass.

Follow recommendations in the resistance management section above for instructions on how to use with successive applications (e.g., no more than 2 applications followed by a different mode of action fungicide) or for use in mixtures (e.g., 0.3 oz/1,000 sq. ft. of VELISTA™ and the recommended label rate of the mixture partner).

For broader spectrum control, VELISTA™ can be tank mixed with other fungicides. Check compatibility before tank mixing. Observe all directions, precautions, and limitations on labeling of all products used in tank mixes. Tank mixtures or other applications of products referenced on this label are permitted only in those states in which the referenced products are registered. Do not combine VELISTA™ in the spray tank with pesticides, surfactants, or fertilizers unless your prior use has shown the combination is physically compatible.

**Table 1
APPLICATION DILUTION CHART FOR USE OF VELISTA™ IN TURFGRASS AREAS**

Application Volume (Gallons Water per 1,000 Square Feet)	Application Rate (Ounces of Product per Acre)	Application Rate (Ounces of Product per 1,000 Square Feet)	Application Rate (Lb of A.I. per Acre)	Ounces of VELISTA™ diluted to these volumes of finished spray			
				1 gallon	5 gallons	10 gallons	100 gallons
1	13.1	0.3	0.41	0.3	1.5	3.0	30.0
	21.8	0.5	0.68	0.5	2.5	5.0	50.0
	30.5	0.7	0.95	0.7	3.5	7.0	70.0
2	13.1	0.3	0.41	0.15	0.75	1.5	15.0
	21.8	0.5	0.68	0.25	1.25	2.5	25.0
	30.5	0.7	0.95	0.35	1.75	3.5	35.0
3	13.1	0.3	0.41	0.1	0.5	1.0	10.0
	21.8	0.5	0.68	0.17	0.85	1.7	17.0
	30.5	0.7	0.95	0.23	1.15	2.3	23.0
4	13.1	0.3	0.41	0.075	0.375	0.75	7.5
	21.8	0.5	0.68	0.125	0.625	1.25	12.5
	30.5	0.7	0.95	0.175	0.875	1.75	17.5

To convert from ounces to grams, multiply by 28.35.

1 ounce = 28.35 grams

Do not use household utensils to measure VELISTA™.

Table 2
SPECIFIC USE RATES FOR DUPONT™ VELISTA™ FUNGICIDE TO CONTROL DISEASES OF TURFGRASS

Disease	Applications Sites	Ounces of VELISTA™/ 1,000 sq ft	Application Interval (days)	Comments
Dollar Spot (<i>Sclerotinia homeocarpa</i>)	Golf course tees and greens	0.3 - 0.5	14	Begin application prior to or at the early stages of disease development. Use the high rate for high disease pressure and/or for curative situations. When using 0.3 oz / 1,000 sq ft rate, apply as a tank mix using a different FRAC group fungicide approved for this use site and disease.
	Golf course fairways All other turfgrass	0.3 - 0.5	14 - 21	
Brown Patch (<i>Rhizoctonia solani</i>)	Golf course tees and greens	0.3 - 0.5	14	Begin application at the early stages of disease development. Use high rate for high disease pressure or for curative situations.
	Golf course fairways All other turfgrass	0.3 - 0.5	14 - 21	
Anthracnose (<i>Colletotrichum cereale</i> <i>Colletotrichum graminicola</i>)	All turfgrass	0.3 - 0.5	14	Begin application at the early stages of disease development; consider using tank mixtures with chlorothalonil or strobilurin fungicides at label rates under heavy disease situations. Use high rate for high disease pressure or for curative situations.
Large Patch (<i>Rhizoctonia solani</i>)	All turfgrass	0.7	14 - 28	Make 1 - 2 applications in the fall as a preventative. Apply an early spring application if disease pressure is high.
Leaf spots (such as <i>Bipolaris spp.</i>)	All turfgrass	0.3 - 0.5	14	Begin application at the early stages of disease development. Use high rate for high disease pressure or for curative situations.
Red Thread (<i>Laetisaria fuciformis</i>) Pink Patch (<i>Limonomyces roseipellis</i>)	All turfgrass	0.3 - 0.5	14	Apply when conditions are favorable for disease development.
Powdery Mildew (<i>Erysiphe graminis</i>) Rust (<i>Puccinia graminis</i>)	All turfgrass	0.3 - 0.5	14	Apply when conditions are favorable for disease development.
Gray Snow Mold (<i>Typhula spp.</i>) Pink Snow Mold (<i>Microdochium nivale</i>)	All turfgrass	0.7	NA (only 1 application)	When using 0.7 oz / 1,000 sq ft rate, apply as a tank mix using a different FRAC group fungicide approved for this use site and disease.

DIRECTIONS FOR LANDSCAPE FLOWERING PLANTS AND TREES

- DuPont™ VELISTA™ controls key diseases on landscape ornamental plants and other landscape plantings

Applications of VELISTA™ will control powdery mildews rusts leafspots scabs and other listed diseases of landscape ornamental plants and trees

Guidance on How to Use Tables 3a and 3b for Developing Disease Control Programs In Landscape Plantings

- 1 Use VELISTA™ fungicide in a preventative disease control program
- 2 To determine the use directions for controlling a disease on a landscape plant species select the plant species and target disease in Table 3a (Landscape Trees and Non Bearing Fruit/Nut Tree Diseases) or Table 3b (Landscape Herbaceous Plants)
- 3 The treatment use directions for each table are given at the bottom of each table
- 4 Allow treated areas to dry before watering any ornamental plants growing in landscaped areas
- 5 Optimum benefit of VELISTA™ is obtained when used in conjunction with sound disease management practices

Specific Treatment Rate Recommendations for Landscape Flowering Plants and Trees

VELISTA™ may be used at rates of 6 - 8 oz of VELISTA™/100 gallons of water every 7 - 14 days For most effective control begin applications with VELISTA™ before disease development

Do not apply more than 8 oz of VELISTA™ per single application in 100 gallons of water per acre

Do not apply more than 6 lbs of VELISTA™ fungicide product (equivalent to 3 lb of active ingredient) per acre per year

Plant tolerances to VELISTA™ have been found acceptable for the specific genera and species of plants listed under the Directions for Use

Other plant species may be sensitive to VELISTA™ and diseases other than those listed may not be controlled Before using VELISTA™ on plants or for diseases that are not listed in the Directions for Use test VELISTA™ on a small scale basis first Apply the recommended rates for a particular type of disease i.e. rust powdery mildew etc and evaluate for phytotoxicity and disease control prior to widespread use

Table 3a Landscape Trees and Non Bearing Fruit/Nut Tree Diseases

Landscape Trees and Non Bearing Fruit/Nut Trees	Target Diseases
Landscape Trees	Refer to diseases below
Non bearing Pome fruits Apple crabapple loquat mayhaw pear pear oriental quince	Alternaria leafspots <i>Alternaria spp</i> Scab apple or pear <i>Venturia inaequalis</i> <i>Venturia pirina</i> Powdery mildew <i>Podosphaera leucotricha</i> Rusts <i>Gymnosporangium spp</i>
Non bearing Stone fruits Apricot cherry sweet cherry tart nectarine peach plum plum Chickasaw plum Damson plum Japanese plumcot	Brown rot blossom blight <i>Monilinia spp</i> Cherry leafspot <i>Blumeriella jaapu</i> Powdery mildew rusty spot on peaches <i>Podosphaera clandestina</i> <i>Sphaerotheca pannosa</i> Rust <i>Tranzschelia discolor</i> Shot hole <i>Wilsonomyces carpophilus</i>
Non bearing Tree nuts Almond beech nut Brazil nut butternut cashew chestnut chinquapin filbert (hazelnut) hickory nut macadamia nut walnut black and English pistachio	Alternaria leafspot blight <i>Alternaria spp</i> Anthracnose <i>Colletotrichum spp</i> Brown rot blossom blight <i>Monilinia spp</i> Botrytis blights <i>Botrytis cinerea</i> Panicle and shoot blight <i>Botryosphaeria dothidea</i> Powdery mildew <i>Podosphaera tridactyla var tridactyla</i> <i>Sphaerotheca pannosa</i> <i>Phyllactinia angulata</i> <i>Phyllactinia guttata f sp coryli</i> <i>Microsphaera spp</i> <i>Oidium spp</i> Rust <i>Tranzschelia discolor</i> <i>Uromyces spp</i> <i>Pucciniastrum coryli</i> Scab <i>Cladosporium carpophila</i> Sclerotinia shoot blight <i>Sclerotinia sclerotiorum</i> Seedling blight <i>Rhizoctonia solani</i> Septoria leafspot <i>Septoria spp</i> Shot hole <i>Wilsonomyces carpophilu</i>

Specific Treatment Rates and Application Information for Control of Diseases of Landscape Trees (including Non Bearing Fruit/Nut Trees)

Mix 8 oz of VELISTA™ in 100 gals of water and apply 100 gallons per acre as a full coverage spray to the point of drip

Apply every 7 - 14 days during the period of primary infection or when conditions are favorable for disease development

Do not apply more than 8 oz of VELISTA™ per single application in 100 gallons of water per acre

- DO NOT apply more than 6 lbs of DuPont™ VELISTA™ (equivalent to 3 lbs of active ingredient) per acre per year
- DO NOT apply to any fruit and nut trees that will bear harvestable fruit within 12 months

Table 3b
Landscape Flowering Plants

Landscape Flowering Plants	Target Diseases
Geranium	<i>Botrytis cinera</i>
Daylily	<i>Puccinia hemerocallidis</i>
Gerber Daisy	<i>Erysiphe cichorearum</i>

Specific Application Rates for Landscape Flowering Plants

Mix 6.8 oz of VELISTA™ fungicide in 100 gals of water and apply as a full coverage spray to the point of drip

Apply every 7-14 days during the period of primary infection or when conditions are favorable for disease development

Do not apply more than 8 oz of VELISTA™ per single application in 100 gallons of water per acre

DO NOT apply more than 6 lbs of VELISTA™ (equivalent to 3 lbs of active ingredient) per acre per year

Follow recommendations in the resistance management section for how to use with successive applications (no more than 2 successive applications followed by a different mode of action fungicide)

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal

PESTICIDE STORAGE Store product in original container only in a location inaccessible to children and pets. Do not contaminate water, other pesticides, fertilizer, food or feed in storage.

PESTICIDE DISPOSAL Do not contaminate water, food or feed by storage or disposal. Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional office for guidance.

CONTAINER HANDLING Refer to the Net Contents section of this product's labeling for the applicable "Nonrefillable Container" or "Refillable Container" designation.

Nonrefillable Plastic and Metal Containers (Capacity Equal to or Less Than 50 Pounds) Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then, for Plastic Containers, offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration. Do not burn, unless allowed by state and local ordinances. For Metal Containers, offer for recycling if available or reconditioning if appropriate, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

Nonrefillable Plastic and Metal Containers (Capacity Greater Than 50 Pounds) Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then, for Plastic Containers, offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration. Do not burn, unless allowed by state and local ordinances. For Metal Containers, offer for recycling if available or reconditioning if appropriate, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

Nonrefillable Plastic and Metal Containers, e.g., Intermediate Bulk Containers (IBC) (Size or Shape Too Large to be Tipped, Rolled or Turned Upside Down) Nonrefillable container Do not reuse or refill this container Clean container promptly after emptying the contents from this container into application equipment or mix tank and before final disposal using the following pressure rinsing procedure Insert a lance fitted with a suitable tank cleaning nozzle into the container and ensure that the water spray thoroughly covers the top bottom and all sides inside the container The nozzle manufacturer generally provides instructions for the appropriate spray pressure spray duration and/or spray volume If the manufacturer's instructions are not available pressure rinse the container for at least 60 seconds using a minimum pressure of 30 PSI with a minimum rinse volume of 10% of the container volume Drain pour or pump rinsate into application equipment or rinsate collection system Repeat this pressure rinsing procedure two more times Then for Plastic Containers offer for recycling if available or puncture and dispose of in a sanitary landfill or by incineration For Metal Containers offer for recycling if available or reconditioning if appropriate or puncture and dispose of in a sanitary landfill or by other procedures approved by state and local authorities

Nonrefillable Paper or Plastic Bags, Fiber Sacks including Flexible Intermediate Bulk Containers (FIBC) or Fiber Drums With Liners Nonrefillable container Do not reuse or refill this container Completely empty paper or plastic bag fiber sack or drum liner by shaking and tapping sides and bottom to loosen clinging particles Empty residue into application or manufacturing equipment Then offer for recycling if available or dispose of empty paper or plastic bag fiber sack or fiber drum and liner in a sanitary landfill or by incineration Do not burn unless allowed by state and local ordinances

Refillable Fiber Drums With Liners Refillable container (fiber drum only) Refilling Fiber Drum Refill this fiber drum with DuPont™ VELISTA™ fungicide containing penthiopyrad only Do not reuse this fiber drum for any other purpose Cleaning before refilling is the responsibility of the refiller Completely empty liner by shaking and tapping sides and bottom to loosen clinging particles Empty residue into application or manufacturing equipment Disposing of Fiber Drum and/or Liner Do not reuse this fiber drum for any other purpose other than refilling (see preceding) Cleaning the container (liner and/or fiber drum) before final disposal is the responsibility of the person disposing of the container Offer the liner for recycling if available or dispose of liner in a sanitary landfill or by incineration Do not burn unless allowed by state and local ordinances If drum is contaminated and cannot be reused dispose of it in the manner required for its liner To clean the fiber drum before final disposal completely empty the fiber drum by shaking and tapping sides and bottom to loosen clinging particles Empty residue into application or manufacturing equipment Then offer the fiber drum for recycling if available or dispose of in a sanitary landfill or by incineration Do not burn unless allowed by state and local ordinances

All Other Refillable Containers Refillable container Refilling Container Refill this container with VELISTA™ fungicide containing penthiopyrad only Do not reuse this container for any other purpose Cleaning before refilling is the responsibility of the refiller Prior to refilling inspect carefully for damage such as cracks punctures abrasions worn out threads and closure devices If damage is found do not use the container contact DuPont at the number below for instructions Check for leaks after refilling and before transporting If leaks are found do not reuse or transport container contact DuPont at the number below for instructions Disposing of Container Do not reuse this container for any other purpose other than refilling (see preceding) Cleaning the container before final disposal is the responsibility of the person disposing of the container To clean the container before final disposal use the following pressure rinsing procedure Insert a lance fitted with a suitable tank cleaning nozzle into the container and ensure that the water spray thoroughly covers the top bottom and all sides inside the container The nozzle manufacturer generally provides instructions for the appropriate spray pressure spray duration and/or spray volume If the manufacturer's instructions are not available pressure rinse the container for at least 60 seconds using a minimum pressure of 30 PSI with a minimum rinse volume of 10% of the container volume Drain pour or pump rinsate into application equipment or rinsate collection system Repeat this pressure rinsing procedure two more times Then for Plastic Containers offer for recycling if available or puncture and dispose of in a sanitary landfill or by incineration Do not burn unless allowed by state and local ordinances For Metal Containers offer for recycling if available or reconditioning if appropriate or puncture and dispose of in a sanitary landfill or by other procedures approved by state and local authorities

Do not transport if this container is damaged or leaking If the container is damaged leaking or obsolete or in the event of a major spill fire or other emergency contact DuPont at 1 800 441 3637 day or night

NOTICE TO BUYER Purchase of this material does not confer any rights under patents of countries outside of the United States

The DuPont Oval Logo DuPont™ and VELISTA™ are trademarks or registered trademarks of E I du Pont de Nemours and Company

D - 1486MSTR 022912

LIMITATION OF WARRANTY AND LIABILITY

NOTICE Read This Limitation of Warranty and Liability Before Buying or Using This Product If the Terms Are Not Acceptable Return the Product at Once Unopened and the Purchase Price Will Be Refunded

It is impossible to eliminate all risks associated with the use of this product Such risks may arise from factors such as weather conditions soil factors off target movement unconventional technique presence of other materials the manner of use or application or other unknown factors all of which are beyond the control of DuPont or Seller These risks can cause ineffectiveness of the product or other unintended consequences

DuPont does not agree to be an insurer of these risks

WHEN YOU BUY OR USE THIS PRODUCT YOU AGREE TO ACCEPT THESE RISKS

DuPont warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for the purpose stated in the Directions for Use subject to the inherent risks described above when used in accordance with the Directions for Use under normal conditions

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW DUPONT MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR OF MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW IN NO EVENT SHALL DUPONT OR SELLER BE LIABLE FOR ANY INCIDENTAL CONSEQUENTIAL OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT TO THE EXTENT CONSISTENT WITH APPLICABLE LAW THE EXCLUSIVE REMEDY OF THE USER OR BUYER AND THE EXCLUSIVE LIABILITY OF DUPONT OR SELLER FOR ANY AND ALL CLAIMS LOSSES INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY OR CONTRACT NEGLIGENCE TORT STRICT LIABILITY OR OTHER LEGAL THEORY) RESULTING FROM THE STORAGE USE OR HANDLING OF THIS PRODUCT SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR AT THE ELECTION OF DUPONT OR SELLER THE REPLACEMENT OF THE PRODUCT

To the extent consistent with applicable law that allows such requirement DuPont or its distributor must have prompt notice of any claim so that an immediate investigation of buyer s or user s claim can be made Buyer and all users shall promptly notify DuPont or a DuPont distributor of any claims whether based on contract negligence strict liability other tort or otherwise or be barred from any remedy

This Limitation of Warranty and Liability may not be amended by any oral or written agreement

For product information call 1 888 6 DUPONT (1 888 638 7668)

© 2012 E I DuPont de Nemours and Company, 1007 Market Street, Wilmington, Delaware 19898

All rights reserved