



352-838

08/30/2011

1/9

 <p align="center">U.S. ENVIRONMENTAL PROTECTION AGENCY Office of Pesticide Programs Registration Division (7505C) Ariel Rios Building 1200 Pennsylvania Ave., NW Washington, D.C. 20460</p> <p align="center">NOTICE OF PESTICIDE: <input checked="" type="checkbox"/> Registration <input type="checkbox"/> Reregistration (under FIFRA, as amended)</p>	EPA Reg. Number: 352-838	Date of Issuance: AUG 30 2011
	Term of Issuance: Conditional	
	Name of Pesticide Product: DuPont Dermacor X-151	
Name and Address of Registrant (include ZIP Code): E. I. DuPont de Nemours & Company Crop Protection P. O. Box 30 Newark, DE 19714-0030		
<p>Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.</p> <p>On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act. Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.</p> <p>This product is conditionally registered in accordance with FIFRA sec. 3(c)(7)(A) provided that you:</p> <ol style="list-style-type: none"> 1. Submit and/or cite all data required for registration/reregistration/registration review of your product when the Agency requires all registrants of similar products to submit such data. 2. Make the following label changes before you release the product for shipment: Add the phrase "EPA Registration Number 352-838". 3. Submit the following data: One year storage stability (830.6317) and corrosion characteristics (830.6320) studies. These studies can be run concurrently. It is recommended that the observation must be made at 0, 3, 6, 9, and 12 months period and the results must be submitted to the Agency on completion. These studies must be submitted within 18 months from today's date. 		
Signature of Approving Official:  Driss Benmhend, Acting Product Manager 1	Date: AUG 30 2011	


page 2
EPA Reg. No. 352-838

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(a). Your release for shipment of the product constitutes acceptance of these conditions.

Please be advised that 40 CFR 156.140 requires that a batch code, lot number, or other code identifying the batch of the pesticide distributed and sold be placed on nonrefillable containers. The code may appear either on the label or durably marked on the container itself.

A stamped copy of the label is enclosed for your records. Copies of the product chemistry review, acute toxicity review, and human health science assessment for the new seed treatment uses are also enclosed for your perusal. The fate and ecotoxicity assessment will be sent to you in an electronic file. If you have any questions, please contact Rita Kumar at (703) 308-8291, or kumar.rita@epa.gov.

Sincerely,



Driss Benmhend
Acting Product Manager 1
Insecticide Rodenticide Branch
Registration Division (7505P)

Enclosures (4)



DuPont™ Dermacor® X-151

lettuce and spinach seed treatment

ACCEPTED
With COMMENTS
In EPA Letter Dated:
AUG 30 2011

Under the Federal Insecticide, Fungicide
and Rodenticide Act, As amended, for the
pesticide Registered under EPA Reg. No:

352-838

DRAFT LABEL

4/9



DuPont™ Dermacor® X-151 lettuce and spinach seed treatment

GROUP	28	INSECTICIDE
-------	----	-------------

DERMACOR® X-151 is a flowable suspension.
Contains 5.21 lb. active ingredient per gallon.

<i>Active Ingredient</i>	<i>By Weight</i>
Chlorantraniliprole 3-Bromo-N-[4-chloro-2-methyl-6- [(methylamino)carbonyl]phenyl]-1- (3-chloro-2-pyridinyl)-1H-pyrazole- 5-carboxamide	50%
<i>Other Ingredients</i>	50%
TOTAL	100%

EPA Reg. No. 352-XXX
Nonrefillable Container

Net: _____
OR

Refillable Container
Net: _____

E. I. du Pont de Nemours and Company
1007 Market Street
Wilmington, Delaware 19898
Phone: 1-800-441-7515 (Toll Free)

PRECAUTIONARY STATEMENTS

KEEP OUT OF REACH OF CHILDREN

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

FIRST AID

For questions regarding emergency medical treatment, you may contact 1-800-441-3637 for information.

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

When used as directed this product does not present a hazard to humans or domestic animals.

PERSONAL PROTECTIVE EQUIPMENT

Applicators and other handlers must wear:
Long-sleeved shirt and long pants.
Shoes plus socks.

After the product has been diluted in accordance with label directions for use, shirt, pants, socks, and shoes are sufficient Personal Protective Equipment. Follow manufacturer's instructions for cleaning/maintaining personal protective equipment (PPE). If no such instructions for washables are available, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

USERS SHOULD: Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to aquatic invertebrates, oysters, and shrimp. Do not apply directly to water. Drift and runoff may be hazardous to aquatic organisms in water adjacent to use sites.

Surface Water Advisory-

This product may impact surface water quality due to runoff of rain water. This is especially true for poorly draining soils and soils with shallow ground water. This product is classified as having high potential for reaching surface water via runoff for several months or more after application. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of chlorantraniliprole from runoff water and sediment. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 48 hours.

Ground Water Advisory-

This chemical has properties and characteristics associated with chemicals detected in ground water. This chemical may leach into ground water if used in areas where soils are permeable, particularly where the water table is shallow.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

- Use this product only as a commercial seed treatment in lettuce or spinach.
- Treated seed must not be used for or mixed with food or animal feed, or processed for oil.

AGRICULTURAL USE REQUIREMENTS

DuPont™ DERMACOR® X-151 must be used only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on the label about personal protective equipment, restricted-entry interval, and notification to workers (as applicable).

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

For any requirements specific to your State or Tribe, consult the State or Tribal agency responsible for pesticide regulation.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 4 hours.

Exception: If the seed is treated with the product and the treated seed is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

For early entry into treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, wear:

- Long-sleeved shirt and long pants
- Shoes plus socks

When applied according to this label, DERMACOR® X-151 will provide early season control of beet armyworm, corn earworm, cabbage looper and leafminer larvae (*Liriomyza* species) in lettuce and spinach. DERMACOR® X-151 is recommended for control of *Liriomyza* species except suppression only for *L. huidabrensis* and *L. langei*. Consult your local DuPont Crop Protection company representative or agricultural advisor for guidance on predicted pest pressure based on historical field data for these pests in your area. Do not use more than 0.1 lb ai per acre or less than 0.05 lb ai per acre.

DERMACOR® X-151 seed treatment should be used only in accordance with directions on this label or in separate DuPont supplemental labeling.

DuPont will not be responsible for losses or damages resulting from the use of this product in any manner not specifically stated on this label or other labels or bulletins published by DuPont. User assumes all risks associated with such non-recommended use.

DERMACOR® X-151 may be applied to lettuce and spinach seed. Lettuce and spinach seed is typically planted at a soil depth of 1/8th to 1/4th inch.

Treated seed exposed on soil surface may be hazardous to aquatic invertebrates. Cover or collect seeds spilled during loading.

DERMACOR® X-151 treated lettuce or spinach may be replanted if an emergency replanting due to crop failure occurs.

Use of DERMACOR® X-151 as a seed treatment must be communicated to all personnel involved in seasonal insect control recommendations.

Following the planting of DERMACOR® X-151 treated seed, total foliar applications for a crop must not exceed a total rate of 0.1 lb chlorantraniliprole per acre. Soil applications of chlorantraniliprole containing products are not permitted.

INTEGRATED PEST MANAGEMENT

DuPont supports the use of Integrated Pest Management (IPM) programs to control pests. This product may be used as part of an IPM program, which can include biological, cultural, and genetic practices, aimed at preventing economic pest damage. IPM principles and practices include field scouting or other detection methods, correct target pest identification, population monitoring, rotation of insecticides with different modes-of-action, and treating when target pest populations reach locally determined action thresholds. Consult your state cooperative extension service, professional consultants or other qualified authorities to determine appropriate action treatment threshold levels for treating specific pest/crop or site systems in your area.

RESISTANCE MANAGEMENT

DERMACOR® X-151 is a Group 28 Insecticide. Repeated and exclusive use of DERMACOR® X-151 or other Group 28 Insecticides may lead to the buildup of resistant strains of insects in some crops.

Because the development of resistance cannot be predicted, this product may be used as part of resistant management strategies established for the use area. These strategies may include incorporation of cultural and biological control practices, alternation of mode-of-action classes of insecticides (non-Group 28) on succeeding generations and targeting the most susceptible life stage. Consult your local or state agricultural authorities for details.

For crops grown from DERMACOR® X-151 treated seeds:

- Including the DERMACOR® X-151 seed treatment, make no more than two applications of Chlorantraniliprole or other Group 28 products per generation to the same insect species on a crop or within a 30 day period. Application(s) to the next generation of target pest(s) must be with an effective product with a different mode of action (non-Group 28 insecticide).
- For short season crops (planting to harvest of approximately 30 - 45 days), applications to the next generation of target pest(s) on the same farm acre must be made with an effective product with a different mode of action (non-Group 28 insecticide).

If you experience difficulty with control and resistance is a reasonable cause, immediately consult your local DuPont

Crop Protection company representative or agricultural advisor for the best alternate method of control for your area. For additional information on insect resistance monitoring, visit the Insecticide Resistance Action Committee (IRAC) on the web at <http://www.irac-online.org>.

CROP ROTATION

Crops on this label and the following crops or crop groups may be planted immediately following harvest: Artichoke; Asparagus; Brassica (Cole) Leafy Vegetables (Crop Group 5); Corn (field, pop, seed, and sweet); Cotton; Cucurbit Vegetables (Crop Group 9); Fruiting Vegetables (Crop Group 8); Grape; Grass Forage, Fodder and Hay (Crop Group 17); Herbs subgroup (Crop Group subgroup 19A); Leafy Vegetables (non-brassica, Crop Group 4); Legume Vegetables (Crop Group 6); Foliage of Legume Vegetables (Crop Group 7); Spearmint; Peppermint; Nongrass Animal Feeds (Forage, Fodder, Straw, and Hay Crop Group 18); Okra; Potato; Prickly Pear Cactus; Root and Tuber Vegetables (Crop Group 1); Strawberries; Sugarcane; Tree Nuts (Crop Group 14); Tobacco and Tops of Root and Tuber Vegetables (Crop Group 2).

The following crops or crop groups may be planted 30 days following the planting of DuPont™ DERMACOR® X-151 treated seed: garlic, greathheaded garlic, dry bulb onion, leek, green onion, Welsh onion, shallot; Cereal Grains (Crop Group 15); soybean and peanut.

All other crops cannot be planted until 12 months after the last planting of DERMACOR® X-151 treated seed.

APPLICATION AND MIXING GUIDANCE

DERMACOR® X-151 may be applied as a water-based slurry with other registered seed treatment fungicides and insecticides. Potential mixing partners should be tested for physical compatibility with DERMACOR® X-151 and added dyes or polymers before mixing in a large tank for seed treatment. Avoid mixtures of several materials and very concentrated spray mixtures. Observe the most restrictive of the labeling limitations and precautions of all products used in mixtures. Do not mix this product with any product that prohibits such mixing.

USE RATE

Use DERMACOR® X-151 as recommended in the following Use-Rate Table for early-season control of beet armyworm, corn earworm, cabbage looper and leafminer larvae (*Liriomyza* species) in lettuce and spinach. DERMACOR® X-151 is recommended for control of *Liriomyza* species except suppression only for *L. huidabrensis* and *L. langei*. The seed treating rate of DERMACOR® X-151 should be determined based on planned seeding density. Consult your DuPont Crop Protection company representative for further guidance.

DERMACOR® X-151 Seed Treatment Use-Rate Table

Planting Density (seed per acre)	DERMACOR® X-151 Use Rate (fl. oz/lb of seed)	DERMACOR® X-151 Use Rate (lb ai/A)	DERMACOR® X-151 Use Rate (Milligrams ai/Seed)
75,000-250,000	2.3 - 15.2	0.05 - 0.10	0.091 - 0.605
2,000,000-3,500,000	0.3 - 0.5* 0.032 - 0.056**	0.10	0.013 - 0.023

Do not use more than 0.1 lb ai per acre or less than 0.05 lb ai per acre.

*Spring Mix lettuce

**Baby spinach

MIXING GUIDANCE

For use in commercial seed treaters only, operating standard, calibrated seed treatment equipment. Treatment equipment must be clean and free of previous pesticide deposits before applying DERMACOR® X-151. Add DERMACOR® X-151 directly to mixing tank. Mix thoroughly recommended amount of DERMACOR® X-151 with the required amount of water.

Add an EPA approved dye or colorant to treat the seeds per 40 CFR 153.155(b)(1) during the seed treatment process. Ensure that all treated seed are dyed an unnatural color.

MIXING TANK CLEANOUT

Prior to application, start with clean, well maintained application equipment. Immediately following application, thoroughly clean all mixing equipment to reduce the risk of forming hardened deposits, which might become difficult to remove.

Drain mixing equipment. Thoroughly rinse mixing equipment and flush with clean water.

Clean all other associated application equipment. Take all necessary safety precautions when cleaning equipment. Do not clean near wells, water sources or desirable vegetation. Dispose of waste rinse water in accordance with local regulations.

COMMERCIAL SEED BAG LABELING

In accordance with the Federal Seed Act and the US Environmental Protection Agency requirements, seeds commercially treated with a pesticide must be labeled as follows:

- Treated Seed. Do not use for food, feed or oil purposes.
- This seed has been treated with DuPont™ DERMACOR® X-151 seed treatment (Chlorantraniliprole).
- Treated seed exposed on soil surface may be hazardous to aquatic invertebrates. Cover or collect seeds spilled during loading.
- Store away from food or feed.
- Dispose of all excess treated seed by burying seed away from bodies of water.
- Lettuce and spinach seed is typically planted at a soil depth of 1/8th to 1/4th inch.
- DERMACOR® X-151 treated lettuce or spinach may be replanted if an emergency replanting due to crop failure occurs.
- Use of DERMACOR® X-151 as a seed treatment must be communicated to all personnel involved in seasonal insect control recommendations.
- Including the DERMACOR® X-151 seed treatment, make no more than two applications of Chlorantraniliprole or other Group 28 products per generation to the same insect species on a crop or within a 30 day period. Application(s) to the next generation of target pest(s) must be with an effective product with a different mode of action (non-Group 28 insecticide).
- For short season crops (planting to harvest of approximately 30 - 45 days), applications to the next generation of target pest(s) on the same farm acre must be made with an effective product with a different mode of action (non-Group 28 insecticide).
- Following the planting of DERMACOR® X-151 treated seed, total foliar applications for a crop must not exceed a total rate of 0.1 lb chlorantraniliprole per acre. Soil applications of chlorantraniliprole-containing products are not permitted.

Crop Rotation

Crops on this label and the following crops or crop groups may be planted immediately following harvest: Artichoke; Asparagus; Brassica (Cole) Leafy Vegetables (Crop Group 5); Corn (field, pop, seed, and sweet); Cotton; Cucurbit Vegetables (Crop Group 9); Fruiting Vegetables (Crop Group 8); Grape; Grass Forage, Fodder and Hay (Crop Group 17); Herbs subgroup (Crop Group subgroup 19A); Leafy Vegetables (nonbrassica, Crop Group 4); Legume Vegetables (Crop Group 6); Foliage of Legume Vegetables (Crop Group 7); Spearmint;Peppermint; Nongrass Animal Feeds (Forage, Fodder, Straw, and Hay Crop Group 18); Okra; Potato; Prickly Pear Cactus; Root and Tuber Vegetables (Crop Group 1); Strawberries; Sugarcane; Tobacco and Tops of Root and Tuber Vegetables (Crop Group 2). The following crops or crop groups may be planted 30 days following the last planting of DERMACOR® X-151 treated seed: garlic; greatheaded garlic; dry bulb onion; leek; green onion; Welsh onion; shallot; Cereal Grains (Crop Group 15); soybean and peanut. All other crops cannot be planted until 12 months after the last planting of DERMACOR® X-151 treated seed.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Do not subject to temperatures below 32 degrees F. Store product in original container only in a location inaccessible to children and pets. Do not contaminate water, other pesticides, fertilizer, food or feed in storage. Not for use or storage in or around the home.

PESTICIDE DISPOSAL: Do not contaminate water, food or feed by storage or disposal. Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

CONTAINER HANDLING:

Refer to the Net Contents section of this product's labeling for the applicable "Refillable Container" or "Nonrefillable Container" designation.

Nonrefillable Rigid Plastic and Metal Containers (Capacity Equal to or Less Than 5 Gallons):

Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then, for Plastic Containers, offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration. Do not burn, unless allowed by state and local ordinances. For Metal Containers, offer for recycling if available or reconditioning if appropriate or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

Nonrefillable Plastic and Metal Containers

(Capacity Greater Than 5 Gallons): Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then, for Plastic Containers, offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration. Do not burn, unless allowed by state and local ordinances. For Metal Containers, offer for recycling if available or reconditioning if appropriate or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

Nonrefillable Plastic and Metal Containers, e.g., Intermediate Bulk Containers [IBC] (Size or Shape Too Large to be Tipped, Rolled or Turned Upside Down): Nonrefillable container. Do not reuse or refill this container. Clean container promptly after emptying the contents from this container into application equipment or mix tank and before final disposal using the following pressure rinsing procedure. Insert a lance fitted with a suitable tank cleaning nozzle into the container and ensure that the water spray thoroughly covers the top, bottom and all sides inside the container. The nozzle manufacturer generally provides instructions for the appropriate spray pressure, spray duration and/or spray volume. If the manufacturer's instructions are not available, pressure rinse the container for at least 60 seconds using a minimum pressure of 30 PSI with a minimum rinse volume of 10% of the container volume. Drain, pour or pump rinsate into application equipment or rinsate collection system. Repeat this pressure rinsing procedure two more times. Then, for Plastic Containers, offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration. For Metal Containers, offer for recycling if available or reconditioning if appropriate, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

All Refillable Containers: Refillable container.
Refilling Container: Refill this container with DuPont™ DERMACOR® X-151 containing chlorantraniliprole only. Do not reuse this container for any other purpose. Cleaning before refilling is the responsibility of the refiller. Prior to refilling, inspect carefully for damage such as cracks, punctures, abrasions, worn out threads and closure devices. If leaks are found, do not reuse or transport container, contact DuPont at the number below for instructions.
Disposing of Container: Do not reuse this container for any other purpose other than refilling (see preceding). Cleaning the container before final disposal is the responsibility of the person disposing of the container. To clean the container before final disposal, use the following pressure rinsing procedure. Insert a lance fitted with a suitable tank cleaning nozzle into the container and ensure that the water spray thoroughly covers the top, bottom and all sides inside the container. The nozzle manufacturer generally provides instructions for the appropriate spray pressure, spray duration and/or spray volume. If the manufacturer's instructions are not available, pressure rinse the container for at least 60 seconds using a minimum pressure of 30 PSI with a minimum rinse volume of 10% of the container volume. Drain, pour or pump rinsate into application equipment or rinsate collection system. Repeat this pressure rinsing procedure two more times. Then, for Plastic Containers, offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration. Do not burn, unless allowed by state and local ordinances. For Metal Containers, offer for recycling if available or reconditioning if appropriate, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

Do not transport if container is damaged or leaking. If the container is damaged, leaking or obsolete, or in the event of a major spill, fire or other emergency, contact DuPont at 1-800-441-3637, day or night.

NOTICE TO BUYER—Purchase of this material does not confer any rights under patents of countries outside of the United States.

The DuPont Oval Logo, DuPont™ and DERMACOR® are trademarks or registered trademarks of E. I. du Pont de Nemours and Company.

D - 1533 051011

LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read this Limitation of Warranty and Liability Before Buying or Using This Product. If the Terms Are Not Acceptable, Return the Product at Once, Unopened, and the Purchase Price Will Be Refunded.

It is impossible to eliminate all risks associated with the use of this product. Such risks arise from weather conditions, soil factors, off target movement, unconventional farming techniques, presence of other materials, the manner of use or application, or other unknown factors, all of which are beyond the control of DuPont. These risks can cause: ineffectiveness of the product, crop injury, or injury to non-target crops or plants. **WHEN YOU BUY OR USE THIS PRODUCT, YOU AGREE TO ACCEPT THESE RISKS.**

DuPont warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for the purpose stated in the Directions for Use, subject to the inherent risks described above, when used in accordance with the Directions for Use under normal conditions.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, DUPONT MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR OF MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL DUPONT OR SELLER BE LIABLE FOR ANY INCIDENTAL, CONSEQUENTIAL OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. BUYER'S OR USER'S BARGAINED-FOR EXPECTATION IS CROP PROTECTION. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER AND THE EXCLUSIVE LIABILITY OF DUPONT OR SELLER, FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY OR CONTRACT, NEGLIGENCE, TORT OR STRICT LIABILITY), WHETHER FROM FAILURE TO PERFORM OR INJURY TO CROPS OR OTHER PLANTS, AND RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT, OR AT THE ELECTION OF DUPONT OR SELLER, THE REPLACEMENT OF THE PRODUCT.

To the extent consistent with applicable law that allows such requirement, DuPont or its Ag Retailer must have prompt notice of any claim so that an immediate inspection of buyer's or user's growing crops can be made. Buyer and all users shall promptly notify DuPont or a DuPont Ag Retailer of any claims, whether based on contract, negligence, strict liability, other tort or otherwise, or be barred from any remedy.

This Limitation of Warranty and Liability may not be amended by any oral or written agreement.

For product information call: 1-888-6-DUPONT [1-888-638-7668]

© 2011 E. I. du Pont de Nemours and Company, Crop Protection, 1007 Market Street, Wilmington, Delaware 19898.

All rights reserved.