

352-840

11/30/2012

1/13



U.S. ENVIRONMENTAL PROTECTION AGENCY

Office of Pesticide Programs
Registration Division (7505P)
Ariel Rios Building
1200 Pennsylvania Ave., NW
Washington, D.C. 20460

EPA Reg. Number: 352-840

Date of Issuance: NOV 30 2012

NOTICE OF PESTICIDE:

[x] Registration
[] Reregistration
(under FIFRA, as amended)

Term of Issuance: Unconditional

Name of Pesticide Product: Dupont™ Aproach™

Jack Cain
E.I. du Pont de Nemours and Company
1007 Market Street
Wilmington, DE 19898

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act. Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is unconditionally registered in accordance with FIFRA sec. 3(c)(5) provided that you:

- 1. Revise the EPA Reg. No. to 352-840
2. Assure that the establishment number and net content are added to the label.
3. Submit and/or cite all data required for registration/reregistration review of your product when the Agency requires all registrants of similar products to submit data.

A stamped copy of the label is enclosed for your records. Submit one (1) copy of the final printed label before you release the product for shipment.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

If you have any questions regarding this notice, please contact Grant Rowland at (703) 347-0254 or at rowland.grant@epa.gov.

Kathryn V. Montague
Product Manager 23
Herbicide Branch
Registration Division (7505P)
Office of Pesticide Programs

[Handwritten signature of Kathryn V. Montague]

Date: NOV 30 2012



DuPont™ Aproach™

FUNGICIDE

GROUP 11 FUNGICIDE

Suspension Concentrate

Active Ingredients

By Weight

Picoxystrobin

Methyl (αE)-α-(methoxymethylene)-2-[[[6-(trifluoromethyl)-2-pyridinyl]oxy]methyl]benzeneacetate22.5%

Other Ingredients77.5%

TOTAL100.0%

Contains 2.08 pounds of picoxystrobin per gallon of product

EPA Reg. No. 352-~~HN~~ 890

EPA Est. No. _____


Nonrefillable Container

Net: _____

OR

Refillable Container

Net: _____



KEEP OUT OF REACH OF CHILDREN CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand this label, find someone to explain it to you in detail.)

FIRST AID

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice.

IF SWALLOWED: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to by a poison control center or doctor. Do not give anything to an unconscious person.

IF ON SKIN: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For medical emergencies involving this product, call toll-free 1-800-441-3637.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION! Causes moderate eye irritation. Harmful if swallowed. Harmful if absorbed through skin. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Avoid contact with eyes, skin or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Mixers, loaders, applicators, and other handlers must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks
- Chemical resistant gloves made of any waterproof material such as polyethylene or polyvinyl chloride

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry. Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them.

ENGINEERING CONTROL STATEMENTS

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

USERS SHOULD: Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and put on clean clothing.

ENVIRONMENTAL HAZARDS

This product is toxic to fish and aquatic invertebrates, including shrimp and oysters. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash water or rinsate.

SURFACE WATER ADVISORY: Picoxystrobin has the potential to contaminate surface water through spray drift. Under some conditions, picoxystrobin may also have a high potential for runoff into surface water, especially in areas with poorly-draining soils, and areas with shallow water tables. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water will reduce the potential for runoff. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 48 hours.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

DuPont™ APPROACH™ must be used only in accordance with instructions on this label or separately published DuPont labeling.

DuPont will not be responsible for losses or damages resulting from use of this product in any manner not specifically instructed by DuPont. User assumes all risks associated with such non-labeled use.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the State or Tribal agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on the label about personal protective equipment (PPE), and restricted-entry interval, and notification to workers (as applicable). The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours. PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Shoes and socks
- Chemical resistant gloves made of any waterproof material (such as Natural Rubber, selection Category A)

4/10

PRODUCT INFORMATION

DuPont™ APROACH™ is a broad-spectrum fungicide for control of foliar and soil-borne plant diseases and has preventive, curative, and systemic activity. APROACH™ must be applied in a regularly scheduled protective spray program in rotation with other fungicides. When used in a disease control program, APROACH™ improves plant health, vigor, and yield. See directions below for specific crop/disease instructions.

APROACH™ rapidly penetrates into plant tissues and is rainfast within 1-hour after application.

This product may be applied to crop sites that contain areas of temporary surface water caused by collection of water between planting beds, in equipment ruts, or in other depressions caused by management activities.

INTEGRATED PEST MANAGEMENT

This product may be used as part of an Integrated Pest Management (IPM) program that can include biological, cultural, and genetic practices aimed at preventing economic pest damage. IPM principles and practices include field scouting or other detection methods, correct target pest identification, population monitoring, and treating when disease forecasting models reach locally determined action thresholds. Consult your state cooperative extension service, professional consultants, or other qualified authorities to determine appropriate action treatment threshold levels for treating specific pest/crop or site systems in your area.

RESISTANCE

DuPont™ APROACH™, which contains the active ingredient picoxystrobin, is a Group 11 fungicide based on the mode of action classification system of the Fungicide Resistance Action Committee of Croplife International.

Repeated use of products for control of specific plant pathogens may lead to selection of resistant strains of fungi and result in a reduction of disease control. A disease management program that includes rotation and/or tank mixing with fungicides that have a different mode of action is essential to reduce the risk of fungicide resistance development. When making applications to pathogens that have tolerance or resistance to APROACH™ and other strobilurin fungicides, tank mix with a fungicide with a different mode-of-action that is effective for controlling the target disease. For guidance on a particular crop and disease control situation, consult your agricultural dealer, consultant, applicator or appropriate state agricultural extension service representative for specific area practices and/or requirements.

APPLICATION INFORMATION

APPLICATION EQUIPMENT

APROACH™ may be applied with ground, air or chemigation equipment.

APPLICATION VOLUME

Use a sufficient volume of water to ensure thorough coverage when applying APROACH™ as a broadcast spray. Select a spray volume and delivery system that will ensure thorough coverage and a uniform spray pattern. An increased volume of water may be required as foliage density increases.

TANK MIXTURES

Do not use an adjuvant or crop oil when applying APROACH™ on corn between the V8 and VT stages of growth. Do not use an adjuvant or crop oil when applying APROACH™ on dry beans or peas.

The crop safety of all tank mixtures with APROACH™ which may include physically compatible pesticides, fertilizers, adjuvants, and/or additives, has not been tested. When considering the use of a tank mixture on a labeled crop without prior experience, or which is not specifically described on APROACH™ product labeling or in other DuPont product use instruction, it is important to first understand crop safety. To test for crop safety prepare a small volume of the intended tank mixture, apply it to an area of the target crop as directed by both this and the tank mix partner product labels, and observe the treated crop to ensure that a phytotoxic response does not occur. DuPont will not be responsible for any crop injury arising from the use of a tank mixture that is not specifically described on APROACH™ product labeling or in other DuPont product use instruction.

Always follow the tank mix instructions of the product label that is most restrictive.

Consult a DuPont representative or local agricultural authorities for more information concerning tank mixtures.

Physical Compatibility

APROACH™ is physically compatible with many commonly used fungicides, herbicides, insecticides, biological control products, liquid fertilizers, non-ionic surfactants, crop oils, methylated seed oils and drift control additives. However, since the formulations of products change, it is important to test the physical compatibility of desired tank mixes and check for undesirable physical effects, including settling out or flocculation. To determine physical compatibility, add the proportions of the tank mix products and water to a small container, mix thoroughly and allow to stand for 20 minutes. If the combination remains mixed, or can be re-mixed readily, it may be considered physically compatible.

MIXING INSTRUCTIONS

1. Fill clean spray tank 1/4 - 1/2 full of water.
2. While agitating, add the required amount of APROACH™, continuing agitation until the product is completely dispersed.

3. Continue filling the tank, with agitation, adding desired additives or tank mix partners, following the sequence listed below in 'tank mixing sequence'.

Tank Mixing Sequence

Add different formulation types in the sequence indicated below. Allow time for complete mixing and dispersion after addition of each product.

1. water-soluble bag
2. water-dispersible granules
3. wettable powders
4. water-based suspension concentrates (DuPont™ APROACH™)
5. water-soluble concentrates
6. oil-based suspension concentrates
7. emulsifiable concentrates
8. adjuvants, surfactants, and oils
9. soluble fertilizers
10. drift control additives

CHEMIGATION

Apply APROACH™ only through sprinkler irrigation systems (such as center pivot, lateral move, end tow, side (wheel) roll, traveler, big gun, solid set or hand move irrigation systems).

Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water. If you have questions about calibration, contact your State Extension Service Specialists, equipment manufacturers or other experts.

Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place.

A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, must shut the system down and make necessary adjustments should the need arise.

Specific Instructions for Public Water Systems:

Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.

Chemigation systems connected to public water systems must contain a functional, reduced-pressure zone, backflow preventer (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.

The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.

The pesticide injection pipeline must contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected.

Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

Specific Instructions for Sprinkler Irrigation Systems:

The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.

The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.

The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.

The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.

5/13

Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

Do not apply when wind speed favors drift beyond the area to be treated.

Good agitation is required in the injection tank. In moving systems, apply specified dosage of DuPont™ APROACH™ as a continuous injection. In nonmoving systems inject APROACH™ for 15 to 30 minutes at end of cycle. Use the least amount of water possible consistent with uniform coverage.

Mix the amount of APROACH™ needed for acreage to be treated into the quantity of water determined during prior calibration. For moving systems inject into the system continuously for one complete revolution of the field. For nonmoving systems inject into system for the time established during calibration.

Stop injection equipment after completing treatment; continue to operate irrigation equipment until all APROACH™ is flushed from the system.

7/13

POST-EMERGENCE APPLICATION TIMING AND USE RATES

Table 1: DuPont™ APROACH™ Labeled Crop and Crop Groups, Pre-Harvest Intervals, Maximum Single Application Rates, and Total Crop Use Rates.

Crop, Crop Group or Subgroup (with examples)	Minimum Time from last Application to Harvest (PHI days or crop stage)	Maximum Rate per Acre per Application (fluid ounces product)	Maximum Product per Acre per Crop (fluid ounces product)
Canola	28-days	12	24
Cereal grains (except rice)	Apply no later than the beginning of flowering (Feekes 10.5) for grain 45-days, wheat and barley grain and straw 14-days, wheat and barley hay 7-days, wheat forage	12	36
Sorghum	Do not apply after flowering Do not use for hay or forage	12	36
Corn	7-days, grain and ear, 0-days, forage	12	36
Legume vegetables Dried shelled beans, peas	14-days, seed 0-day, vines and hay	12	24
Soybean (forage, hay and grain)	14-days	12	36 (grain) 12 (forage, hay)

Table 2: APROACH™ Specific Crop/Crop Group Disease Treatment Use Rates, and Treatment Instructions.

Crop/ Crop Group	Disease Controlled or Suppressed	Rate (fluid ounces per acre)	Treatment Instructions
Canola	Alternaria blackspot, leaf and stem spots (<i>Alternaria spp.</i>) Blackleg (<i>Leptosphaeria maculans, L. biglobosa</i>)	6 to 12	Begin applications prior to disease development and continue on a 7- to 14-day interval. Use higher rate and shorter interval when disease pressure is high.
	Sclerotinia stem rot (<i>Sclerotinia spp.</i>)	8 to 12	Sclerotinia stem rot: Begin application at 20-50% bloom prior to disease development and continue on a 7- to 14-day interval. Use higher rate and shorter interval when disease pressure is high.

- Make no more than 2 sequential applications of APROACH™ before switching to a fungicide with a different mode of action.
- The minimum pre-harvest interval (PHI) between the last application and harvest is 28-days.
- Do not exceed 24 fl ounces per acre per crop.

Crop/ Crop Group	Disease Controlled or Suppressed	Rate (fluid ounces per acre)	Treatment Instructions
Cereal grains Sorghum, (milo), sorghum spp. (sudangrass and hybrids)	Anthraxnose (<i>Colletotrichum graminicola</i>) Grey leafspot (<i>Cercospora sorghi</i>) Rust, common (<i>Puccinia sorghi</i>)	6 to 12	Begin applications prior to disease development and continue on a 7- to 14-day interval. Use higher rate and shorter interval when disease pressure is high. Do not apply after flowering.

- Make no more than 2 sequential applications of APROACH™ before switching to a fungicide with a different mode of action.
- Do not apply after flowering.
- Do not use for hay or forage.
- Do not exceed 36 fluid ounces per acre per crop.

Crop/ Crop Group	Disease Controlled or Suppressed	Rate (fluid ounces per acre)	Treatment Instructions
Cereal grains Barley, wheat, rye, oats, triticale	Black point (<i>Alternaria spp.</i> , <i>Helminthosporium spp.</i>) Leaf and glume blotch (<i>Stagonospora spp.</i> , <i>Septoria spp.</i>) Powdery mildew (<i>Erysiphe graminis f. sp. tritici</i>) Rusts (<i>Puccinia spp.</i>) Spot blotch (<i>Cochliobolus sativus</i>) Tan spot (<i>Pyrenophora tritici-repentis</i>)	3 to 4	Make a single application between tillering through jointing for early season disease control/suppression. Starting no sooner than 7-days later, additional 6 fl oz to 12 fl oz treatments can be made depending on disease pressure and environmental conditions.
		6 to 12	Begin applications prior to disease development and continue on a 7- to 14-day interval, depending on the targeted disease. Use higher rate and shorter interval when disease pressure is high.
	Disease suppression Scab (<i>Fusarium spp.</i>)	6 to 12	To optimize yields in cereals, it is important to protect the flag leaf from foliar diseases. For optimizing yield and flag leaf disease control, apply DuPont™ APROACH™ at Feekes' 9, 'flag leaf out'. Apply no later than the beginning of flowering (Feekes 10.5).

- Make no more than 2 sequential applications of APROACH™ before switching to a fungicide with a different mode of action.
- For grain, apply no later than the beginning of flowering (Feekes 10.5).
- The minimum pre-harvest interval (PHI) between the last application and harvest for grain is 45 days, 7-days for forage and 14-days for hay.
- Do not exceed 36 fluid ounces per acre per crop.

Crop/ Crop Group	Disease Controlled or Suppressed	Rate (fluid ounces per acre)	Treatment Instructions
Corn, field corn, sweet corn, seed popcorn	Anthracnose leaf blight and stalk rot (<i>Colletotrichum graminicola</i>) Eye spot (<i>Aureobasidium zeae</i> , <i>Kabatiella zeae</i>) Gray leaf spot (<i>Cercospora zeae-maydis</i>) Leaf spots (<i>Alternaria spp.</i>) Northern corn leaf blight (<i>Septosphaeria turcica</i> , <i>Exserohium turcicum</i>) Northern corn leaf spot (<i>Cochliobolus carbonum</i>) Physoderma brown spot (<i>Physoderma maydis</i>) Rust, common (<i>Puccinia sorghi</i>) Rust, southern (<i>Puccinia polyspora</i>) Southern corn leaf blight (<i>Cochliobolus heterostrophus</i> , <i>Bipolaris maydis</i>) Yellow leaf blight (<i>Phyllosticta maydis</i>)	3 to 4	Make a single 3-4 oz application between V4 to V7 for early season disease control/suppression.
		6 to 12	Make additional 6 to 12 fl oz applications at 7 to 14-day intervals. For best results apply between VT to R3 and make applications prior to disease development. Use the higher rate and shorter interval when disease pressure is high.

- Make no more than 2 sequential applications of APROACH™ before switching to a fungicide with a different mode of action.
- Do not tank mix APROACH™ with an adjuvant or crop oil when spraying corn between the V8 and VT stages of growth.
- The minimum pre-harvest interval (PHI) between the last application and harvest for grain or ear is 7-days and for forage is 0-days.
- Do not exceed 36 fluid ounces per acre per crop.

Crop/ Crop Group	Disease Controlled or Suppressed	Rate (fluid ounces per acre)	Treatment Instructions
Legume vegetables (dry shelled beans and peas) Broad bean (dry); chickpea (garbanzo); guar; lablab bean; lentil; pigeon pea; (Lupinus) grain lupin, sweet lupin, white lupin, white sweet lupin; (Phaseolus) field bean, kidney bean, lima bean, navy bean, pinto bean, teparty bean, (Vigna) adzuki bean, blackeyed pea, catjang, cowpea, crowder pea, moth bean, mung bean, rice bean, southern pea, urd bean, (Pisum) field pea	Alternaria blight, leafspot (<i>Alternaria spp.</i>) Anthracnose (<i>Colletotrichum spp.</i>) Ascochyta blight, leafspot (<i>Ascochyta spp.</i>) Cercospora leafspot (<i>Cercospora spp.</i>) Downy mildew (<i>Phytophthora nicotianae</i>) Mycosphaerella blight (<i>Mycosphaerella spp.</i>) Powdery mildew (<i>Erysiphe spp.</i>) Rust (<i>Uromyces spp. Phakopsora spp.</i>) Septoria blotch (<i>Septoria spp.</i>)	6 to 12	Begin applications prior to disease development and continue on a 7- to 14-day interval. Use higher rate and shorter interval when disease pressure is high.
	Sclerotinia rot, white mold (<i>Sclerotinia spp.</i>)	8 to 12	For white mold: make initial preventive application at beginning bloom and follow with 2nd application 7-10 days later at full bloom.

- Make no more than 2 sequential applications of DuPont™ APROACH™ before switching to a fungicide with a different mode of action.
- The minimum pre-harvest interval (PHI) between last application and harvest of seed is 14-days and vines and hay is 0-days.
- Do not exceed 24 fluid ounces per acre per crop.
- Do not tank mix APROACH™ with an adjuvant or crop oil when spraying dry beans or peas.

Crop/ Crop Group	Disease Controlled or Suppressed	Rate (fluid ounces per acre)	Treatment Instructions
Soybean	Aerial web blight (<i>Rhizoctonia solani</i>) Anthracnose (<i>Colletotrichum truncatum</i>) Alternaria leaf spot (<i>Alternaria spp.</i>) Brown Spot (<i>Septoria glycines</i>) Cercospora blight and leaf spot, purple seed stain (<i>Cercospora kikuchii</i>) Downy mildew (<i>Peronospora manshurica</i>) Frogeye leafspot (<i>Cercospora soja</i>) Pod and stem blight (<i>Diaporthe phaseolum</i>) Powdery mildew (<i>Erysiphe spp.</i>) Rust (<i>Puccinia spp., Phakospora spp.</i>) Target Spot (<i>Corynespora cassicola</i>)	6 to 12	Begin applications prior to disease development and continue on a 7- to 14-day interval. Use higher rate and shorter interval when disease pressure is high.
	Sclerotinia stem rot (<i>Sclerotinia sclerotiorum</i>)	8 to 12	For white mold: make initial preventive application at 100% bloom (1 flower blooming on all plants) and follow with 2nd application 7-10 days later at full bloom.

- Make no more than 2 sequential applications of APROACH™ before switching to a fungicide with a different mode of action.
- The minimum pre-harvest interval (PHI) between last application and harvest of grain, forage, and hay is 14-days.
- Do not exceed 12 fluid ounces per acre per crop if grown for forage and hay.
- Do not exceed 36 fluid ounces per acre per crop if grown for grain (seed).

16/13

ADDITIONAL INSTRUCTIONS, PRECAUTIONS AND RESTRICTIONS FOR ALL USES

IMPORTANT RESTRICTIONS

- Do not use DuPont™ APROACH™ on residential plantings.

SPRAY DRIFT RESTRICTIONS

- Where states have more stringent regulations they must be observed.

AERIAL APPLICATIONS

- Applicators are required to use upwind swath displacement, and displacement distance must increase with increasing drift potential.
- Applications into temperature inversions are prohibited.
- Spray must be released at the lowest height consistent with pest control objectives and flight safety.

GROUND APPLICATIONS

- Applications into temperature inversions are prohibited.
- Apply spray at the lowest height that is consistent with pest control objectives.

See Spray Drift Management Section of this label for additional information.

IMPORTANT PRECAUTIONS

- Not all crops within a crop group, and not all varieties, cultivars or hybrids of crops, have been individually tested for crop safety. It is not possible to evaluate for crop safety all applications of APROACH™ on all crops within a crop group, on all varieties, cultivars, or hybrids of those crops, or under all environmental conditions and growing circumstances. To test for crop safety, apply the product in accordance with the label instructions to a small area of the target crop to ensure that a phytotoxic response will not occur, especially where the application is a new use of the product by the applicator.

CROP ROTATION

Treated areas may be replanted immediately after harvest with any crop appearing on this label.

All other crops may be planted 6 months following the last application of APROACH™.

EQUIPMENT CLEANING

Prior to application, start with clean, well maintained application equipment. Immediately following application, thoroughly clean all spray equipment to reduce the risk of forming hardened deposits which might become difficult to remove.

Drain spray equipment. Thoroughly rinse sprayer and flush hoses, boom and nozzles with clean water. Clean all other associated application equipment. Take all necessary safety precautions when cleaning equipment. Do not clean near wells, water sources or desirable vegetation.

Dispose of waste rinse water in accordance with local regulations.

SPRAY DRIFT MANAGEMENT

The interaction of many equipment and weather-related factors determines the potential for spray drift. The applicator is responsible for considering all these factors when making application decisions. Avoiding spray drift is the responsibility of the applicator.

IMPORTANCE OF DROPLET SIZE

The most effective drift management strategy is to apply the largest droplets which are consistent with pest control objectives. The presence of sensitive species nearby, the environmental conditions, and pest pressure may affect how an applicator balances drift control and coverage. Applying larger droplets reduces drift potential, but will not prevent drift if applications are made improperly or under unfavorable environmental conditions.

A droplet size classification system describes the range of droplet sizes produced by spray nozzles. The American Society of Agricultural and Biological Engineers (ASABE) provide a Standard that describes droplet size spectrum categories defined by a number of reference nozzles (fine, coarse, etc.). Droplet spectra resulting from the use of a specific nozzle may also be described in terms of volume mean diameter (VMD). Coarser droplet size spectra have larger VMD's and lower drift potential.

CONTROLLING DROPLET SIZE - GROUND APPLICATION

- Nozzle Type - Select a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. The use of low-drift nozzles will reduce drift potential.
- Pressure - The lowest spray pressures recommended for the nozzle produce the largest droplets. Higher pressure reduces droplet size and does not improve canopy penetration. When higher flow rates are needed, using a higher-capacity nozzle instead of increasing pressure results in the coarsest droplet spectrum.
- Flow Rate/Orifice Size - Using the highest flow rate nozzles (largest orifice) that are consistent with pest control objectives reduces the potential for spray drift. Nozzles with higher rated flows produce coarser droplet spectra.

CONTROLLING DROPLET SIZE – AIRCRAFT

- Nozzle Type - Solid stream, or other low drift nozzles produce the coarsest droplet spectra.
- Number of Nozzles - Using the minimum number of nozzles with the highest flow rate that provide uniform coverage will produce a coarser droplet spectrum
- Nozzle Orientation - Orienting nozzles in a manner that minimizes the effects of air shear will produce the coarsest droplet spectra. For some nozzles such as solid stream, pointing the nozzles straight back parallel to the airstream will produce a coarser droplet spectrum than other orientations.
- Pressure – Selecting the pressure that produces the coarsest droplet spectrum for a particular nozzle and airspeed reduces spray drift potential. For some nozzle types such as solid streams, lower pressures can produce finer droplet spectra and increase drift potential

BOOM LENGTH (AIRCRAFT), AND APPLICATION HEIGHT

- Boom Length (aircraft) - Using shorter booms decreases drift potential. Boom lengths are expressed as a percentage of an aircraft’s wingspan or a helicopter’s rotor blade diameter. Shorter boom length and proper positioning can minimize drift caused by wingtip or rotor vortices.
- Application Height (aircraft) - Applications made at the lowest height that are consistent with pest control objectives and the safe operation of the aircraft will reduce the potential for spray drift.
- Application Height (ground) - Applications made at the lowest height consistent with pest control objectives, and that allow the applicator to keep the boom level with the application site and minimize bounce, will reduce the exposure of spray droplets to evaporation and wind, and reduce spray drift potential.

WIND

Drift potential is lowest when applications are made in light to gentle sustained winds (2-10 mph), which are blowing in a constant direction. Many factors, including droplet size and equipment type also determine drift potential at any given wind speed. AVOID GUSTY OR WINDLESS CONDITIONS.

Local terrain can also influence wind patterns. Every applicator is expected to be familiar with local wind patterns and how they affect spray drift.

TEMPERATURE AND HUMIDITY

Setting up equipment to produce larger droplets to compensate for droplet evaporation can reduce spray drift potential. Droplet evaporation is most severe when conditions are both hot and dry.

SURFACE TEMPERATURE INVERSIONS

Drift potential is high during a surface temperature inversion. Surface inversions restrict vertical air mixing, which may cause small suspended droplets to remain close to the ground and move laterally in a concentrated cloud. Surface inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Mist or fog may indicate the presence of an inversion in humid areas. Inversions may also be identified by producing smoke and observing its behavior. Smoke that remains close to the ground, or moves laterally in a concentrated cloud under low wind conditions indicates a surface inversion. Smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

SHIELDED SPRAYERS

Shielding the boom or individual nozzles can reduce the effects of wind. However, it is the responsibility of the applicator to verify that the shields are minimizing drift potential, and not interfering with uniform deposition of the product.

AIR ASSISTED (AIR BLAST) FIELD CROP SPRAYERS

Air assisted field crop sprayers carry droplets to the target via a downward directed air stream. Some may reduce the potential for drift, but if a sprayer is unsuitable for the application and/or set up improperly, high drift potential can result. It is the responsibility of the applicator to determine that a sprayer is suitable for the intended application, that it is configured properly, and that drift potential has been minimized.

Note: Air assisted field sprayers can affect product performance by affecting spray coverage and canopy penetration. Read the specific crop use and application equipment instructions to determine if an air assisted field crop sprayer can be used.

SENSITIVE AREAS

Making applications when there is a sustained wind moving away from adjacent sensitive areas (e.g., residential areas, bodies of water, known habitat for threatened or endangered species, non-target crops) is an effective way to minimize the effect of spray drift.

DRIFT CONTROL ADDITIVES

Using product compatible drift control additives can reduce drift potential. When a drift control additive is used, read and carefully observe cautionary statements and all other information on the additive’s label. If using an additive that increases viscosity, ensure that the nozzles and other application equipment will function properly with a viscous spray solution. Preferred drift control additives have been certified by the Chemical Producers and Distributors Association (CPDA).

STORAGE AND DISPOSAL

PESTICIDE STORAGE: Store product in original container only. Do not contaminate water, other pesticides, fertilizer, food or feed in storage.

PESTICIDE DISPOSAL: Do not contaminate water, food or feed by disposal. Waste resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

CONTAINER HANDLING: Refer to the Net Contents section of this product's labeling for the applicable "Nonrefillable Container" or "Refillable Container" designation.

Nonrefillable Rigid Plastic and Metal Containers (Capacity Equal to or Less Than 5 Gallons): Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then, for Plastic Containers, offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration. Do not burn, unless allowed by state and local ordinances. For Metal Containers, offer for recycling if available or reconditioning if appropriate, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

Nonrefillable Rigid Plastic and Metal Containers (Capacity Greater Than 5 Gallons): Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then for Plastic Containers, offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration. Do not burn, unless allowed by state and local ordinances. For Metal Containers, offer for recycling if available or reconditioning if appropriate, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

Nonrefillable Rigid Plastic and Metal Containers, e.g., Intermediate Bulk Containers [IBC] (Size or Shape Too Large to be Tipped, Rolled or Turned Upside Down): Nonrefillable container. Do not reuse or refill this container. Clean container promptly after emptying the contents from this container into application equipment or mix tank and before final disposal using the following pressure rinsing procedure. Insert a lance fitted with a suitable tank cleaning nozzle into the container and ensure that the water spray thoroughly covers the top, bottom and all sides inside the container. The nozzle manufacturer generally provides instructions for the appropriate spray pressure, spray duration and/or spray volume. If the manufacturer's instructions are not available, pressure rinse the container for at least 60 seconds using a minimum pressure of 30 PSI with a minimum rinse volume of 10% of the container volume. Drain, pour or pump rinsate into application equipment or rinsate collection system. Repeat this pressure rinsing procedure two more times. Then, for Plastic Containers, offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration. For Metal Containers, offer for recycling if available or reconditioning if appropriate or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

All Refillable Containers: Refillable container. Refilling Container: Refill this container with DuPont™ APROACH™ Fungicide containing Picoxystrobin only. Do not reuse this container for any other purpose. Cleaning before refilling is the responsibility of the refiller. Prior to refilling, inspect carefully for damage such as cracks, punctures, abrasions, worn out threads and closure devices. If damage is found, do not use container, contact DuPont at the number below for instructions. Check for leaks after refilling and before transporting. If leaks are found, do not reuse or transport container, contact DuPont at the number below for instructions. Disposing of Container: Do not reuse this container for any other purpose other than refilling (see preceding). Cleaning the container before final disposal is the responsibility of the person disposing of the container. To clean the container before final disposal, use the following pressure rinsing procedure. Insert a lance fitted with a suitable tank cleaning nozzle into the container and ensure that the water spray thoroughly covers the top, bottom and all sides inside the container. The nozzle manufacturer generally provides instructions for the appropriate spray pressure, spray duration and/or spray volume. If the manufacturer's instructions are not available, pressure rinse the container for at least 60 seconds using a minimum pressure of 30 PSI with a minimum rinse volume of 10% of the container volume. Drain, pour or pump rinsate into application equipment or rinsate collection system. Repeat this pressure rinsing procedure two more times. Then, for Plastic Containers, offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration. Do not burn, unless allowed by state and local ordinances. For Metal Containers, offer for recycling if available or reconditioning if appropriate, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

Do not transport if container is damaged or leaking. If the container is damaged, leaking or obsolete, or in the event of a major spill, fire or other emergency, contact DuPont at 1-800-441-3637, day or night.

NOTICE TO BUYER: Purchase of this material does not confer any rights under patents of countries outside of the United States.

The DuPont™ Oval Logo, DuPont™ and APROACH™ are trademarks or registered trademarks of E. I. du Pont de Nemours and Company.

D - 1541 112912

LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read this Limitation of Warranty and Liability Before Buying or Using This Product. If the Terms Are Not Acceptable, Return the Product at Once, Unopened, and the Purchase Price Will Be Refunded.

It is impossible to eliminate all risks associated with the use of this product. Such risks arise from weather conditions, soil factors, off target movement, unconventional farming techniques, presence of other materials, the manner of use or application, or other unknown factors, all of which are beyond the control of DuPont. These risks can cause: ineffectiveness of the product, crop injury, or injury to non-target crops or plants. WHEN YOU BUY OR USE THIS PRODUCT, YOU AGREE TO ACCEPT THESE RISKS.

DuPont warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for the purpose stated in the Directions for Use, subject to the inherent risks described above, when used in accordance with the Directions for Use under normal conditions.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, DUPONT MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR OF MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL DUPONT OR SELLER BE LIABLE FOR ANY INCIDENTAL, CONSEQUENTIAL OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. BUYER'S OR USER'S BARGAINED-FOR EXPECTATION IS CROP PROTECTION. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER AND THE EXCLUSIVE LIABILITY OF DUPONT OR SELLER, FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY OR CONTRACT, NEGLIGENCE, TORT OR STRICT LIABILITY), WHETHER FROM FAILURE TO PERFORM OR INJURY TO CROPS OR OTHER PLANTS, AND RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT, OR AT THE ELECTION OF DUPONT OR SELLER, THE REPLACEMENT OF THE PRODUCT.

To the extent consistent with applicable law that allows such requirement, DuPont or its Ag Retailer must have prompt notice of any claim so that an immediate inspection of buyer's or user's growing crops can be made. Buyer and all users shall promptly notify DuPont or a DuPont Ag Retailer of any claims, whether based on contract, negligence, strict liability, other tort or otherwise, or be barred from any remedy.

This Limitation of Warranty and Liability may not be amended by any oral or written agreement.

For product information call: 1-888-6-DUPONT [1-888-638-7668]

Internet address: www.cropprotection.dupont.com

© 2012 E. I. du Pont de Nemours and Company, Crop Protection, 1007 Market Street, Wilmington, DE 19898

All rights reserved.