

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, DC 20460



OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

July 29, 2014

Bonnie Bieber
E.I. du Pont de Nemours & Company
Crop Protection
P.O. Box 30
Newark, DE 19714

Subject: Label Amendment – Supplemental label for legume vegetables
Product Name: Dupont Prevathon Insect Control
EPA Registration Number: 352-844
Application Date: June 26, 2014
Decision Number: 492944

Dear Ms. Bieber:

The amended label referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act, as amended, is acceptable. A stamped copy of your labeling is enclosed for your records. You must submit one (1) copy of the final printed labeling before you release the product for shipment. Please note that the supplemental label expires on 1/17/2016 and must not be used or distributed after this date. If you have any questions, you may contact Jennifer Urbanski at 703-347-0156 or via email at urbanski.jennifer@epa.gov.

Sincerely,

A handwritten signature in black ink, appearing to read "Venus Eagle".

Venus Eagle, Product Manager 01
Insecticide-Rodenticide Branch
Registration Division (7505P)
Office of Pesticide Programs

Enclosure



**DuPont Crop
Protection**

SUPPLEMENTAL LABELING

DUPONT™ PREVATHON®
INSECT CONTROL WITH ACTIVE INGREDI-
ENT RYNAXYPYR® FOR USE ON LEGUME
VEGETABLES - DRIED SHELLED PEA AND
BEAN SUBGROUP AND FOLIAGE OF LEGUME
VEGETABLES (EXCEPT SOYBEAN)

DUPONT™ PREVATHON® INSECT CONTROL

EPA Reg. No. 352-844

This supplemental label expires on 1/17/2016 and must not be used or distributed after this date.

FOR FOLIAR INSECT CONTROL ON LEGUME VEGETABLES - DRIED SHELLED PEA AND BEAN SUBGROUP AND FOLIAGE OF LEGUME VEGETABLES (EXCEPT SOYBEAN)

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

This labeling must be in the possession of the user at the time of pesticide application. Read the label affixed to the container for PREVATHON® before applying. Use of PREVATHON® according to the labeling is subject to the use precautions and limitations imposed by the label affixed to the container for PREVATHON®.

See Federal Section 3 label accompanying product for full resistance management instructions.


APPLICATION

Apply at the specified rates when insect populations reach locally determined economic thresholds. Consult the cooperative extension service, professional consultants or other qualified authorities to determine appropriate threshold levels for treatment in your area. Apply follow-up treatments of PREVATHON®, as specified, to keep pest populations within threshold limits. Refer to the Resistance Management section of the Federal Section 3 label for further guidance on follow-up treatments. See individual crop sections of this label for specific minimum spray intervals. Use sufficient water to obtain thorough, uniform coverage.

IMPORTANT

Before using PREVATHON®, read and follow all applicable directions: restrictions; and precautions on the EPA-registered label. This supplemental label contains new or supplemental instructions for use of these products in combination which does not appear on the package label. Follow the instructions carefully.

Read the Limitation of Warranty and Liability on the Section 3 Federal product label before buying or using THIS product. If terms are not acceptable, return the unopened package at once to Seller for full refund of purchase price paid. Otherwise, use by Buyer or any other User constitutes acceptance of the terms of the Limitation of Warranty and Liability on the Section 3 Federal product label.



© 2014 E. I. du Pont de Nemours and Company, Crop Protection, 1007 N. Market St., Wilmington, Delaware 19898

Crop	Application Method	Target Pest	DUPONT™ PREVATHON® RATE		Last Application (Days to Harvest)	REI (Hours)
			Lb ai per acre	fluid ounces product per acre		
Legume vegetables, Dried shelled pea and bean (for soybean see separate soybean crop section below.) subgroup: Dried cultivars of bean (Lupins); bean (Phaseolus) (includes field bean, kidney bean, lima bean (dry), navy bean, pinto bean, tepary bean); bean (Vigna) includes adzuki bean, blackeyed pea, catjang, cowpea, crowder pea moth bean, mung bean, rice bean, southern pea, urd bean); broad bean (dry); chickpea; guar; lablab bean; lentil; pea (Pisum) (includes field pea); pigeon pea	FOLIAR	Corn earworm Beet armyworm European corn borer Fall armyworm	0.047 - 0.067	14.0 - 20.0	1	4
		Leafminers (larvae)*	0.097	29.0		
<p>USE RESTRICTIONS Make no more than 4 applications per acre per crop. Minimum interval between treatments in 3 days. Do not apply more than 60 fl. oz. PREVATHON® or 0.2 lbs a.i. of chlorantraniliprole containing products per acre per year.</p> <p>*Control of Liriomyza species except suppression only for L. huidabrensis and L. langei.</p>						
Foliage of Legume vegetables except soybean. Plant parts of any legume vegetable included in the legume vegetables that will be used as animal feed.	FOLIAR	Corn earworm Beet armyworm European corn borer Fall armyworm	0.047 - 0.067	14.0 - 20.0		
<p>USE RESTRICTIONS Make no more than 4 applications per acre per crop. Minimum interval between treatments in 3 days. Do not apply more than 60 fl. oz. PREVATHON or 0.2 lbs a.i. of chlorantraniliprole containing products per acre per year.</p>						