



U.S. ENVIRONMENTAL PROTECTION AGENCY

Office of Pesticide Programs  
Registration Division (7505P)  
1200 Pennsylvania Ave., N.W.  
Washington, D.C. 20460

EPA Reg. Number:

352-940

Date of Issuance:

9/21/21

NOTICE OF PESTICIDE:

Registration  
 Reregistration  
(under FIFRA, as amended)

Term of Issuance:

Unconditional

Name of Pesticide Product:

Lumiante™ FST

Name and Address of Registrant (include ZIP Code):

Samantha Miller  
Regulator Leader for CORTEVA™ Agriscience  
c/o E.I. du Pont de Nemours and Company  
9330 Zionsville Road  
Indianapolis, IN 46268

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is unconditionally registered in accordance with FIFRA section 3(c)(5) provided that you:

1. Submit and/or cite all data required for registration/reregistration/registration review of your product when the Agency requires all registrants of similar products to submit such data.

Signature of Approving Official:

Carmen J. Rodia, Jr., Acting Product Manager 22  
Fungicide Branch, Registration Division (7505P)

Date:

9/21/21

2. Make the following label changes before you release the product for shipment:
  - Revise the EPA Registration Number to read, “EPA Reg. No. 352-940.”
3. Submit one copy of the revised final printed label for the record before you release the product for shipment.

Should you wish to add/retain a reference to the company’s website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product’s label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA’s Office of Enforcement and Compliance.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6. Your release for shipment of the product constitutes acceptance of these conditions. A stamped copy of the label is enclosed for your records. Please also note that the record for this product currently contains the following CSFs:

- Basic CSF dated 4/26/2021

If you have any questions, please contact Elisha Graham via email at [Graham.Elisha@epa.gov](mailto:Graham.Elisha@epa.gov).

Enclosure

(Base Label)

ETHABOXAM	GROUP	22	FUNGICIDE
-----------	-------	----	-----------

[Bracketed text is optional]

[Bracketed *Italicized text is information for the reviewer*]

# Lumiante™ FST


**A FUNGICIDE SEED TREATMENT PRODUCT PROVIDING SYSTEMIC FUNGICIDE PROTECTION AGAINST SEED AND SEEDLING DISEASES CAUSED BY *PYTHIUM***

Active Ingredient	By Wt
Ethaboxam*.....	34.2%
Other Ingredients .....	65.8%
Total	100.0%

\*(RS)-N-( $\alpha$ -cyano-2-thenyl)-4-ethyl-2-(ethylamino)-1,3-thiazole-5-carboxamide

Lumiante™ FST contains 3.2 lb ethaboxam per gallon.

**KEEP OUT OF REACH OF CHILDREN  
CAUTION**



## PERSONAL PROTECTIVE EQUIPMENT (PPE)

**Workers loading product, treating seed and conducting maintenance and calibration of the treater must wear:** long-sleeved shirt, long pants, socks and shoes, and chemical-resistant gloves made of any waterproof material.

**Workers involved with handling treated seed (such as baggers, sewers and bag stackers) must wear:** long-sleeved shirt, long pants, socks and shoes.

**Workers who load and plant treated seed must wear:** long-sleeved shirt, long pants, socks and shoes and chemical-resistant gloves made of any waterproof material when handling treated seed.

Follow the manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

## USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

## ENVIRONMENTAL HAZARDS

This pesticide is toxic to aquatic invertebrates, oysters, and shrimp. Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash water or rinsate. Do not apply where runoff is likely

to occur. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Apply this product only as specified on label.

### **Groundwater Advisory**

This chemical has properties and characteristics associated with chemicals detected in groundwater. This chemical may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow.

### **Surface Water Advisory**

This product may impact surface water quality due to runoff of rain water. This is especially true for poorly draining soils and soils with shallow ground water. This product is classified as having high potential for reaching surface water via runoff for several days after application. A level, well maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams and springs will reduce the potential loading of ethaboxam from runoff water and sediment. Runoff of this product will be reduced by avoiding applications when rainfall or irrigation is expected to occur within 48 hours.

## **STORAGE AND DISPOSAL**

Do not contaminate water, food or feed by storage or disposal.

### **PESTICIDE STORAGE**

Keep pesticide in original container.

Store in a dry place away from excessive heat.

Do not store near food or feed.

Do not put pesticide into food or drink containers.

Not for use or storage around the home.

To close package, replace and tighten cap to form an airtight seal.

### **PESTICIDE DISPOSAL**

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

### **CONTAINER HANDLING**

[Use the following statement for containers equal to or less than 5 gallons]

[Nonrefillable container. Do not reuse or refill this container. Clean container promptly after emptying.

Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Add water – at least 2% of the container volume, and up to 1/3 of the volume of water needed to make the proper slurry composition with a maximum of 1/4 of the container volume – and recap. Shake for 30 seconds. Pour rinsate into a rinsate collection system for disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Offer for recycling, if available; otherwise dispose of in a sanitary landfill or other procedures allowed by State and local authorities.]

[Use the following statement for rigid nonrefillable containers greater than 5 gal or 50 lb]

[Nonrefillable container. Do not reuse or refill this container. Clean container promptly after emptying.

Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Add water – at least 2% of the container volume, and up to 1/3 of the volume of water needed to make the proper slurry composition with a maximum of 1/4 of the container volume. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 60 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into a rinsate collection system for disposal. Repeat this procedure two more times. Offer for recycling, if available; otherwise dispose of in a sanitary landfill or other procedures allowed by State and local authorities.]

[Use the following statement for all refillable container types]

[Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents into application equipment or mix tank. Add water – at least 2% of the container volume, and up to 1/3 of the volume of water needed to make the proper slurry composition with a maximum of 1/4 of the container volume. Replace and tighten closure. Agitate vigorously or recirculate the rinsate with a pump for at least 2 minutes ensuring that the rinsate rinses the walls of the container. Empty the rinsate into a rinsate collection system for disposal. Repeat this rinsing procedure two more times. Offer for recycling, if available; otherwise dispose of in a sanitary landfill or other procedures allowed by State and local authorities.]

EPA Reg No. 352-XXX

EPA Est. No. \_\_\_\_\_

®™Trademarks of Dow AgroSciences, DuPont or Pioneer and their affiliated companies or respective owners

Manufactured  
E. I. du Pont de Nemours and Company  
Chestnut Run Plaza  
974 Centre Road  
Wilmington, DE 19805

**NET CONTENTS** \_\_\_\_\_

(cover, shipping container)

ETHABOXAM	GROUP	22	FUNGICIDE
-----------	-------	----	-----------

[Bracketed text is optional]

[Bracketed *Italicized text is information for the reviewer*]

# Lumiante™ FST

## A FUNGICIDE SEED TREATMENT PRODUCT PROVIDING SYSTEMIC FUNGICIDE PROTECTION AGAINST SEED AND SEEDLING DISEASES CAUSED BY *PYTHIUM*

Active Ingredient	By Wt
Ethaboxam* .....	34.2%
Other Ingredients .....	65.8%
Total	100.0%

\*(*RS*)-*N*-( $\alpha$ -cyano-2-thenyl)-4-ethyl-2-(ethylamino)-1,3-thiazole-5-carboxamide

Lumiante™ FST contains 3.2 lb ethaboxam per gallon.

### KEEP OUT OF REACH OF CHILDREN CAUTION

#### SEE NEXT PAGE FOR PRECAUTIONARY STATEMENTS

EPA Reg No. 352-XXX

EPA Est. No. \_\_\_\_\_

®™Trademarks of Dow AgroSciences, DuPont or Pioneer and their affiliated companies or respective owners

Manufactured for  
E. I. du Pont de Nemours and Company  
Chestnut Run Plaza  
974 Centre Road  
Wilmington, DE 19805

NET CONTENTS \_\_\_\_\_

(page 1 through end)

## PRECAUTIONARY STATEMENTS

### CAUTION

#### PERSONAL PROTECTIVE EQUIPMENT (PPE)

**Workers loading product, treating seed and conducting maintenance and calibration of the treater must wear:** long-sleeved shirt, long pants, socks and shoes, and chemical-resistant gloves made of any waterproof material.

**Workers involved with handling treated seed (such as baggers, sewers and bag stackers) must wear:** long-sleeved shirt, long pants, socks and shoes.

**Workers who load and plant treated seed must wear:** long-sleeved shirt, long pants, socks and shoes and chemical-resistant gloves made of any waterproof material when handling treated seed.

Follow the manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

#### USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

#### ENVIRONMENTAL HAZARDS

This pesticide is toxic to aquatic invertebrates, oysters, and shrimp. Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash water or rinsate. Do not apply where runoff is likely to occur. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Apply this product only as specified on label.

#### Groundwater Advisory

This chemical has properties and characteristics associated with chemicals detected in groundwater. This chemical may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow.

#### Surface Water Advisory

This product may impact surface water quality due to runoff of rain water. This is especially true for poorly draining soils and soils with shallow ground water. This product is classified as having high potential for reaching surface water via runoff for several days after application. A level, well maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams and springs will reduce the potential loading of ethaboxam from runoff water and sediment. Runoff of this product will be reduced by avoiding applications when rainfall or irrigation is expected to occur within 48 hours.

#### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training,

decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard (WPS).

**DO NOT** enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

*Exception:* Once the seeds are planted in soil, the WPS, under certain circumstances, allows workers to enter the treated area without restriction if there will be no worker contact with the soil subsurface or treated seed. PPE required for early entry to treated areas that is permitted under the WPS and that involves contact with anything that has been treated, such as plants, soil, or water, is: long-sleeved shirt and long pants, socks and shoes, and chemical-resistant gloves made of any waterproof material.

### **STORAGE AND DISPOSAL**

Do not contaminate water, food, or feed by storage or disposal.

#### **PESTICIDE STORAGE: Keep pesticide in original container.**

Store in a dry place away from excessive heat.

Do not store near food or feed.

Do not put pesticide into food or drink containers.

Not for use or storage around the home.

To close package, replace and tighten cap to form an airtight seal.

#### **PESTICIDE DISPOSAL**

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

#### **CONTAINER HANDLING**

[Use the following statement for containers equal to or less than 5 gallons]

[Nonrefillable container. Do not reuse or refill this container. Clean container promptly after emptying.

Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Add water – at least 2% of the container volume, and up to 1/3 of the volume of water needed to make the proper slurry composition with a maximum of 1/4 of the container volume – and recap. Shake for 30 seconds. Pour rinsate into a rinsate collection system for disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Offer for recycling, if available; otherwise dispose of in a sanitary landfill or other procedures allowed by State and local authorities.]

[Use the following statement for rigid nonrefillable containers greater than 5 gal or 50 lb]

[Nonrefillable container. Do not reuse or refill this container. Clean container promptly after emptying.

Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Add water – at least 2% of the container volume, and up to 1/3 of the volume of water needed to make the proper slurry composition with a maximum of 1/4 of the container volume. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 60 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into a rinsate collection system for disposal. Repeat this procedure two more times. Offer for recycling, if available; otherwise dispose of in a sanitary landfill or other procedures allowed by State and local authorities.]

[Use the following statement for all refillable container types]

[Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents into application equipment or mix tank. Add water – at least 2% of the container volume, and up to 1/3 of the volume of water needed to make the proper slurry composition



with a maximum of 1/4 of the container volume. Replace and tighten closure. Agitate vigorously or recirculate the rinsate with a pump for at least 2 minutes ensuring that the rinsate rinses the walls of the container. Empty the rinsate into a rinsate collection system for disposal. Repeat this rinsing procedure two more times. Offer for recycling, if available; otherwise dispose of in a sanitary landfill or other procedures allowed by State and local authorities.]

## **DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

## **READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH PRECAUTIONARY STATEMENTS AND DIRECTIONS AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.**

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the State or Tribal agency responsible for pesticide regulation.

## **APPLICATION INSTRUCTIONS**

### **RESTRICTIONS**

Lumiante FST must be used only in accordance with use instructions on this label utilizing mechanical, slurry or mist-type seed treating equipment, provided that the equipment can be calibrated to accurately and uniformly apply the product to seed. This product is for both commercial and on-farm application. Do not apply this product in a hopper-box or planter-box at planting time. Always mix product thoroughly before use.

As stated by regulations specified in 40 CFR 153.155, all seed treated with Lumiante FST must be colored with an EPA-approved dye or colorant to visually distinguish seed is treated with a pesticide. It is mandatory that such dye or colorant be added into the slurry mixture containing Lumiante FST, prior to application. This regulation is necessary to prevent treated seed from accidental use as a food for man or feed for animals.

### **MODE OF ACTION**

Lumiante FST contains ethaboxam, which is a Group 22 fungicide as classified by the Fungicide Resistance Action Committee (FRAC). Its mode of action is the inhibition of cell division.

### **SEED QUALITY AND STORAGE OF TREATED SEED**

Seed treatments applied to low vigor or poor quality seed or to mechanically damaged seed may result in loss of seed germination or reduced seedling vigor. To verify seed safety, treat a small portion of seed with the desired slurry mixture (Lumiante FST alone or in combination with other commercial seed treatment products) prior to treatment of an entire seed lot. Utilize standard seed germination methodologies to determine and verify seed safety. Use proper storage conditions for treated seed. Due to seed quality and seed storage conditions beyond the control of DuPont, no claims are made to guarantee the germination of carry-over treated seed.

### **SEED BAG TAG**

The Federal Seed Act requires that bags containing treated seed be labeled or tagged with the following information:

- This seed has been treated with ethaboxam.
- Do not use for feed, food or oil processing.
- Excess seed may be used for ethanol production only if (1) by-products are not used for livestock feed and (2) no measurable residues of pesticide remain in ethanol by-products that are used in agronomic practices.

In addition, the following statements are required on containers of seed treated with Lumiante FST Fungicide:

- Store away from feeds and other foodstuffs.
- Wear long-sleeved shirt, long pants, shoes and socks, and chemical-resistant gloves made of any waterproof material when handling treated seed.
- Dispose of all excess treated seed. Left over treated seed may be double-sown around the headland or buried away from water sources in accordance with local requirements. Do not contaminate water bodies when disposing of planting equipment washwaters.
- Dispose of seed packaging in accordance with local requirements.
- Treated seeds exposed on soil surface may be hazardous to wildlife. Cover or collect treated seeds spilled during loading.
- Treated seed must be planted into the soil at a minimum depth of 0.5 inch.

### PRODUCT INFORMATION

Lumiante FST is a systemic fungicide seed protection product for use on corn. The active ingredient, ethaboxam, provides protection against Oomycetes including *Pythium* spp. that can cause seed decay, seedling dieback and root rot.

For best results, use Lumiante FST combined with other Oomycete-active seed treatment fungicides, such as metalaxyl or mefenoxam, to broaden the spectrum of activity. In addition, use with a broad-spectrum seed treatment fungicide(s) having activity against *Rhizoctonia solani* and other fungal pathogens inciting seed and seedling disease.

### APPLICATION DIRECTIONS

Accurate application of Lumiante FST to seed is required for best performance. Thoroughly mix Lumiante FST in the desired seed slurry mixture, prior to application to seed. **Pre-test tank mixtures with other seed treatment products to evaluate formulation compatibility and to ensure proper physical compatibility of products. Follow the most restrictive limitations and precautions on labels of all products used in a seed treatment mixture. Do not tank mix with any products which contain a prohibition on tank mixing.**

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

The following table provides use rates of Lumiante FST by labeled crop, with disease protection guidelines.

Corn (field, sweet and popcorn)	
Disease Protection Claims	Application Rate of Lumiante FST (fl oz/100 lb seed)

<i>Pythium</i> Seed Decay, Seedling Dieback and Root Rot	<p style="text-align: center;">0.2 to 0.3 (5 to 7.5 g ai/100 kg seed)</p> <p><b>Field Corn:</b> To deliver 5 g ai/100 kg seed, apply 0.1 fl oz per 80,000 seed (equals 0.0141 mg ai/seed) based on 1600 seed/lb count</p> <p><b>Sweet Corn:</b> To deliver 5 g ai/100 kg seed, apply 0.064 fl oz per 100,000 seed (equals 0.0072 mg ai/seed) based on 3150 seed/lb count</p> <p><b>Popcorn:</b> To deliver 5 g ai/100 kg seed, apply 0.066 fl oz per 100,000 seed (equals 0.0074 mg ai/seed) based on 3061 seed/lb count</p>
--	--

### LIMITATION OF WARRANTY AND LIABILITY

**NOTICE: Read this Limitation of Warranty and Liability Before Buying or Using This Product. If the Terms Are Not Acceptable, Return the Product at Once, Unopened, and the Purchase Price Will Be Refunded.**

DuPont will not be responsible for losses or damages resulting from use of this product in any manner not specifically published by DuPont and stated on this label or other labels or recommendations which conform with Section 2(ee) of the Federal Insecticide, Fungicide and Rodenticide Act. User assumes all risks associated with such non-specified use.

It is impossible to eliminate all risks associated with the use of this product. Such risks arise from weather conditions, soil factors, off target movement, unconventional farming techniques, presence of other materials, the manner of use or application, or other unknown factors, all of which are beyond the control of DuPont. These risks can cause: ineffectiveness of the product, crop injury, or injury to non-target crops or plants. **WHEN YOU BUY OR USE THIS PRODUCT, YOU AGREE TO ACCEPT THESE RISKS.**

DuPont warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for the purpose stated in the Directions for Use, subject to the inherent risks described above, when used in accordance with the Directions for Use under normal conditions.

**TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, DUPONT MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR OF MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL DUPONT OR SELLER BE LIABLE FOR ANY INCIDENTAL, CONSEQUENTIAL OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. BUYER'S OR USER'S BARGAINED-FOR EXPECTATION IS CROP PROTECTION. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER AND THE EXCLUSIVE LIABILITY OF DUPONT OR SELLER, FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY OR CONTRACT, NEGLIGENCE, TORT OR STRICT LIABILITY), WHETHER FROM FAILURE TO PERFORM OR INJURY TO CROPS OR OTHER PLANTS, AND RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT, OR AT THE ELECTION OF DUPONT OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

To the extent consistent with applicable law that allows such requirement, DuPont or its Ag Retailer must have prompt notice of any claim so that an immediate inspection of buyer's or user's growing crops can be made. Buyer and all users shall promptly notify DuPont or a DuPont Ag Retailer of any claims, whether based on contract, negligence, strict liability, other tort or otherwise, or be barred from any remedy.

#### COMMERCIAL TANK MIXES

**NOTICE:** Commercial tank mixing or use of this product with any other product which is not specifically and expressly authorized by the label shall be the exclusive risk of user, applicator and/or application advisor.

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

®™Trademarks of Dow AgroSciences, DuPont or Pioneer and their affiliated companies or respective owners