

ACCEPTED
 11/18/81
 UNDER THE FEDERAL INSECTICIDE
 FUNGICIDE AND ANTICIDE ACT
 FOR ECONOMIC POISON REGISTERED
 UNDER NO. 358-42

"Branched" crab grass usually requires two or three applications and if crab grass "yellows" and revives, it will be evident that additional applications are needed and these should begin, before seed head formation, and continued at 5-7 day intervals.

RATES FOR TEMPERATURES ABOVE 80°

super ARTOX required	add water to make*	to cover
1½ oz. (3 tablespoonfuls)	1 gal.	200 sq. ft.
3 oz. (6 tablespoonfuls)	2 gal.	400 sq. ft.
8 oz. (1 cup)	5 gal.	1000 sq. ft.
1 pint	10 gal.	2000 sq. ft.
2.75 gallons	220 gal.	1 acre

ANTIDOTE: If taken internally, induce vomiting and call a physician at once.

CAUTION: Do not get in eyes or on skin and if spilled on skin, wash off immediately with warm water and soap. Wash hands after using. Flush eyes with clear water. Avoid inhalation of spray mist. Avoid contamination of food, feedstuffs, streams or other water sources. Keep beyond the reach of children and domestic animals. Keep children and animals off treated areas until washed into soil. Harmful if swallowed. Do not apply to St. Augustine grass, Carpet grass or Centipede grass. Will temporarily injure bent grass and fescues. Do not reuse container. Wash thoroughly with water and detergent. Discard in a safe place.



PLEASANT VALLEY, NEW YORK 12569
 MADE IN U.S.A.

N&T
Super
ARTOX
LIQUID
CRAB GRASS KILLER

ACTIVE INGREDIENTS	8
INERT INGREDIENTS	84%
TOTAL	100

WARNING: Keep Beyond Reach of Children
 SEE SIDE PANEL FOR ADDITIONAL CAUTIONS.
 Contents: 1 pint (16 fluid ounces) EPA Reg. No. 358-42

DIRECTIONS

Do not use super ARTOX on lawns, ornamentals, or other plants susceptible to injury under unfavorable conditions. On newly seeded lawns do not apply until after the first mowing. Water lawns thoroughly previous to application of Artox and water evenly again for 3 days. Begin watering 3 days after treatment. Do not water for 3 days after watering is resumed.

WHEN TO APPLY: Early control should be practiced beginning in June when young crab grass plants are less than 2" tall. Repeat treatments for control of crab grass may be necessary to control low germination rates. Use Pressure Sprayer or garden hose with nozzle. Wet thoroughly the foliage of crab grass. Use sprayer or spraying equipment with clean water.

RATES FOR TEMPERATURES BELOW 80°

super ARTOX required	add water to make*	to cover
3 oz. (6 tablespoonfuls)	1 gal.	200 sq. ft.
6 oz. (12 tablespoonfuls)	2 gal.	400 sq. ft.
1 pint	5 gal.	1000 sq. ft.
1 quart	10 gal.	2000 sq. ft.
5½ gallons	220 gal.	1 acre

Application by Sprinkling can: Use 3 oz. per 200 sq. ft. of lawn. Use 6 oz. per 400 sq. ft. of lawn. Use 1 pint per 1000 sq. ft. of lawn. Use 1 quart per 2000 sq. ft. of lawn. Use 5½ gallons per acre.

ACCEPTED WITH COMMENTS