

RATES FOR JULY • AUGUST

Special super ARTOX required	add water to make*	TO COVER
1 1/2 oz. (3 tablespoonfuls)	1 gal.	200 sq. ft.
3 oz. (6 tablespoonfuls)	2 gal.	400 sq. ft.
8 oz. (1 cup)	5 gal.	1000 sq. ft.
1 pint	10 gal.	2000 sq. ft.
2 1/2-3 gallons	220 gal.	1 acre

**IMPORTANT:** Avoid spray drift to susceptible plants as this product may injure cotton, beans, peas, grapes, ornamentals and other plants. (Coarse sprays are less likely to drift.) Store away from seeds, fertilizers, insecticides and fungicides. Spray equipment should be rinsed several times with hot water and washing soda before using again for other spraying purposes. Do not reuse empty container. Wash thoroughly with water and detergent. Discard in a safe place. Do not contaminate water used for irrigation or domestic purposes.

**CAUTION:**

Avoid contact with skin, eyes or clothing. Avoid breathing vapor or spray mist. Wash hands after using. Avoid contamination of feed and foodstuffs and keep beyond the reach of children and domestic animals. **Poisonous if swallowed.** Keep children and animals off treated areas until washed into soil. Vapors Harmful.


**FIRST AID:** If taken internally induce vomiting and call a Physician at once.



Pleasant Valley, N. Y. 12569

Made in U.S.A.

EPA EST 358-NY-1



**SPECIAL**



**LIQUID**

**A.M.A. & 2,4-D**

**KILLS Lawn Weeds - Dandelion, Crab Grass, Chickweed, Plantain, Dallis Grass.**

**ACTIVE INGREDIENTS:**

Octylammonium Methane arsonate	8.00%
Dodecylammonium Methane arsonate	8.00%
*Octyl Ammonium Salt of 2,4-Dichlorophenoxyacetic Acid	5.44%
<b>INERT INGREDIENTS</b>	<b>78.56%</b>

**TOTAL 100.00%**

Total arsenic, expressed as metallic, all in water soluble form. 4.00%  
\*Equivalent to 2,4-Dichlorophenoxyacetic acid. 3.43%

**CAUTION: Keep Beyond Reach of Children**

SEE SIDE PANEL FOR ADDITIONAL CAUTIONS.

EPA Reg. No. 358-56 CONTENTS 1 pint

Do not use Special, Super ARTOX Liquid, when grasses are stunted or dormant due to drought or other unfavorable factors. On newly seeded lawns do not apply until after the 3rd mowing. Water lawn 1 day previous to applying and water lightly again for 3 days, beginning 2 days after treatment. If soil is moist from rain, no water is required.

**WHEN TO APPLY:** Early Spring—April and May for the control of Chickweed, Plantain and Dandelion. Early June for Crab Grass Seedlings, Dallis Grass and simultaneous control of common lawn weeds. July-August for branched Crab Grass. Since germination of many lawn weeds extends over several months, repeat applications, at 7-10 day intervals, may be necessary to control new growth. Delay re-seeding bare spots until two weeks after last application.

**HOW TO APPLY:** Follow chart rates for applicable months, using PRESSURE SPRAYER or GARDEN HOSE JAR SPRAYER. Wet thoroughly the foliage of all weeds to be controlled. If discoloration occurs with first application, on lawns predominantly Bent, Fescue or White Clover, reduce application rate by 50%. Do not use on St. Augustine, Carpet, Centipede, Bahia, Bent Putting Greens or Dichondra lawns. May possibly injure bentgrass and clover lawns.

RATES FOR APRIL • MAY • JUNE

Special super ARTOX required	add water to make	TO COVER
3 oz. (6 tablespoonfuls)	1 gal.	200 sq. ft.
6 oz. (12 tablespoonfuls)	2 gal.	400 sq. ft.
1 pint	5 gal.	1000 sq. ft.
1 quart	10 gal.	2000 sq. ft.
5 1/2 gallons	220 gal.	1 acre

**APPLICATION BY SPRINKLING CAN:** Use 1 1/2 ozs. (3 table-spoonfuls) in 1 gal. water (Imperial 4/5 gal.) to cover 100 sq. ft. \*Use 4 5 Imperial gals. for each U. S. Gal. water specified.