



Rhodia

CHEMICALS FOR AGRICULTURE AND INDUSTRY

PRODUCT NO. 402

NET WEIGHT 100 POUNDS

ATLACIDE® WITH 2,4-D

A Chlorate Type Weed Killer KILLS WEEDS AND GRASSES

ACTIVE INGREDIENTS:

Sodium chlorate (NaClO₂) 58.00%
2,4-dichlorophenoxyacetic acid 10.00%

INERT INGREDIENTS 41.40%

CAUTION: Keep Out of Reach of Children

CAUTION: Harmful if swallowed. Strong Oxidant. Contact with combustible material may cause fire. Clothing and vegetation contaminated with chlorate or its solutions are dangerously flammable. After application remove clothing and wash thoroughly with water. Store separately from flammable materials. Keep away from fire. Avoid drift to susceptible plants as this product may injure cotton, beans, peas, other vegetables, legumes and ornamentals (coarse sprays are less likely to drift). Dusting or spraying equipment in which 2,4-D has been used should be thoroughly cleaned with a suitable chemical before being used for other purposes. Do not transport or store with fertilizers, insecticides, fungicides or seeds. Do not contaminate any body of water by direct application, cleaning of equipment or disposal of wastes. Do not permit livestock to graze on treated areas.

Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

GENERAL RECOMMENDATIONS: For control of weeds and grasses along fence rows, railroad rights-of-way and other similar weed infested areas. Use 2 to 3 pounds per square rod, applied dry or as a spray (¾ to 3 pounds per 100 square feet). It is important not to plow, cultivate, burn or pasture treated areas for several months after treatment. Do not apply during hot, dry weather.

HOW TO APPLY: As a Spray—Dissolve 1 to 1½ pounds per gallon of water. Apply uniformly, wetting entire plant. Use pressure sprayer with coarse spray to obtain good coverage. Do not cultivate, burn, or pasture before treatment. Dry—Use undiluted and spread evenly by hand or mechanical spreader. Top growth should be disced in, or mowed and removed, before dry treatment is made.

WHEN TO APPLY: In areas of light winter rainfall (8 to 20 inches), apply in fall or early winter. In areas of heavy winter rainfall (20 inches or more), apply in the spring.

Local conditions may affect the use of this chemical. Consult State Agricultural Extension or Experiment Station weed specialists for specific recommendations for local weed problems.

USDA Reg. No. 359-5

Do NOT Store with Feed and Foodstuffs

Part. 1M-1-128

