

PRODUCT NO.

808

ACCEPTED
8-30-68

359-42

NET WEIGHT
POUNDS

CHIPMAN CALCIUM ARSENATE

And Other Calcium Compounds

For control of certain insects of vegetables as given in the directions.

ACTIVE INGREDIENT:

Tricalcium arsenate

75.00%

INERT INGREDIENTS

30.00%

Total Arsenic expressed as metallic

26.5%

Arsenic in water soluble form, expressed as metallic.

Not more than

0.15%

Density not less than 80 nor more than 100 cu. in. per pound

POISON-DANGER

Keep Out of Reach of Children

ANTIDOTE

Give 100 cc. of 1% solution of sodium bicarbonate in a glass of warm water. Repeat every 15 minutes until the patient is clear. Then give 2 tablespoonfuls of 1% solution of sodium bicarbonate in water and plenty of milk. If the patient is unconscious, give 100 cc. of 1% solution of sodium bicarbonate in water. **CALL A PHYSICIAN**

WARNING: Poisonous if Swallowed. Avoid contact with skin, eyes or clothing. Avoid breathing dust or spray mist. Keep out of reach of children and domestic animals. Do not store with feeds and foodstuffs. Do not reuse container. Destroy by burying in a safe place. Do not burn.

This product is toxic to wildlife. Wildlife feeding on treated areas may be harmed.

CAUTIONS: Do not use in combination with soap or fish oil. Avoid spray or dust drift that may cause arsenical burn on sensitive crops such as soybeans, rice, peaches and pecans, flowers and ornamentals, or contamination of forage crops. Do not use near bee hives, nor on blossoms that bees may work at a later time. Do not use in fields where rice may be planted, as a residue in the soil may be injurious.

DIRECTIONS

VEGETABLES Tomatoes, peppers and cabbage. Colorado Potato Beetle, flea beetle, tomato hornworm and fruit worm, larvae (caterpillars), diamond-back moth, imported cabbage butterfly and cabbage looper. Make first application as soon as insects threaten damage. Repeat applications as needed for control, but do not use more than 25 pounds per acre on cabbage, or not more than 12 pounds on peppers and tomatoes. Do not apply to cabbage after edible parts begin to form. Remove excess residue at harvest on peppers and tomatoes.

AS A DUST: Mix 1 part calcium arsenate with 3 parts of hydrated lime or fungicide mixture where this is also used. Apply 20 to 40 lbs. per acre of this mixture. Apply dust preferably when foliage is wet with dew and apply to obtain uniform coverage.

FOR SPRAY APPLICATION: Use 4 lbs. per acre applied with lime in sufficient water to give uniform coverage. Unless applied with lime sulfur or bordeaux mixture, add twice as much hydrated lime as calcium arsenate in the spray.

Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

U.S.D.A. Reg.
No. 359-42

N.B. 75M-1-88

Atlanta, Ga.
Bound Brook, N. J.
Chicago, Ill.
N. Kansas City, Mo.

Manufactured by

RHODIA INC.
CHIPMAN DIVISION
NEW BRUNSWICK, NEW JERSEY

Palo Alto, Calif.
Pasadena, Tex.
Portland, Ore.
St. Paul, Minn.

