

SUPPLEMENTAL LABELING

EPA Reg. No. 359-170

Product Number
2280

359-170
CHIPTOX[®]

For selective control of certain weeds in GRAIN SORGHUM

Chiptox used in grain sorghum will control weeds such as:

Pigweed
Lambsquarters
Shepherdspurse
Purslane
Narrowleaf Goosefoot

See label for other weeds controlled.

Apply when sorghum is 6 to 12 inches tall but before the boot stage of development. Use 3 pints in 10 to 20 gallons of water per acre. For best results, apply before weeds reach a height of 5 inches or branches of weeds exceed a length of 5 inches. Avoid cultivation after treatment while sorghum is brittle.

Hybrids vary in tolerance to MCPA. Hybrids of Wheatland parentage may be particularly susceptible. Spray only varieties known to be tolerant. Consult seed company or your Agricultural Experiment Station or Extension Service Weed Specialist for guidance.

Do not graze dairy animals on treated areas until 7 days after treatment.
Do not graze meat animals on treated areas within 7 days of slaughter.

NOTICE: Before using Chiptox, read and carefully observe the cautionary statements and all other information appearing on the product label.

This supplemental label contains directions for use for this product which may not appear on the package label. Follow directions carefully.

PL-1076

Manufactured by
RHODIA INC.
AGRICULTURAL DIVISION
MONMOUTH JUNCTION, NEW JERSEY 08852

