PRODUCT NO.
**2280**

# CHIPTOX

NET CONTENTS
GALLONS

## HERBICIDE

**For selective control of certain weeds in crops, grassland, turf, and in non-crop areas such as roadsides, fence rows, and rights-of-way.**

ACTIVE INGREDIENT
[illegible]

INERT INGREDIENTS
[illegible]

## CAUTION: Keep Out of Reach of Children

**CAUTION:** Avoid contact [illegible] to desirable plants as this product may [illegible] ornamentals. Coarse sprays are less likely to drift. Thoroughly clean spray equipment with [illegible] at the chemical cleaner before using it for other purposes. Avoid contamination of water [illegible] container by burying or burning.

Do not use or [illegible]

Do not transport with [illegible]

Keep out of lakes, streams and ponds. [illegible]

**GENERAL INFORMATION** Chiptox [illegible] research indicated that Chiptox [illegible]

**SELECTIVE SPRAYING: Wheat, Oats, Barley, and Rye** [illegible] 1 to 2 pints per acre [illegible] Stage. Do not [illegible]

**SMALL GRAINS Underseeded with Alfalfa, Birdsfoot Trefoil, Lespedeza, Red and White Clover** [illegible] weeds apply [illegible] after cereals are [illegible] stage. Do not [illegible] pressure as this may damage [illegible] with other [illegible] legumes if any thinning [illegible]

**GRAIN SORGHUM:** [illegible] Shepherdspurse, Pigweed, Smartweed [illegible]

Apply when sorghum is [illegible] 10 to 12 gallons water per acre [illegible] Do not [illegible]

Do not [illegible]

**RICE** [illegible]

**In California-** [illegible]

**FLAX** [illegible] 1 pint per acre in [illegible] gallons water. Apply after flax [illegible] at least [illegible] tall and before flax [illegible] bud. Treatment [illegible] Flax Variety Resistant Weeds [illegible] spraying [illegible] Chiptox per acre to [illegible] product by [illegible] ICA [illegible] susceptible grass weeds and [illegible] grass [illegible] has [illegible] with [illegible] consult [illegible]

**PEAS** [illegible] Chiptox per acre [illegible] gallons water per acre [illegible] after they emerge at [illegible] Do not apply after [illegible] Chiptox [illegible] Before [illegible] Chiptox [illegible] Carefully [illegible] differences in these [illegible]

**ESTABLISHED GRASSLANDS & PASTURES** [illegible] pints per acre in sufficient water [illegible] Canada [illegible] Spring [illegible] and [illegible] Other [illegible]

**ESTABLISHED LAWNS AND TURFS** [illegible] 3 to 5 gallons water by [illegible] 2 to 12 gallons water. Apply when [illegible] actively growing [illegible]

**GRASSES GROWN FOR SEED** [illegible] after [illegible]

**NON CROP SPRAYING: Canada Thistle, White Top, Meadow Buttercup, Field Bindweed (Morning glory), and Texas Blueweed** [illegible]

### CONTROLS THESE WEEDS

**SUSCEPTIBLE (Most Areas)**

[illegible]

**LESS SUSCEPTIBLE**

[illegible] Tall [illegible] Purslane, Silver leaf [illegible]

**IN RICE -CALIFORNIA**

[illegible]

LOT No.

**Do Not Store with Foodstuffs**

EPA Reg No 359-170
EPA Est 359-MO-1
EPA Est 359-OR-1

FL-1277


Manufactured for
**RHODIA INC.**
**AGRICULTURAL DIVISION**
MONMOUTH JUNCTION, NEW JERSEY 08852


Rhodia
A SUBSIDIARY OF RHÔNE-POULENC