

PRODUCT NO.

510

# CHIPMAN® NO BUNT LIQUID (SEED DISINFECTANT)

NET CONTENTS  
5 GALLONS

ACCEPTED  
3-27-69  
359-385  
ATTACHED COMMENTS

No Bunt Liquid contains hexachlorobenzene (HCB), a non-mercurial seed treating disinfectant specially formulated for use on wheat and sorghum in slurry and automatic and mist-type treaters. Controls Bunt or Stinking Smut of Wheat, Loose and Covered Kernel Smut of Sorghums. (1 gallon will treat 153 bushels of wheat or 196 bushels of sorghum in slurry treaters.)

ACTIVE INGREDIENT:

Hexachlorobenzene\* ..... 33.0%  
INERT INGREDIENTS ..... 67.0%

\*Equivalent to 4 pounds of hexachlorobenzene per gallon.

DIRECTIONS FOR USE

**GENERAL—MIXING:** For equipment without an agitator, always mix required amount of No Bunt Liquid with an equal volume of water before final dilution.

**FOR SLURRY TREATERS:** Add to water in treater at rates given below with agitator running.

	Amount of Liquid No Bunt	Amount of Water	Total Slurry	Slurry Cup	Dump Wt.
WHEAT	15 pts.	8 gal. 1 pt.	10 gal.	23 cc	10.5 lbs.
SORGHUM	12 pts.	8 gal. 4 pts.	10 gal.	23 cc	9.5 lbs.

### CAUTION: Keep Out of Reach of Children

Avoid contact with skin and eyes. Reasonable care should be exercised in the use of this product. Avoid inhalation. Wash thoroughly after handling.

Avoid contamination of feed and foodstuffs. Treated seed should not be used for feed, food or oil purposes.

Do not store at temperatures below 32°F.

**NOTE:** No Bunt Liquid is recommended only for treatment of seed wheat and sorghum for control of the diseases indicated above.

Local conditions may affect the use of this chemical. Consult State agricultural extension specialists for specific recommendations and for information on possible lower dosages.

Label treated seed as follows:

"This seed treated with hexachlorobenzene, NO BUNT LIQUID. Do not use for food or feed."

Use the correct slurry cup, bucket or rotor and correct seed gate (depending on the type of machine and seed used).

**FOR ON THE FARM TREATMENT:** Add 4 pts. of No Bunt Liquid to 2 gallons water. Mix thoroughly. Use 1/3 pint of this mixture to each bushel of seed, wheat or sorghum. Apply in a drum or revolving barrel or similar device to obtain uniform coverage.

**FOR AUTOMATIC AND OTHER READY MIX AND MIST TYPE TREATERS:** Make up treating strength by first diluting liquid No Bunt with equal volume of water, mix thoroughly and set machine to apply 2 fl. oz. per bushel for wheat, and 1 1/2 fl. oz. per bushel for sorghum. Keep well agitated.

Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

U.S.D.A. Reg.  
No. 359-385

Post BC 1 29

Atlanta, Ga.  
Bound Brook, N. J.  
Chicago, Ill.  
St. Paul, Minn.  
N. Kansas City, Mo.

Manufactured by  
**RHODIA INC.**  
CHIPMAN DIVISION  
NEW BRUNSWICK, N. J.

Palo Alto, Calif.  
Pasadena, Tex.  
Portland Ore.  
Billings, Mont.

