

PRODUCT NO.

404

BOREA K-10

(PELLETIZED)

NET WEIGHT
50 POUNDS

A Semi-Soil Sterilant Type Weed Killer

ACCEPTED
5-28-71
359-492

For use on railroads, industrial sites and similar non-crop areas to kill weeds and grasses where lasting residual effect is desired.

ACTIVE INGREDIENTS:

Diuron [3-(3,4-dichlorophenyl)-1,1-dimethylurea*]	8.0%
Sodium metaborate (Na ₂ B ₄ O ₇ ·5H ₂ O)**	30.0%
INERT INGREDIENTS	42.0%

*Equivalent to 10% "Kormex"

**Equivalent to 4.88% boron expressed as elemental boron.

"Kormex" a trademark of E. I. duPont de Nemours & Co., Inc.

U. S. Patent Nos. 2,655,445 and 2,886,425

CAUTION: Keep Out of Reach of Children

CAUTION: May cause irritation of eyes, nose and throat and the skin of some individuals. Avoid contact with skin, eyes and clothing. Avoid handling with cuts or abrasions on hands and wear gloves. Avoid breathing dust. Wash clothing and equipment after application. Keep away from children and domestic animals. Do not contaminate water intended for irrigation or domestic purposes. Do not use same equipment for applying insecticides, fungicides or fertilizers, as plant injury may result. Avoid contamination of feed and foodstuffs.

Do not reuse empty container. Destroy it by perforating or crushing. Bury or discard in a safe place.

PRECAUTIONS: Apply only as recommended. The chemical should be used only where unproductive soil is not objectionable. Do not apply on or near turf, valuable trees or other plants or on areas into which their roots may extend. Do not apply in locations where the chemical may be washed or moved into contact with their roots.

GENERAL RECOMMENDATIONS: May be used on non-cropland areas such as railroad rights-of-way, along pipe lines, around telephone and power poles, and other such areas where total plant kill is desired.

RECOMMENDATIONS: For general weed control, spread evenly by hand or mechanical spreader as follows:

For initial Application in Southern areas with long growing season, apply 400 to 800 lbs. per acre (1 to 2 lbs. per 100 sq. ft.).

In Northern areas with shorter growing season, apply 200 to 800 lbs. per acre (1/2 to 2 lbs. per 100 sq. ft.).

For Follow Up Treatments: Northern and Southern areas — apply 200 to 600 lbs. per acre (1/2 to 1 1/2 lb. per 100 sq. ft.)

The amount to use will depend on the type of weeds present, the degree of control and duration of effect required. These depend on soil type, rainfall and other conditions.

NOTE: Borea K-10 should be used wherever annual rainfall is in excess of 30 inches

WHEN TO APPLY: May be applied at any time of year, except when ground is frozen, as long as adequate moisture is supplied, either by rainfall or irrigation. Best results are obtained on annual and perennial weeds in high rainfall areas when applied just before weed growth begins. To avoid fire hazard from dead and dry vegetation, apply when weeds and grass are small.

Local conditions may affect the use of this chemical. Consult state agricultural extension or experiment station weed specialist for specific recommendations for local weed problems.

Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

DO NOT Store with Feed, Foodstuffs or Fertilizer.

Store in a Cool, Dry Place.

BEST DOCUMENT AVAILABLE

USDA Reg. No.
359-492

Atlanta, Ga.
Bound Brook, N. J.
Hinsdale, Ill.
N. Kansas City, Mo.

Manufactured by

RHODIA INC.
CHIPMAN DIVISION
NEW BRUNSWICK, NEW JERSEY

Fresno, Calif.
Pasadena, Tex.
Portland, Ore.
St. Paul, Minn.



PRODUCT NO.

404

BOREA

(PELLET)

A Semi-Soil Sterilant

For use on railroads, industrial sites and similar
where lasting residual

ACTIVE INGREDIENTS:

Diuron [3,3,4-dichlorophenyl-1,1-dimethylurea*]	8.0%
Sodium metaborate (Na ₂ B ₄ O ₇ ·5H ₂ O)**	30.0%
INERT INGREDIENTS	42.0%

*Equivalent to 10% "Karmex"
 **Equivalent to 4.88% boron expressed as elemental boron.
 "Karmex" a trademark of E. I. duPont deNemours & Co., Inc.
 U. S. Patent Nos. 2,655,445 and 2,886,425

CAUTION: Keep Out of Reach of Children

CAUTION: May cause irritation of eyes, nose and throat and the skin of some individuals. Avoid contact with skin, eyes and clothing. Avoid handling with cuts or abrasions on hands and wear gloves. Avoid breathing dust. Wash clothing and equipment after application. Keep away from children and domestic animals. Do not contaminate water intended for irrigation or domestic purposes. Do not use same equipment for applying insecticides, fungicides or fertilizers, as plant injury may result. Avoid contamination of feed and foodstuffs.

Do not reuse empty container. Destroy it by perforating or crushing. Bury or discard in a safe place.

PRECAUTIONS: Apply only as recommended. The chemical should be used only where unproductive soil is not objectionable. Do not apply on or near turf, valuable trees or other plants or on areas into which their roots may extend. Do not apply in locations where the chemical may be washed or moved into contact with their roots.

GENERAL RECOMMENDATIONS: May be used on non-cropland areas such as railroad rights-of-way, along pipe lines, around telephone and power poles, and other such areas where total plant kill is desired.

USDA Reg. No.
359-492

PAS. 1M-1-41

Atlanta Ga
 Bound Brook N J
 Hinsdale Ill
 N Kansas City Mo

Manufactu

RHODIA
CHIPMAN
 NEW BRUNSWICK